Imperial College London
Department of Earth Science and Engineering
MSc Environmental Data Science and Machine Learning

Comparison of Direct and Cascaded High-Resolution Spatial Downscaling of Wildfire Data

Author:
Tsuyu Huang
tsuyu.huang25@imperial.ac.uk

Supervisors:
Jorge Veiras (jorge@artiocarbon.com)
Prof. Anna Korre (a.korre@imperial.ac.uk)

Code Repository:
https://github.com/ese-ada-lovelace-2025/irp-th25

Company:
Artio
www.artiocarbon.com

July 13, 2026

Abstract

Wildfires are destructive and a rapidly evolving hazard. With climate change, they become increasingly severe and frequent. This increases the need for a higher-resolution wildfire data for better monitoring and risk assessment. Geostationary satellites (GEO) have high-temporal resolution, but lacks the resolution needed for a detailed wildfire risk assessment. While low orbit satellites (LEO) provide better spatial resolution, but low temporal resolution to monitor rapid wildfire spread. Existing downscaling models have demonstrated their effectiveness in producing high-resolution remote sensing data. But many still suffers from artefacts, blurriness, and hallucinations. This project aims to achieve a high-resolution spatial downscaling of wildfire data by integrating additional physical covariates to the model, such as topography and landcover to mitigate physically implausible outputs. The project compares direct (2km to 30m) and cascaded (2km to 375m to 30m) downscaling pipelines and their effectiveness in producing high-fidelity, high-resolution wildfire data. The project compares diffusion (EDiffSR), attention U-Net (HAUNet), and transformer-based (ESTNet) downscaling models.

Contents

1 Introduction **3**

2 Data and Materials **3**
2.1 Input Datasets 3
2.2 Target Fire Data 3

3 Methods **4**
3.1 HAUNet (Hybrid Attention U-Net) 4

4 Results **5**

5 Discussions **5**

6 Conclusions **5**

1 Introduction

In recent years, due to rising popularity in Net-Zero initiatives, carbon investors are increasingly funding carbon reduction and removal projects, such as reforestation (Lambert and Turner, 2024). However, these carbon projects are at risk because wildfires can destroy and release the carbon back into the atmosphere. With climate change, wildfires are becoming increasingly severe and frequent (Jones et al., 2020). This causes economic losses for the insurance industry. For instance, The January 2025 Los Angeles County wildfires set an all-time record, causing $40 billion in insured losses alone from a single event (Dwyer and Nam, 2026). This project, conducted with Artio, aims to address this gap. By providing highly accurate risk assessment models, this research seeks to give better confidence and enable better risk assessment.

High-spatial and high-temporal resolution wildfire data is highly desired in risk assessment. However, high-temporal resolution data obtained from geostationary satellites (GEO) have limited spatial resolution, which makes them difficult to use for accurate risk assessment. While low orbit satellites (LEO) offers high-spatial resolution, but low-temporal resolution.

To bridge this gap, existing methods uses Deep Learning (DL) models for downscaling due to their ability in recognising non-linear spatiotemporal patterns. However, the resolution gap between GEO and LEO is significant (2km - 30m). This increases the risk of producing physically implausible or unrealistic outputs, as a single low-resolution (LR) pixel can correspond to different high-resolution (HR) details (Z. Wang et al., 2020). In this project, to mitigate this limitation, the model integrates additional physical information, such as HR topography and landcover to help guide the downscaling process. To further improve downscaling, the project aims to investigate whether cascaded downscaling (2km to 375m to 30m) produces more accurate and better results compared to direct downscaling (2km to 30m). By breaking up the downscaling process, the model trains on a smaller resolution gap, aiming to mitigate physically implausible outputs. This project will compare the cascaded and direct pipeline performance of different downscaling architectures, including hybrid attention U-Net, a transformer-based model, and a diffusion-based model.

2 Data and Materials

2.1 Input Datasets

- GOES-17 channel 7 (IR shortwave) of Level 1B (L1B) Radiances product (ABI-L1B-Rad). Has spatial resolution of 2km at every 5 mins. This channel is selected because it is commonly used to identify hot regions (Badhan et al., 2024). This will be the main coarse input.
- Copernicus DEM GLO-30: Global 30m Digital Elevation Model with 30m resolution (European Space Agency and Airbus, 2022). DEM band will be used.
- ESA WorldCover 10m v100 landcover dataset with 10m resolution. This dataset have 11 land cover classes, including tree cover, shrubland, grassland, and permanent water bodies (Zanaga et al., 2021).

2.2 Target Fire Data

- Landsat-8 active fire dataset includes data for fire scenes taken from August and September 2020 with dimensions 256x256 at 30m resolution (de Almeida Pereira et al., 2021). This will be the final target HR target data. Fire-sensitive SWIR and thermal bands will be used.

- VIIRS (S-NPP) Band Active Fire with 375 m spatial resolution (Earth Science Data Systems, 2025) will be used as the target data for the 2km to 375m stage downscaling. The I-4 channel brightness temperature (3.55-3.93 µm) band will be used.

3 Methods

3.1 HAUNet (Hybrid Attention U-Net)

HAUNet is a hybrid attention U-Net model (J. Wang et al., 2023) that uses two variants of attention blocks:

- **CEM**: Applies channel attention.
- **S_CEM**: Additionally applies spatial attention in parallel.

The model uses cross-scale attention module as skip connections across different scale outputs in the encoder, to capture dependency between different scales. The overall framework of wildfire-adapted HAUNet is shown in Figure 1. The original HAUNet S_CEM does full spatial attention, which is $O((H \cdot W)^2)$ and will run into OOM crashes. Hence the input GOES and covariates are sampled to a smaller image before feeding into the model. But this loses covariates' high-resolution information, so an additional covariate pyramid encoder was added form the original model to run in parallel. This performs windowed spatial attention on the covariates at fine resolution. The output of the covariate encoders are fused at different scales during refinement.




Figure 1: Architecture overview of the wildfire-adapted HAUNet. The full-resolution stream processes native covariates, which are progressively fused via refinement blocks with the low-resolution stack after CIM.

4 Results

5 Discussions

6 Conclusions

References

Badhan, M., Shamsaei, K., Ebrahimian, H., Bebis, G., Lareau, N. P., & Rowell, E. (2024). Deep Learning Approach to Improve Spatial Resolution of GOES-17 Wildfire Boundaries Using VIIRS Satellite Data. *Remote Sensing*, *16*(4), 715. https://doi.org/10.3390/rs16040715

de Almeida Pereira, G. H., Fusioka, A. M., Nassu, B. T., & Minetto, R. (2021). Active fire detection in Landsat-8 imagery: A large-scale dataset and a deep-learning study. *ISPRS Journal of Photogrammetry and Remote Sensing*, *178*, 171–186. https://doi.org/10.1016/j.isprsjprs.2021.06.002

Dwyer, O., & Nam, H. W. (2026, March 31). *How wildfires and storms drove insurance losses in 2025 – in three charts*. Carbon Brief. Retrieved June 10, 2026, from https://www.carbonbrief.org/how-wildfires-and-storms-drove-insurance-losses-in-2025-in-three-charts/

Earth Science Data Systems, N. A. S. A. (2025, May 27). *VIIRS (S-NPP) I Band 375 m Active Fire Product NRT (Vector data) | NASA Earthdata*. Earth Science Data Systems, NASA. Retrieved June 8, 2026, from https://www.earthdata.nasa.gov/data/catalog/lancemodis-vnp14imgtdl-nrt-2

European Space Agency & Airbus. (2022). *Copernicus DEM*. https://doi.org/10.5270/ESA-c5d3d65

Jones, M. W., Smith, A. J. P., Betts, R., Canadell, J. G., Prentice, I. C., & Quéré, C. L. (2020). Climate Change Increases the Risk of Wildfires.

Lambert, J., & Turner, G. (2024, November 14). *Investment Trends and Outcomes in the Global Carbon-Credit Market-2024 | MSCI*. Retrieved June 5, 2026, from https://www.msci.com/research-and-insights/paper/investment-trends-and-outcomes-in-the-global-carbon-credit-market-2024

Wang, J., Wang, B., Wang, X., Zhao, Y., & Long, T. (2023). Hybrid Attention-Based U-Shaped Network for Remote Sensing Image Super-Resolution. *IEEE Transactions on Geoscience and Remote Sensing*, *61*, 1–15. https://doi.org/10.1109/TGRS.2023.3283769

Wang, Z., Chen, J., & Hoi, S. C. H. (2020, February 8). *Deep Learning for Image Super-resolution: A Survey*. arXiv: 1902.06068 [cs.CV]. https://doi.org/10.48550/arXiv.1902.06068

Zanaga, D., Van De Kerchove, R., De Keersmaecker, W., Souverijns, N., Brockmann, C., Quast, R., Wevers, J., Grosu, A., Paccini, A., Vergnaud, S., Cartus, O., Santoro, M., Fritz, S., Georgieva, I., Lesiv, M., Carter, S., Herold, M., Li, L., Tsendbazar, N.-E., ... Arino, O. (2021, October 20). *ESA WorldCover 10 m 2020 v100*. Zenodo. https://doi.org/10.5281/zenodo.5571936