IRP Tsuyu

Project Plan 12th June.

The main focus will be increasing the spatial resolution of wildfire satellite images. Step 1 of the big project. If we have time we can go to step 2.

Paper to follow:

https://www.mdpi.com/2072-4292/16/4/715

https://www.sciencedirect.com/science/article/abs/pii/S092427162100160X?via%3Dihub
(30m spatial resolution)

1. GOES vs **MTG**. If it's in Google earth engine it's easier to collect data. What bands are we using? Etc
2. Sentinel-2(20m) vs Landsat(30m) what do we want?
3. Cascade approach or direct approach, or both and compare
4.

Coarse Input Data: GOES vs MTG

	GOES	**MTG**
Spatial Res	2km	Visible and NIR: 1km IR: 2km (Active Fire Monitoring EO:EUM:DAT:0682)
Temporal Res	10 mins	Every 10 mins
Bands	3.9μm	FCI channel IR-3.8μm
Spatial coverage	The Americas (North and South)	Lat: -67.5° to 67.5°, Lon: -67.5° to 67.5° (EU, Africa, Atlantic Ocean, parts of Middle East and South

		America?)
Temporal extent	GOES-16: May 2017 - Now GOES-17: 2018 - now GOES-18: Oct 2022 - Now	Jul 2025 - Now
Access	GEE API	EUMETSAT API
Advantages	Can access via GEE API	Directly relevant to the ESA project. Greater spatial extent
Limitations	Only covers the Americas	Data only available from July 2025
Relevant links	GEE GOES-16 FDCF Data Guide	Data guide

- Since GOES likely will have more wildfire data (or could check if the spatial breadth of the MTG compensates for the short time period), we could initially work with GOES as our coarse input data. If time permits, we can try tuning the model with MTG?
- Alternatively we can combine MTG and MSG data to get a larger dataset for MTG (provided that the relevant data have the same spatial resolution. The IR bands of MSG are 3km while 1/2km for MTG.
- For MTG, we find that some bands have different resolution (0.5, 1, 2km) due to the different modes (High Resolution Fast Imagery (HRFI))
- Fire Radiative Power - MTG have data available in 1km from May 2026

High-res Target Data: Landsat vs Sentinel-2

	Landsat	**Sentinel-2**
Spatial res	30m	20m
Temporal res	8/16 days	5 days
Relevant Bands	Band 6 (1.6 μm), Band 7 (2.2 μm)	Band 11 (1.6 μm), Band 12 (2.2 μm)
Temporal extent	Landat 8: Apr 2013 -	March 2017 - present

	present Landsat 9: Sept 2019 - present In NASA FIRMS: June 2022 - present	
Spatial extent	NASA FIRMS: US/Canada only	global
Access	GEE	Copernicus Data Space Ecosystem (API)
Advantages	Contains thermal infrared band Fire dataset is available	Slightly higher resolution
Limitations	NASA FIRMS only have data from 2022, and only available for US/Canada	Lacks thermal infrared bands. As far as I can find, there is no clear 'fire' dataset that I could use

- Active fire detection in Landsat-8 imagery: A large-scale dataset and a deep-learning study Only fire data for Aug/Sept 2020 (worldwide).

Intermediate Resolution Data

- VIIRS:
  - Resolution: 375m
  - Access via NASA FIRMS
  - Temporal Res: Daily
- Main problem: when doing the cascaded approach, we need to spatially and temporally match VIIRS with the high-res target (landsat or Sentinel). There could potentially be a temporal mismatch

Main Research

Experiment	Architecture	Pipeline Structure	Input Resolution	Output resolution	Primary Research Objective
E1	Swin-UNet (deterministic)	Direct	2km (GOES)	30m (Landsat, Pereira et al.)	The baseline. Tests how a deterministic, CNN-based, direct downscaling handles big resolution leap.
E2	Swin-UNet (deterministic)	Cascaded	Stage 1: 2km -> 375m Stage 2: 375m -> 30m	30m (Landsat, Pereira et al.)	Tests if breaking the downscaling into two steps improves a deterministic model's accuracy.
E3	Conditional DDPM with Swin-UNet backbone (Probabilistic)	Direct	2km (GOES)	30m (Landsat, Pereira et al.)	Tests if a generative diffusion approach natively stops spatial blurring during big resolution leap.
E4	Conditional DDPM with Swin-UNet backbone (Probabilistic)	Cascaded	Stage 1: 2km -> 375m Stage 2: 375m -> 30m	30m (Landsat, Pereira et al.)	The proposed SOTA. Tests if combining step-wise pipelines with generative logic yields the highest fidelity.