Laboratory Session: April 30, 2024 Exercises due on: May 13, 2024

Exercise 1: Bayesian Inference for Poisson model

The number of particles emitted by a radioactive source during a fixed interval of time ($\Delta t = 10$ s) follows a Poisson distribution on the parameter μ . The number of particles observed during consecutive time intervals is: 4, 1, 3, 1, 5 and 3.

- (a) assuming a positive uniform prior distribution for the parameter μ
 - determine and draw the posterior distribution for μ , given the data
 - evaluate mean, median and variance, both analytically and numerically in R
- (b) assuming a Gamma prior such that the expected value is $\mu = 3$ with a standard deviation $\sigma = 1$,
 - determine and draw the posterior distribution for μ , given the data
 - evaluate mean, median and variance, both analytically and numerically in R.
- (c) evaluate a 95% credibility interval for the results obtained with different priors. Compare the result with that obtained using a normal approximation for the posterior distribution, with the same mean and standard deviation

Exercise 2: Efficiency using Bayesian approach

A researcher A wants to evaluate the efficiency of detector 2 (Det2). For this purpose, he sets up the apparatus shown in the figure 1, where Det2 is sandwiched between Det1 and Det3. Let $\bf n$ be the number of signals recorded simultaneously by Det1 and Det3, and $\bf r$ be those also recorded by Det2, researcher A obtains $\bf n=500$ and $\bf r=312$.




Figure 1:

Assuming a binomial model where n is the number of trials and r is the number of success out of n trials,

a) Evaluate the *mean* and the *variance* using a Bayesian approach under the hypothesis of:

- uniform prior $\sim \mathcal{U}(0,1)$
- Jeffrey's prior $\sim \text{Beta}(1/2, 1/2)$
- b) Plot the posterior distributions for both cases

Taking into account that the same detector has been studied by researcher B, who has performed only n = 10 measurements and has obtained r = 10 signals,

- c) Evaluate the *mean*, the *variance* and the *posterior* distribution using a uniform prior with the results of researcher B.
- d) Repeat the computation of points a) and b) with the data of researcher A using as a prior the posterior obtained from point c).
- e) [Optional] Compute 95% credible interval using the posterior of the previous point d).

Exercise 3 - Bayesian Inference for Binomial model

• A coin is flipped n = 30 times with the following outcomes:

```
T, T, T, T, H, T, T, H, H, T, T, H, H, T, H, H, H
```

- a) Assuming a flat prior, and a beta prior, plot the likelihood, prior and posterior distributions for the data set.
- b) Evaluate the most probable value for the coin probability p and, integrating the posterior probability distribution, give an estimate for a 95% credibility interval.
- c) Repeat the same analysis assuming a sequential analysis of the data. Show how the most probable value and the credibility interval change as a function of the number of coin tosses (i.e. from 1 to 30).
- d) Do you get a different result, by analysing the data sequentially with respect to a one-step analysis (i.e. considering all the data as a whole)?

Exercise 4 - Poll

A couple of days before an election in which four parties (A,B,C,D) compete, a poll is taken using a sample of 200 voters who express the following preferences 57, 31,45 and 67 for, respectively, parties A,B,C and D.

Using a Bayesian approach, for all parties

- Calculate the expected percentage of votes and a 68% credibility interval by assuming as prior a
 - uniform prior
 - a prior constructed from the results obtained from another poll conducted the previous week on a sample of 100 voters who expressed the following preferences 32,14,26,28 for, respectively, parties A,B,C and D.
- Sample size to obtain a margin of error less or equal than $\pm 3\%$ for each party