Ezequiel Boero <ezequiel.boero@adagioinstitute.org>

Re: Memory Production Paper (6) - My Response to Your Latest Comments

Ezequiel Boero <ezequiel.boero@adagioinstitute.org> Tue, Sep 29, 2026 at 12:07 AM
To: "Benjamin D. Summers" <bsummers@adagiollc.com>
Cc: "ISMAIL-SUTTON, SARA R." <sara.r.ismail-sutton@durham.ac.uk>

Hi Ben,

Revision 2.1 closes the file. All five qualifications are addressed, Corollary 2.8 is correct.
I found nothing to amend.

A couple of things to mention below:

Erdős. I checked the content against two independent source; the copy in the Erdős archive and Hajłasz's paper on the ambiguous locus,
which states the result as the set of points without a unique nearest point being covered by countably many surfaces of finite $(n-1)$-dimensional measure and being Lebesgue-null. Your paraphrase is faithful and the conclusion you draw follows,
Flat_n being closed. The attribution is correct.

Open Problem (vi), which I went after. You give linearised evidence at flat families. I did not reproduce that computation;
I ran the direct test instead, which is the more informative one: a search for the ratio $h_n/\|\Delta_V\|$ over 114 process-induced families
with $n \in \{2,3\}$ and $m \in \{3,4,5\}$ (grids of four, five and six points) nine restarts each. No ratio above 1 appeared**.**
The global maximum was 0.98305, at $m = 3$, and the maxima decrease with m: 0.9806 and 0.9406 at $m = 3$,
0.9304 and 0.8359 at $m = 4$, 0.8687 at $m = 5$. Since the computed h_n over-estimates, a ratio below 1 is strong evidence.
It is evidence and not an answer, and the paper is right to print the question only.

On "comparable" rather than "bi-Lipschitz" in Remark 2.7: the distinction is correct and worth keeping,
since the statement bounds the two functions against each other and says nothing about their moduli of continuity.

Besides that, I have nothing else to add**.** The paper is closed with no outstanding amendments.

Ezequiel

[Quoted text hidden]