

3.14. Show that a function $h(\omega)$ is \mathcal{F}_t -measurable if and only if h is a pointwise limit (for a.a. ω) of sums of functions of the form

$$g_1(B_{t_1}) \cdot g_2(B_{t_2}) \cdots g_k(B_{t_k})$$

where g_1, \dots, g_k are bounded continuous functions and $t_j \leq t$ for $j \leq k$, $k = 1, 2, \dots$

Hint: Complete the following steps:

a) We may assume that h is bounded.

b) For $n = 1, 2, \dots$ and $j = 1, 2, \dots$ put $t_j = t_j^{(n)} = j \cdot 2^{-n}$. For fixed n let \mathcal{H}_n be the σ -algebra generated by $\{B_{t_j}(\cdot)\}_{t_j \leq t}$. Then by Corollary C.9

$$h = E[h|\mathcal{F}_t] = \lim_{n \rightarrow \infty} E[h|\mathcal{H}_n] \quad (\text{pointwise a.e. limit})$$

c) Define $h_n := E[h|\mathcal{H}_n]$. Then by the Doob-Dynkin lemma (Lemma 2.1.2) we have

$$h_n(\omega) = G_n(B_{t_1}(\omega), \dots, B_{t_k}(\omega))$$

for some Borel function $G_n: \mathbf{R}^k \rightarrow \mathbf{R}$, where $k = \max\{j; j \cdot 2^{-n} \leq t\}$. Now use that any Borel function $G: \mathbf{R}^k \rightarrow \mathbf{R}$ can be approximated pointwise a.e. by a continuous function $F: \mathbf{R}^k \rightarrow \mathbf{R}$ and complete the proof by applying the Stone-Weierstrass theorem.

Assume that h is bounded and put $t_j = t_j^{(n)} = j \cdot 2^{-n}$,
 $n = 1, 2, \dots$, $j = 1, 2, \dots$

For a fixed n , let \mathcal{H}_n be the σ -algebra generated by $\{B_{t_j}\}_{t_j \leq t}$. Then

$$h = E[h|\mathcal{F}_t] = \lim_{n \rightarrow \infty} E[h|\mathcal{H}_n] \quad \text{pointwise a.e.}$$

Let $h_n = E[h|\mathcal{H}_n]$. By the Doob-Dynkin lemma,

$$h_n(\omega) = G_n(B_{t_1}(\omega), \dots, B_{t_k}(\omega))$$

for some Borel function G_n , and where $k = \max\{j; j \cdot 2^{-n} \leq t\}$.

Since every Borel function $G: \mathbf{R}^k \rightarrow \mathbf{R}$ can be approximated pointwise a.e. by a continuous function $F: \mathbf{R}^k \rightarrow \mathbf{R}$, by the Stone-Weierstrass Theorem applied on $[t_1, t_k]$, G can be approximated by a polynomial function g_n , completing the proof.

Corollary C.9. *Let $X \in L^1(P)$, let $\{\mathcal{N}_k\}_{k=1}^\infty$ be an increasing family of σ -algebras, $\mathcal{N}_k \subset \mathcal{F}$ and define \mathcal{N}_∞ to be the σ -algebra generated by $\{\mathcal{N}_k\}_{k=1}^\infty$. Then*

$$E[X|\mathcal{N}_k] \rightarrow E[X|\mathcal{N}_\infty] \quad \text{as } k \rightarrow \infty ,$$

a.e. P and in $L^1(P)$.

Lemma 2.1.2. *If $X, Y: \Omega \rightarrow \mathbf{R}^n$ are two given functions, then Y is \mathcal{H}_X -measurable if and only if there exists a Borel measurable function $g: \mathbf{R}^n \rightarrow \mathbf{R}^n$ such that*

$$Y = g(X) .$$