

1 Wavy heliospheric current sheet

Both components of the drift velocity are modified by the waviness:

- There is a Heaviside step function in the Parker field:

$$\vec{B} = A_c B_0 \left(\frac{r_0}{r} \right)^2 \left(\hat{r} - \frac{\Omega r \sin \theta}{V_{sw}} \hat{\phi} \right) H(\theta' - \theta),$$

where θ' is the angular extent of the HCS. In Strauss et al 2012 this is given by eq 11:

$$\theta' = \frac{\pi}{2} + \sin^{-1} \left[\sin \alpha \sin \left(\phi - \phi_0(t) + \frac{\Omega r}{V_{sw}} \right) \right],$$

where $\phi_0(t) = \phi_0 + \Omega t$ and ϕ_0 is an arbitrary phase. In the code, the step function appears when calculating the gradient/curvature part of the drift velocity.

TODO:

- How is ϕ_0 determined?
- α should also be time dependent. How should I implement this? Parameter file should probably point to table of α vs t , which the parameter object can then parse. The date at which the particle was observed at Earth can be specified in the run file.
- The HCS drift velocity is also affected. It takes the form (eq 17)

$$\vec{v}_{hcs} = v_{hcs} \left[\cos A_\xi \xi \sin \Psi \hat{r} + \sin A_\xi \xi \hat{\theta} + \cos A_\xi \xi \cos \Psi \hat{\phi} \right] A_c \operatorname{sgn} q.$$

The angle ξ (β in the reference) between the tangent to the HCS and the radial line passing through the point on the HCS is given by

$$\begin{aligned} \tan^2 \xi &= \left[-r \frac{\partial \theta'}{\partial r} \right]^2 + \left[\frac{1}{\sin \theta} \frac{\partial \theta'}{\partial \phi} \right]^2 \\ \implies \tan \xi &= \frac{\Omega r}{V_{sw}} \frac{1}{\sin \Psi} \frac{\sqrt{\sin^2 \alpha - \cos^2 \theta'}}{\sin \theta'} \quad (\text{for Parker field}). \end{aligned}$$

The sign of ξ , $A_\xi = \pm 1$, is equal to the sign of $\partial \theta' / \partial r$:

$$A_\xi = \operatorname{sgn} \cos \left(\phi - \phi_0(t) + \frac{\Omega r}{V_{sw}} \right).$$

v_{hcs} is given by the usual approximation:

$$v_{hcs} = \left[0.457 - 0.412 \frac{L}{r_L} + 0.915 \frac{L^2}{r_L^2} \right] v,$$

where L is the smallest distance from the particle's position to the HCS. If $L > 2r_L$, v_{hcs} is zero. Finally, ξ is evaluated at the HCS point which minimizes L .

The hard part of these modifications is calculating L , which cannot be done analytically and is expensive. If

- $|\theta - \frac{\pi}{2}| \leq \alpha$,

- $\theta < \frac{\pi}{2} - \alpha$ and $L_+^{th} \leq 2r_L$, where $L_+^{th} = r \cos(\alpha + \theta)$ is the distance from the particle to the surface bounding the HCS above,
- Or $\theta > \frac{\pi}{2} + \alpha$ and $L_-^{th} = -r \cos(\theta - \alpha) \leq 2r_L$,

we compute L using the Nelder-Mead simplex algorithm. Here are the steps for minimizing $L(r_s, \phi_s)$, where r_s and ϕ_s are the coordinates of a point in the sheet:

0. Choose an initial set of points $\vec{x}_1, \vec{x}_2, \vec{x}_3$.
1. Order vertices: $L(\vec{x}_1) \leq L(\vec{x}_2) \leq L(\vec{x}_3)$.
2. Compute $\vec{x}_O = \frac{1}{2}(\vec{x}_1 + \vec{x}_2)$.
3. Compute reflected point $\vec{x}_R = \vec{x}_O + \alpha(\vec{x}_O - \vec{x}_3)$.
If $L(\vec{x}_1) \leq L(\vec{x}_R) < L(\vec{x}_2)$, replace \vec{x}_3 with \vec{x}_R and go to step 1.
4. If $L(\vec{x}_R) < L(\vec{x}_1)$, compute expanded point $\vec{x}_E = \vec{x}_O + \gamma(\vec{x}_R - \vec{x}_O)$.
If $L(\vec{x}_E) < L(\vec{x}_R)$, replace \vec{x}_3 with \vec{x}_E and go to step 1.
Else, replace \vec{x}_3 with \vec{x}_R and go to step 1.
Else, go to step 5.
5. Compute contracted point $\vec{x}_C = \vec{x}_O + \rho(\vec{x}_3 - \vec{x}_O)$.
If $L(\vec{x}_C) < L(\vec{x}_3)$, replace \vec{x}_3 with \vec{x}_C and go to step 1.
Else, go to step 6.
6. For all but the \vec{x}_1 , replace \vec{x}_i with $\vec{x}_1 + \sigma(\vec{x}_i - \vec{x}_1)$. Go to step 1.

Standard values for the constants are $\alpha = 1$, $\gamma = 2$, $\rho = 1/2$, $\sigma = 1/2$.