

## TTIC 31230 Fundamentals of Deep Learning

### Langevin Dynamics Problems.

**Problem 1.** This problem is on batch size scaling of Langevin dynamics. We consider batched SGD as defined by

$$\Phi \leftarrow \eta \hat{g}^B$$

where  $\hat{g}^B$  is the average of  $B$  sampled gradients. Let  $g$  be the average gradient  $g = E \hat{g}$ .

The covariance matrix at batch size  $B$  is

$$\Sigma^B[i, j] = E (\hat{g}^B[i] - g[i])(\hat{g}^B[j] - g[j]).$$

Langevin dynamics is

$$\Phi(t + \Delta t) = \Phi(t) - g\Delta t + \epsilon\sqrt{\Delta t} \quad \epsilon \sim \mathcal{N}(0, \eta\Sigma^B)$$

Show that for  $\eta = B\eta_0$  the Langevin dynamics is determined by  $\eta_0$  independent of  $B$ .

**Solution:**

$$\begin{aligned} \Sigma^B[i, j] &= E (\hat{g}^B[i] - g[i])(\hat{g}^B[j] - g[j]) \\ &= \frac{1}{B^2} E \left( \sum_b \hat{g}_b[i] - g[i] \right) \left( \sum_b \hat{g}_b[j] - g[j] \right) \\ &= \frac{1}{B^2} E \sum_{b, b'} (\hat{g}_b[i] - g[i]) (\hat{g}_{b'}[j] - g[j]) \\ &= \frac{1}{B^2} \sum_{b, b'} E (\hat{g}_b[i] - g[i]) (\hat{g}_{b'}[j] - g[j]) \\ &= \frac{1}{B^2} \sum_b E (\hat{g}_b[i] - g[i]) (\hat{g}_b[j] - g[j]) \\ &= \frac{1}{B} \Sigma^1[i, j] \end{aligned}$$

So for  $\eta = B\eta_0$  we have  $\eta\Sigma^B = \eta_0\Sigma^1$  which yields the equivalence.