

Contents

1	Sequences	1
1.1	Sequences	1
1.1.1	Sequences	1
1.2	Series	2
1.2.1	Series	2
1.2.2	Summation of natural numbers	4
1.3	Products	4
1.3.1	Products	4
1.3.2	Factorials	5

1 Sequences

1.1 Sequences

1.1.1 Sequences

1.1.1.1 Definition

A sequence is an ordered list of terms.

These are commonly maps from natural numbers to real (or complex) numbers.

We can use $a_i = f(i)$ to denote this.

If $f(i)$ is defined on all $i \in \mathbb{N}$ then the sequence is infinite. Otherwise it is finite.

If a sequence is defined on $n \in \mathbb{N}$ and $n \neq 0$ then the sequence must be defined on $n - 1$.

For example a_0, a_1, a_2, \dots is a sequence, but a_1, a_2, \dots is not.

1.1.1.2 Monotone sequence

A monotone sequence is one where each element is succeeded ordinally by the next entry.

For example:

$\langle 1, 2, 3, 6, 7 \rangle$ is monotone

$\langle 1, 2, 3, 3, 4 \rangle$ is not monotone

An increasing sequence is one where:

$$\forall m \in \mathbb{N} \forall n \in \mathbb{N} [m > n \leftrightarrow a_m \geq a_n]$$

A strictly increasing sequence is one where:

$$\forall m \in \mathbb{N} \forall n \in \mathbb{N} [m > n \leftrightarrow a_m > a_n]$$

A decreasing sequence is one where:

$$\forall m \in \mathbb{N} \forall n \in \mathbb{N} [m > n \leftrightarrow a_m \leq a_n]$$

A strictly decreasing sequence is one where:

$$\forall m \in \mathbb{N} \forall n \in \mathbb{N} [m > n \leftrightarrow a_m < a_n]$$

All strictly decreasing sequences are decreasing, and all strictly increasing sequences are increasing.

A monotone sequence is one which is either increasing or decreasing.

1.1.1.3 Subsequences

A subsequence of a sequence is the original sequence with some elements of the original removed, not changing the order.

For example:

$\langle 1, 3, 5 \rangle$ is a subsequence of $\langle 2, 1, 3, 4, 7, 5 \rangle$

1.1.1.4 Bounded sequence

A function $f(x)$ on set X is bounded if:

$$\exists M \in \mathbb{R} [\forall x \in X f(x) \leq M]$$

A bounded sequence is a special case of a bounded function where:

$$X = \mathbb{N}$$

That is, a sequence is bounded by M iff:

$$\forall n \in \mathbb{N} |f(a_n)| < le M$$

1.2 Series

1.2.1 Series

1.2.1.1 Definition

A series is the summation of a sequence. For a series a_n there is a corresponding series:

$$s_n = \sum_{i=0}^n a_i$$

Where:

$$\sum_{i=0}^n a_i = a_0 + a_1 + a_2 + \dots + a_n$$

1.2.1.2 Multiplication of summations

If all members of a sequence are multiplied by a constant, so is each member of the series.

We can take constants out of the series:

$$s_n = \sum_{i=0}^n a_i$$

$$s_n = \sum_{i=0}^n cb_i$$

$$s_n = a \sum_{i=0}^n b_i$$

1.2.1.3 Summation of constants

If all elements of a sequence are the same, then the series is a multiple of that constant.

$$s_n = \sum_{i=0}^n a_i$$

$$s_n = \sum_{i=0}^n c$$

$$s_n = nc$$

1.2.1.4 Addition of summations

Consider a sequence $a_i = b_i + c_i$.

$$s_n = \sum_{i=0}^n a_i$$

$$s_n = \sum_{i=0}^n (b_i + c_i)$$

We can then split this out.

$$s_n = \sum_{i=0}^n b_i + \sum_{i=0}^n c_i$$

1.2.1.5 Summation from a different start point

$$\sum_{i=0}^n a_i = a_0 + \sum_{i=1}^n a_i$$

1.2.1.6 Multiple summations

$$\sum_{i=0}^n \sum_{j=0}^m a_i = n \sum_{j=0}^m a_i$$

$$\sum_{i=0}^n \sum_{j=0}^m a_i b_j = \sum_{i=0}^n a_i \sum_{j=0}^m b_i$$

1.2.2 Summation of natural numbers

1.2.2.1 Goal

Let's prove that:

$$\sum_{i=0}^n i = \frac{n(n+1)}{2}$$

1.2.2.2 Proof by induction

We use the inference rules Modus Ponens, which says that if X is true, and $X \rightarrow Y$ is true, then Y is true.

1.2.2.3 True for $n = 0$

We know this is true for $n = 0$:

$$0 = \frac{0(0+1)}{2}$$

$$0 = 0$$

1.2.2.4 If it's true for n , it's true for $n + 1$

We can also prove that if it true for n , it is true for $n + 1$.

$$\sum_{i=0}^{n+1} i = \frac{(n+1)(n+2)}{2}$$

$$(n+1) + \sum_{i=0}^n i = \frac{n^2+3n+2}{2}$$

If it is true for n , then:

$$(n+1) + \frac{n(n+1)}{2} = \frac{n^2+3n+2}{2}$$

$$\frac{n^2+3n+2}{2} = \frac{n^2+3n+2}{2}$$

$$1 = 1$$

1.2.2.5 Result

So we know that it is true for $n = 0$, and if it is true for n , then it is true for $n + 1$. As a result it is true for all natural numbers.

1.3 Products

1.3.1 Products

A product is a repeated multiplication of a sequence.

$$p_n = \prod_{i=0}^n s_i$$

1.3.1.1 Multiplication of products

We can take constants out of the product.

$$p_n = \prod_{i=0}^n ca_i$$

$$p_n = a^n \sum_{i=0}^n a_i$$

1.3.1.2 Products of constants

If $a_i = c$ then the summation is then of the form:

$$p_n = \prod_{i=0}^n c$$

$$p_n = c^n \prod_{i=j}^n 1$$

$$p_n = c^n$$

1.3.1.3 Combining products

If a sequence is the product of two other sequences then the product of the sequence is equal to the product of the two individual sequences.

$$p_n = \prod_{i=0}^n a_i$$

$$p_n = \prod_{i=0}^n b_i c_i$$

$$p_n = \prod_{i=0}^n b_i \prod_{i=0}^n c_i$$

1.3.2 Factorials

A factorial is a product across natural numbers. That is:

$$n! := \prod_{i=0}^n i$$