

UNSTABLE $m = 1$ ECCENTRIC MODES IN THERMALLY COOLING, SELF GRAVITATING DISKS

ADAM M. DEMPSEY & YORAM LITHWICK

Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA) and Department of Physics and Astronomy
Northwestern University 2145 Sheridan Road Evanston, IL 60208 USA

Draft version March 23, 2015

ABSTRACT

1. INTRODUCTION

2. EQUATIONS OF MOTION

The equations of motion for a two dimensional, viscous disk are

$$(\partial_t + \mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{\nabla P}{\Sigma} - \nabla \Phi + \frac{1}{\Sigma} \nabla \cdot (\nu \Sigma \mathbf{D}) \quad (1)$$

$$(\partial_t + \mathbf{v} \cdot \nabla) \Sigma = -\Sigma \nabla \cdot \mathbf{v} \quad (2)$$

$$T(\partial_t + \mathbf{v} \cdot \nabla) s = -\frac{T - \bar{T}}{t_{\text{cool}}} \quad (3)$$

Where all quantities are assumed to be a vertical average of their three dimensional counterparts. Here, Σ is the surface density, s is the entropy of the fluid, T is the disk temperature, and \mathbf{D} is the viscous stress tensor. These equations describe a viscous fluid that is thermally heated and or cooled to some "background" temperature profile \bar{T} over a timescale t_{cool} . To close the system of equations we assume the fluid is ideal and perfect, so that $P = c_v(\gamma - 1)\Sigma T$ and $s \equiv c_v \ln(P\Sigma^{-\gamma})$. Following Ostriker, Shu, & Adams (2), we can connect the two dimension adiabatic index γ to the true three dimensional adiabatic index Γ by assuming that the gas is vertically isothermal and gaussian distributed.

We consider a disk with a prescribed azimuthally averaged background profile given by $(\bar{v}_r, \bar{v}_\phi, \bar{\Sigma}, \bar{T}) = (0, r\Omega(r), \bar{\Sigma} \propto r^\mu, \bar{T} \propto r^\delta)$. We look at linear perturbations to this background state. Denoted linear quantities with a prime and background quantities with a bar, the linearized equations of motion become

$$(\partial_t + \Omega \partial_\phi) v'_r - 2\Omega v'_\phi = -\frac{1}{\Sigma} \partial_r P' + \frac{\Sigma'}{\bar{\Sigma}^2} \frac{d\bar{P}}{dr} - \partial_r \Phi' + \text{visc.} \quad (4)$$

$$(\partial_t + \Omega \partial_\phi) v'_\phi + \frac{\kappa^2}{2\Omega} v'_r = -\frac{1}{r\bar{\Sigma}} \partial_\phi P' - \frac{1}{r} \partial_\phi \Phi' + \text{visc.} \quad (5)$$

$$(\partial_t + \Omega \partial_\phi) \Sigma' + v'_r \frac{d\bar{\Sigma}}{dr} = -\bar{\Sigma} \left(\frac{1}{r} \partial_r (r v'_r) + \frac{1}{r} \partial_\phi (v'_\phi) \right) \quad (6)$$

$$(\partial_t + \Omega \partial_\phi) s' + v'_r \frac{d\bar{s}}{dr} = -\beta \Omega \frac{T'}{\bar{T}} \quad (7)$$

Where we've parametrized the cooling time as $t_{\text{cool}}^{-1} = \beta \Omega$ (Gammie 2001). We now Fourier Transform the linear variables in the azimuthal direction as well as in time and write $(v'_r, v'_\phi, \Sigma', P', s', T', \Phi') = \sum_m (u, v, \sigma, p, s, T, \phi) e^{im(\phi - \Omega_p t)}$. We further work under

the assumption that the different m modes only couple to the $m = 0$ mode (i.e the background) and not to other $m \neq 0$ modes. The linearized equations of motion are now

$$im(\Omega - \Omega_p)u - 2\Omega v = -\frac{1}{\bar{\Sigma}} \frac{dp}{dr} + \frac{\sigma}{\bar{\Sigma}^2} \frac{d\bar{P}}{dr} - \frac{d\phi}{dr} + \text{visc.} \quad (8)$$

$$im(\Omega - \Omega_p)v + \frac{\kappa^2}{2\Omega} u = -\frac{imp}{r\bar{\Sigma}} - \frac{im\phi}{r} + \text{visc.} \quad (9)$$

$$im(\Omega - \Omega_p) \frac{\sigma}{\bar{\Sigma}} + u \frac{d \ln \bar{\Sigma}}{dr} = -\left(\frac{1}{r} \frac{d}{dr} (ru) + \frac{imv}{r} \right) \quad (10)$$

$$im \left[\left(1 - i \frac{\beta}{m} \right) \Omega - \Omega_p \right] \frac{T'}{\bar{T}} + u \frac{d \ln \bar{T}}{dr} = -(\gamma - 1) \left(\frac{1}{r} \frac{d}{dr} (ru) + \frac{imv}{r} \right) \quad (11)$$

Where we've replaced the entropy in favor of the temperature through the relations

$$s' = \frac{T'}{\bar{T}} - (\gamma - 1) \frac{\sigma}{\bar{\Sigma}} \quad (12)$$

$$\frac{d\bar{s}}{dr} = \frac{d \ln \bar{T}}{dr} - (\gamma - 1) \frac{d \ln \bar{\Sigma}}{dr} \quad (13)$$

3. $m = 1$ SLOW MODES

We specialize to $m = 1$ modes and to the low pattern speed limit $\Omega_p \ll \Omega$. We can simplify the 4 dimensional set of equations to just one equation describing the global structure and evolution of an eccentricity vector, $e(r)$ (5; 4; 3). The eccentricity is defined through the Lagrangian displacement ξ_r as

$$e \equiv \frac{\xi_r}{r} \quad (14)$$

3.1. Boundary Condition

We adopt a zero Lagrangian pressure boundary condition at both the inner and outer boundaries. Using the equations of motion it can be shown that this is equivalent to satisfying

$$r \frac{de(r)}{dr} = \left[\frac{\beta^2}{\gamma^2 + \beta^2} - i \frac{\gamma\beta}{\gamma^2 + \beta^2} \right] e(r) \quad (15)$$

at each boundary in the slow mode approximation.

4. METHODS

5. RESULTS

We studied the growth rates of unstable modes in the slow mode ($\Omega_p \ll \Omega_k$) approximation. The classical WKB dispersion relation for a self gravitating and isothermal disk (see e.g Binney & Tremaine (1)) is

$$m^2(\Omega - \Omega_p)^2 = \kappa^2 - 2\pi G\Sigma|k_r| + c^2 k_r^2 \quad (16)$$

In the $m = 1$ slow mode limit this simplifies to (3)

$$2\Omega(\omega_p - \Omega_p) = -2\pi G\Sigma|k_r| + c^2 k_r^2 \quad (17)$$

Where $\omega_p = \Omega - \kappa$ is the orbital precession frequency of a fluid element. From the dispersion relation it is clear

that pressure creates retrograde modes ($\Omega_p < 0$) and self gravity tends to create prograde modes ($\Omega_p > 0$). Additionally, global modes (i.e low k_r) also tend to create prograde modes (3).

5.1. *Lowest Order Mode with No Self Gravity*

The mode with zero nodes is perhaps the simplest to understand.

5.2. *Self Gravity*

6. DISCUSSION

7. CONCLUSIONS

REFERENCES

- Binney, J., Tremaine, S. *Galactic Dynamics*. Princeton University Press, 2002.
- Gammie, C. F. (2001). ApJ, 553(1), 174-183.
- Ostriker, E. C., Shu, F. H., & Adams, F. C. (1992). ApJ, 399, 192-212.
- Papaloizou, J. C. B. (2002). Astronomy and Astrophysics, 388(2), 615-631.
- Goodchild, S., & Ogilvie, G. (2006). MNRAS, 368(3), 1123-1131.
- Ogilvie, G. I. (2001). MNRAS, 325(1), 231-248.