

Density Theorems in Analytic function spaces

Pierre-Olivier Parisé (Université Laval)
Advisor, T. Ransford

Colloque panquébécois des étudiantes et étudiants de l'ISM
Université de Sherbrooke

25th-27th, May, 2018

Spaces of continuous functions

Theorem (1885, Weierstrass)

Let $f \in \mathcal{C}[a, b]$ be a continuous function on the closed interval $[a, b]$. Then, there exists a sequence of polynomials (p_n) such that

$$\|f - p_n\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Spaces of continuous functions

Theorem (1885, Weierstrass)

Let $f \in \mathcal{C}[a, b]$ be a continuous function on the closed interval $[a, b]$. Then, there exists a sequence of polynomials (p_n) such that

$$\|f - p_n\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Several proofs:

- ① Weierstrass by using convolution with the Heat Kernel;
- ② Bernstein, with the Bernstein polynomials;
- ③ Stone, with the Stone-Weierstrass theorem;
- ④ With the Hahn-Banach theorem.

Spaces of continuous functions

Theorem (1912, Bernstein)

If $f \in C[0, 1]$ and $p_n(x) := \sum_{k=0}^n \binom{n}{k} f\left(\frac{k}{n}\right) x^k (1-x)^{n-k}$, then

$$\|f - p_n\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Spaces of continuous functions

Theorem (1912, Bernstein)

If $f \in C[0, 1]$ and $p_n(x) := \sum_{k=0}^n \binom{n}{k} f\left(\frac{k}{n}\right) x^k (1-x)^{n-k}$, then

$$\|f - p_n\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Remark. Exact constructive proof method!

Restatement of Weierstrass' theorem

Restatement of Weierstrass' theorem

For $f \in \mathcal{C}[a, b]$, we define

$$\|f\|_{\infty} := \sup_{x \in [a, b]} |f(x)|.$$

Restatement of Weierstrass' theorem

For $f \in \mathcal{C}[a, b]$, we define

$$\|f\|_{\infty} := \sup_{x \in [a, b]} |f(x)|.$$

Theorem (1885, Weierstrass)

The polynomials are dense in $(\mathcal{C}[a, b], \|\cdot\|_{\infty})$.

Restatement of Weierstrass' theorem

For $f \in \mathcal{C}[a, b]$, we define

$$\|f\|_{\infty} := \sup_{x \in [a, b]} |f(x)|.$$

Theorem (1885, Weierstrass)

The polynomials are dense in $(\mathcal{C}[a, b], \|\cdot\|_{\infty})$.

In general, given a space of functions V , one can ask many questions about the structure of V :

- 1 Can we endow V with a norm that makes it a Banach space?

Restatement of Weierstrass' theorem

For $f \in \mathcal{C}[a, b]$, we define

$$\|f\|_{\infty} := \sup_{x \in [a, b]} |f(x)|.$$

Theorem (1885, Weierstrass)

The polynomials are dense in $(\mathcal{C}[a, b], \|\cdot\|_{\infty})$.

In general, given a space of functions V , one can ask many questions about the structure of V :

- 1 Can we endow V with a norm that makes it a Banach space?
- 2 With a given norm, can we find approximants in a “nice” subset of V for an $f \in V$?

Hardy space: Definition

Let $f(z) := \sum_{n=0}^{\infty} a_n z^n$, $z \in \mathbb{D} := \{|z| < 1\}$.

Definition

$$H^2 := \{f \in \mathcal{H}(\mathbb{D}) : \sum_{n=0}^{\infty} |a_n|^2 < +\infty\}.$$

Hardy space: Definition

Let $f(z) := \sum_{n=0}^{\infty} a_n z^n$, $z \in \mathbb{D} := \{|z| < 1\}$.

Definition

$$H^2 := \{f \in \mathcal{H}(\mathbb{D}) : \sum_{n=0}^{\infty} |a_n|^2 < +\infty\}.$$

We define:

- A norm: $\|f\|_2^2 := \sum_{n=0}^{\infty} |a_n|^2$;
- The partial sums: $s_n(f)(z) := \sum_{k=0}^n a_k z^k$.

Hardy space: Polynomial approximants

Theorem

If $f \in H^2$, then $\|f - s_n(f)\|_2 \rightarrow 0$ when $n \rightarrow +\infty$.

Hardy space: Polynomial approximants

Theorem

If $f \in H^2$, then $\|f - s_n(f)\|_2 \rightarrow 0$ when $n \rightarrow +\infty$.

Proof.

We have $f - s_n(f) = \sum_{k=n+1}^{\infty} a_k z^k$ and so

$$\|f - s_n(f)\|_2^2 = \sum_{k=n+1}^{\infty} |a_k|^2 \xrightarrow{n \rightarrow +\infty} 0.$$



Disk Algebra: Definition

Definition

$$\mathcal{A}(\mathbb{D}) := \mathcal{H}(\mathbb{D}) \cap \mathcal{C}(\overline{\mathbb{D}}).$$

Disk Algebra: Definition

Definition

$$\mathcal{A}(\mathbb{D}) := \mathcal{H}(\mathbb{D}) \cap \mathcal{C}(\overline{\mathbb{D}}).$$

For $f \in \mathcal{A}(\mathbb{D})$, we define:

$$\|f\|_{\infty} := \sup_{z \in \mathbb{D}} |f(z)|.$$

Disk Algebra: Definition

Definition

$$\mathcal{A}(\mathbb{D}) := \mathcal{H}(\mathbb{D}) \cap \mathcal{C}(\overline{\mathbb{D}}).$$

For $f \in \mathcal{A}(\mathbb{D})$, we define:

$$\|f\|_{\infty} := \sup_{z \in \mathbb{D}} |f(z)|.$$

A suprising fact (du Bois Reymond):

$$\exists f \in \mathcal{A}(\mathbb{D}) \text{ s.t. } \sup_{n \geq 0} \|s_n f\|_{\infty} = +\infty.$$

Theorem

If $f \in \mathcal{A}(\mathbb{D})$, then $\|\sigma_n(f) - f\|_\infty \rightarrow 0$ as $n \rightarrow \infty$.

Theorem

If $f \in \mathcal{A}(\mathbb{D})$, then $\|\sigma_n(f) - f\|_\infty \rightarrow 0$ as $n \rightarrow \infty$.

Proof.

$g := f|_{\partial\mathbb{D}}$ is continuous. So, from Fejér theorem, the Cesàro means $\sigma_n(f) := \frac{\sum_{k=0}^n s_k(f)}{n+1}$ converge to g uniformly. $\sigma_n(f)$ is the restriction of a polynomial to $\partial\mathbb{D}$ and

$$\|\sigma_n(f) - f\|_\infty = \sup_{z \in \partial\mathbb{D}} |\sigma_n(f)(z) - g(z)| \xrightarrow{n \rightarrow +\infty} 0.$$



De Branges-Rovnyak spaces: Definition

Let $H^\infty := \{f \in \mathcal{H}(\mathbb{D}) : \|f\|_\infty < +\infty\}$.

De Branges-Rovnyak spaces: Definition

Let $H^\infty := \{f \in \mathcal{H}(\mathbb{D}) : \|f\|_\infty < +\infty\}$.

Definition

Let $b \in H^\infty$ with $\|b\|_\infty \leq 1$. Then,

$$\mathcal{H}(b) := \left\{ f \in H^2 : \sup_{g \in H^2} (\|f + gb\|_2^2 - \|g\|_2^2) < +\infty \right\}.$$

We define a norm:

$$\|f\|_b^2 := \sup_{g \in H^2} (\|f + gb\|_2^2 - \|g\|_2^2).$$

Example 1.

Let $b_0(z) := \frac{\tau z}{1 - \tau^2 z}$ where $\tau := \frac{\sqrt{5}-1}{2}$. If $f \in \mathcal{H}(b_0)$, then

$$\|f_r - f\|_{b_0} \rightarrow 0$$

as $r \rightarrow 1^-$ where $f_r(z) := f(rz)$.

Example 2.

However, let $b := b_0 B$ where B is a Blaschke product with zeros at $w_n := 1 - 4^{-n}$ for $n \geq 1$.

Example 2.

However, let $b := b_0 B$ where B is a Blaschke product with zeros at $w_n := 1 - 4^{-n}$ for $n \geq 1$.

If we let

$$f(z) := \sum_{n \geq 1} \frac{2^{-n}}{1 - w_n z},$$

then it is possible to show that $f \in \mathcal{H}(b)$, but

Example 2.

However, let $b := b_0 B$ where B is a Blaschke product with zeros at $w_n := 1 - 4^{-n}$ for $n \geq 1$.

If we let

$$f(z) := \sum_{n \geq 1} \frac{2^{-n}}{1 - w_n z},$$

then it is possible to show that $f \in \mathcal{H}(b)$, but

- $\lim_{r \rightarrow 1^-} \|f_r\|_b = +\infty$;
- $\limsup_{n \rightarrow \infty} \|s_n(f)\|_b = +\infty$;
- $\limsup_{n \rightarrow \infty} \|\sigma_n(f)\|_b = +\infty$.

Example 2.

However, let $b := b_0 B$ where B is a Blaschke product with zeros at $w_n := 1 - 4^{-n}$ for $n \geq 1$.

If we let

$$f(z) := \sum_{n \geq 1} \frac{2^{-n}}{1 - w_n z},$$

then it is possible to show that $f \in \mathcal{H}(b)$, but

- $\lim_{r \rightarrow 1^-} \|f_r\|_b = +\infty$;
- $\limsup_{n \rightarrow \infty} \|s_n(f)\|_b = +\infty$;
- $\limsup_{n \rightarrow \infty} \|\sigma_n(f)\|_b = +\infty$.

We are in trouble...

Characterization

The "quality" of $\mathcal{H}(b)$ depends on the nature of b .

Characterization

The "quality" of $\mathcal{H}(b)$ depends on the nature of b .
When b is a non-extreme point of B_{H^∞} , then

Theorem (1994, Sarason[6])

Let $b \in H^\infty$ and $\|b\|_\infty \leq 1$. Then, the following statements are equivalent:

- ① b is non-extreme;
- ② $\log(1 - |b|^2) \in L^1(\partial\mathbb{D})$;
- ③ $\mathcal{H}(b)$ contains all holomorphic functions in $\overline{\mathbb{D}}$;
- ④ The polynomials form a dense subset of $\mathcal{H}(b)$.

Characterization

Two proofs:

Characterization

Two proofs:

- From Sarason, by showing that \mathcal{P}^\perp is trivial in $\mathcal{H}(b)$.

Characterization

Two proofs:

- From Sarason, by showing that \mathcal{P}^\perp is trivial in $\mathcal{H}(b)$.
- From O. El-Fallah, E. Fricain, K. Kellay, J. Mashreghi, and T. Ransford by using Toeplitz operators and building up polynomials to approach a giving function in $\mathcal{H}(b)$.

Characterization

So, we can ask what happen when b is an extreme point
($\log(1 - |b|^2) = -\infty$ a.e.)

Characterization

So, we can ask what happen when b is an extreme point
($\log(1 - |b|^2) = -\infty$ a.e.)

Theorem (2017, A. Aleman, B. Malman[1])

The set $\mathcal{A}(\mathbb{D}) \cap \mathcal{H}(b)$ is dense in $\mathcal{H}(b)$.

Characterization

So, we can ask what happen when b is an extreme point
 $(\log(1 - |b|^2) = -\infty \text{ a.e.})$

Theorem (2017, A. Aleman, B. Malman[1])

The set $\mathcal{A}(\mathbb{D}) \cap \mathcal{H}(b)$ is dense in $\mathcal{H}(b)$.

Remark.

- $\mathcal{H}(b)$ may contain no function in $\mathcal{H}(\overline{\mathbb{D}})$.

Characterization

So, we can ask what happen when b is an extreme point
($\log(1 - |b|^2) = -\infty$ a.e.)

Theorem (2017, A. Aleman, B. Malman[1])

The set $\mathcal{A}(\mathbb{D}) \cap \mathcal{H}(b)$ is dense in $\mathcal{H}(b)$.

Remark.

- $\mathcal{H}(b)$ may contain no function in $\mathcal{H}(\overline{\mathbb{D}})$.
- The proof is not constructive.

Characterization

So, we can ask what happen when b is an extreme point
($\log(1 - |b|^2) = -\infty$ a.e.)


Theorem (2017, A. Aleman, B. Malman[1])

The set $\mathcal{A}(\mathbb{D}) \cap \mathcal{H}(b)$ is dense in $\mathcal{H}(b)$.

Remark.

- $\mathcal{H}(b)$ may contain no function in $\mathcal{H}(\overline{\mathbb{D}})$.
- The proof is not constructive.
- There is no known constructive proof of this fact!!

Thanks for your attention!



Bibliography I



A. Aleman and B. Malman.

Density of Disk Algebra Functions in de Branges–Rovnyak Spaces.
Comptes Rendus Mathematique, 355(8):871–875, 2017.



J. B. Conway.

Functions of One Complex Variable II.
Graduate Texts in Mathematics. Springer-Verlag, 1995.



P. L. Duren.

Theory of H^p spaces, volume 38 of *Pure and Applied Mathematics*.
Academic Press, 1970.



O. El-Fallah, E. Fricain, K. Kellay, J. Mashregghi, and T. Ransford.
Constructive approximation in de branges–rovnyak spaces.
Constructive Approximation, 44(2):269–281, Oct 2016.

Bibliography II



W. Rudin.

Real and Complex Analysis.

Mathematics series. McGraw-Hill, 1987.



D. Sarason.

Sub-Hardy Hilbert spaces in the unit disk.

University of Arkansas lecture notes in the mathematical sciences.

Wiley, 1994.