

Section 6.2 Volume: The Disk and Washer Methods

$$1. V = \pi \int_0^1 (-x + 1)^2 dx = \pi \int_0^1 (x^2 - 2x + 1) dx = \pi \left[\frac{x^3}{3} - x^2 + x \right]_0^1 = \frac{\pi}{3}$$

$$2. V = \pi \int_0^2 (4 - x^2)^2 dx = \pi \int_0^2 (x^4 - 8x^2 + 16) dx = \pi \left[\frac{x^5}{5} - \frac{8x^3}{3} + 16x \right]_0^2 = \frac{256\pi}{15}$$

$$3. V = \pi \int_1^4 (\sqrt{x})^2 dx = \pi \int_1^4 x dx = \pi \left[\frac{x^2}{2} \right]_1^4 = \frac{15\pi}{2}$$

$$4. V = \pi \int_0^3 (\sqrt{9 - x^2})^2 dx = \pi \int_0^3 (9 - x^2) dx = \pi \left[9x - \frac{x^3}{3} \right]_0^3 = 18\pi$$

$$5. V = \pi \int_0^1 [(x^2)^2 - (x^5)^2] dx = \pi \int_0^1 (x^4 - x^{10}) dx = \pi \left[\frac{x^5}{5} - \frac{x^{11}}{11} \right]_0^1 = \pi \left(\frac{1}{5} - \frac{1}{11} \right) = \frac{6\pi}{55}$$

$$6. \quad 2 = 4 - \frac{x^2}{4}$$

$$8 = 16 - x^2$$

$$x^2 = 8$$

$$x = \pm 2\sqrt{2}$$

$$V = \pi \int_{-2\sqrt{2}}^{2\sqrt{2}} \left[\left(4 - \frac{x^2}{4} \right)^2 - (2)^2 \right] dx$$

$$= 2\pi \int_0^{2\sqrt{2}} \left[\frac{x^4}{16} - 2x^2 + 12 \right] dx$$

$$= 2\pi \left[\frac{x^5}{80} - \frac{2x^3}{3} + 12x \right]_0^{2\sqrt{2}}$$

$$= 2\pi \left[\frac{128\sqrt{2}}{80} - \frac{32\sqrt{2}}{3} + 24\sqrt{2} \right]$$

$$= \frac{448\sqrt{2}\pi}{15} \approx 132.69$$

$$7. \quad y = x^2 \Rightarrow x = \sqrt{y}$$

$$V = \pi \int_0^4 (\sqrt{y})^2 dy = \pi \int_0^4 y dy$$

$$= \pi \left[\frac{y^2}{2} \right]_0^4 = 8\pi$$

$$8. \quad y = \sqrt{16 - x^2} \Rightarrow x = \sqrt{16 - y^2}$$

$$V = \pi \int_0^4 (\sqrt{16 - y^2})^2 dy = \pi \int_0^4 (16 - y^2) dy$$

$$= \pi \left[16y - \frac{y^3}{3} \right]_0^4 = \frac{128\pi}{3}$$

$$9. \quad y = x^{2/3} \Rightarrow x = y^{3/2}$$

$$V = \pi \int_0^1 (y^{3/2})^2 dy = \pi \int_0^1 y^3 dy = \pi \left[\frac{y^4}{4} \right]_0^1 = \frac{\pi}{4}$$

$$10. \quad V = \pi \int_1^4 (-y^2 + 4y)^2 dy = \pi \int_1^4 (y^4 - 8y^3 + 16y^2) dy$$

$$= \pi \left[\frac{y^5}{5} - 2y^4 + \frac{16y^3}{3} \right]_1^4 = \frac{459\pi}{15} = \frac{153\pi}{5}$$

11. Because the axis of revolution is vertical, the integral should be found with respect to y , using

$$y = 4e^x \Rightarrow x = \ln \frac{y}{4} \text{ and } x = 1 \text{ on the interval } [4, 4e].$$

$$V = \pi \int_4^{4e} \left[1 - \left(\ln \frac{y}{4} \right)^2 \right] dy$$

12. Because you are integrating with respect to y , the limits of integration should be

$$y = 4e^0 = 4 \text{ and } y = 4e^1 = 4e.$$

$$V = \pi \int_4^{4e} \left[1 - \ln \left(\frac{y}{4} \right)^2 \right] dy$$

$$13. \quad y = \sqrt{x}, y = 0, x = 3$$


$$(a) \quad R(x) = \sqrt{x}, r(x) = 0$$

$$V = \pi \int_0^3 (\sqrt{x})^2 dx$$

$$= \pi \int_0^3 x dx$$

$$= \pi \left[\frac{x^2}{2} \right]_0^3$$

$$= \frac{9\pi}{2}$$



$$(b) \quad R(y) = 3, r(y) = y^2$$


$$V = \pi \int_0^{\sqrt{3}} [3^2 - (y^2)^2] dy$$

$$= \pi \int_0^{\sqrt{3}} (9 - y^4) dy$$

$$= \pi \left[9y - \frac{y^5}{5} \right]_0^{\sqrt{3}}$$


$$= \pi \left[9\sqrt{3} - \frac{9}{5}\sqrt{3} \right]$$

$$= \frac{36\sqrt{3}\pi}{5}$$




(c) $R(y) = 3 - y^2, r(y) = 0$

$$\begin{aligned} V &= \pi \int_0^{\sqrt{3}} (3 - y^2)^2 dy = \pi \int_0^{\sqrt{3}} (9 - 6y^2 + y^4) dy \\ &= \pi \left[9y - 2y^3 + \frac{y^5}{5} \right]_0^{\sqrt{3}} = \pi \left[9\sqrt{3} - 6\sqrt{3} + \frac{9\sqrt{3}}{5} \right] \\ &= \frac{24\sqrt{3}\pi}{5} \end{aligned}$$



(d) $R(y) = 3 + (3 - y^2) = 6 - y^2, r(y) = 3$


$$\begin{aligned} V &= \pi \int_0^{\sqrt{3}} [(6 - y^2)^2 - 3^2] dy = \pi \int_0^{\sqrt{3}} (y^4 - 12y^2 + 27) dy \\ &= \pi \left[\frac{y^5}{5} - 4y^3 + 27y \right]_0^{\sqrt{3}} = \pi \left[\frac{9\sqrt{3}}{5} - 12\sqrt{3} + 27\sqrt{3} \right] \\ &= \frac{84\sqrt{3}\pi}{5} \end{aligned}$$



14. $y = 2x^2, y = 0, x = 2$


(a) $R(y) = 2, r(y) = \sqrt{y/2}$

$$V = \pi \int_0^8 \left(2 - \frac{y}{2} \right) dy = \pi \left[4y - \frac{y^2}{4} \right]_0^8 = 16\pi$$




(b) $R(x) = 2x^2, r(x) = 0$

$$V = \pi \int_0^2 4x^4 dx = \pi \left[\frac{4x^5}{5} \right]_0^2 = \frac{128\pi}{5}$$




(c) $R(x) = 8, r(x) = 8 - 2x^2$

$$\begin{aligned} V &= \pi \int_0^2 [64 - (64 - 32x^2 + 4x^4)] dx \\ &= \pi \int_0^2 (32x^2 - 4x^4) dx = 4\pi \int_0^2 (8x^2 - x^4) dx \\ &= 4\pi \left[\frac{8}{3}x^3 - \frac{1}{5}x^5 \right]_0^2 \\ &= \frac{896\pi}{15} \end{aligned}$$



(d) $R(y) = 2 - \sqrt{y/2}, r(y) = 0$

$$\begin{aligned} V &= \pi \int_0^8 \left(2 - \sqrt{\frac{y}{2}} \right) dy \\ &= \pi \int_0^8 \left(4 - 4\sqrt{\frac{y}{2}} + \frac{y}{2} \right) dy \\ &= \pi \left[4y - \frac{4\sqrt{2}}{3} y^{3/2} + \frac{y^2}{4} \right]_0^8 = \frac{16\pi}{3} \end{aligned}$$





15. $y = x^2$, $y = 4x - x^2$ intersect at $(0, 0)$ and $(2, 4)$.

(a) $R(x) = 4x - x^2$, $r(x) = x^2$

$$\begin{aligned} V &= \pi \int_0^2 [(4x - x^2)^2 - x^4] dx \\ &= \pi \int_0^2 (16x^2 - 8x^3) dx \\ &= \pi \left[\frac{16}{3}x^3 - 2x^4 \right]_0^2 = \frac{32\pi}{3} \end{aligned}$$

(b) $R(x) = 6 - x^2$, $r(x) = 6 - (4x - x^2)$


$$\begin{aligned} V &= \pi \int_0^2 [(6 - x^2)^2 - (6 - 4x + x^2)^2] dx \\ &= 8\pi \int_0^2 (x^3 - 5x^2 + 6x) dx \\ &= 8\pi \left[\frac{x^4}{4} - \frac{5}{3}x^3 + 3x^2 \right]_0^2 = \frac{64\pi}{3} \end{aligned}$$



16. $y = 4 + 2x - x^2$, $y = 4 - x$ intersect at $(0, 4)$ and $(3, 1)$.


(a) $R(x) = 4 + 2x - x^2$, $r(x) = 4 - x$

$$\begin{aligned} V &= \pi \int_0^3 [(4 + 2x - x^2)^2 - (4 - x)^2] dx \\ &= \pi \int_0^3 (x^4 - 4x^3 - 5x^2 + 24x) dx \\ &= \pi \left[\frac{x^5}{5} - x^4 - \frac{5x^3}{3} + 12x^2 \right]_0^3 = \frac{153\pi}{5} \end{aligned}$$




(b) $R(x) = (4 + 2x - x^2) - 1$, $r(x) = (4 - x) - 1$

$$\begin{aligned} V &= \pi \int_0^3 [(3 + 2x - x^2)^2 - (3 - x)^2] dx \\ &= \pi \int_0^3 (x^4 - 4x^3 - 3x^2 + 18x) dx \\ &= \pi \left[\frac{x^5}{5} - x^4 - x^3 + 9x^2 \right]_0^3 = \frac{108\pi}{5} \end{aligned}$$




17. $R(x) = 4 - x$, $r(x) = 1$

$$\begin{aligned} V &= \pi \int_0^3 [(4 - x)^2 - (1)^2] dx \\ &= \pi \int_0^3 (x^2 - 8x + 15) dx \\ &= \pi \left[\frac{x^3}{3} - 4x^2 + 15x \right]_0^3 \\ &= 18\pi \end{aligned}$$




18. $R(x) = 4 - \frac{x^3}{2}$, $r(x) = 0$

$$\begin{aligned} V &= \pi \int_0^1 \left(4 - \frac{x^3}{2} \right)^2 dx \\ &= \pi \int_0^1 \left[16 - 4x^3 + \frac{x^6}{4} \right] dx \\ &= \pi \left[16x - x^4 + \frac{x^7}{28} \right]_0^1 \\ &= \pi \left(32 - 16 + \frac{128}{28} \right) \\ &= \frac{144}{7}\pi \end{aligned}$$




$$19. R(x) = 4, r(x) = 4 - \frac{3}{1+x}$$

$$\begin{aligned} V &= \pi \int_0^3 \left[4^2 - \left(4 - \frac{3}{1+x} \right)^2 \right] dx \\ &= \pi \int_0^3 \left[\frac{24}{1+x} - \frac{9}{(1+x)^2} \right] dx \\ &= \pi \left[24 \ln|1+x| + \frac{9}{1+x} \right]_0^3 \\ &= \pi \left[\left(24 \ln 4 + \frac{9}{4} \right) - 9 \right] \\ &= \left(48 \ln 2 - \frac{27}{4} \right) \pi \approx 83.318 \end{aligned}$$




$$20. R(x) = 4, r(x) = 4 - \sec x$$

$$\begin{aligned} V &= \pi \int_0^{\pi/3} \left[(4)^2 - (4 - \sec x)^2 \right] dx \\ &= \pi \int_0^{\pi/3} (8 \sec x - \sec^2 x) dx \\ &= \pi \left[8 \ln|\sec x + \tan x| - \tan x \right]_0^{\pi/3} \\ &= \pi \left[(8 \ln|2 + \sqrt{3}| - \sqrt{3}) - (8 \ln|1 + 0| - 0) \right] \\ &= \pi \left[8 \ln(2 + \sqrt{3}) - \sqrt{3} \right] \approx 27.66 \end{aligned}$$



$$21. R(y) = 5 - y, r(y) = 0$$


$$\begin{aligned} V &= \pi \int_0^4 (5 - y)^2 dy \\ &= \pi \int_0^4 (25 - 10y + y^2) dy \\ &= \pi \left[25y - 5y^2 + \frac{y^3}{3} \right]_0^4 \\ &= \pi \left[100 - 80 + \frac{64}{3} \right] \\ &= \frac{124\pi}{3} \end{aligned}$$



$$22. y = 3 - x, x = 3 - y$$


$$R(y) = 5, r(y) = 5 - (3 - y) = 2 + y$$

$$\begin{aligned} V &= \pi \int_0^2 \left[5^2 - (2 + y)^2 \right] dy \\ &= \pi \int_0^2 (-y^2 - 4y + 21) dy \\ &= \pi \left[-\frac{y^3}{3} - 2y^2 + 21y \right]_0^2 = \frac{94\pi}{3} \end{aligned}$$



$$23. R(y) = 5 - y^2, r(y) = 1$$


$$\begin{aligned} V &= \pi \int_{-2}^2 \left[(5 - y^2)^2 - 1 \right] dy \\ &= 2\pi \int_0^2 \left[y^4 - 10y^2 + 24 \right] dy \\ &= 2\pi \left[\frac{y^5}{5} - \frac{10y^3}{3} + 24y \right]_0^2 \\ &= 2\pi \left[\frac{32}{5} - \frac{80}{3} + 48 \right] = \frac{832\pi}{15} \end{aligned}$$



$$24. xy = 3, x = \frac{3}{y}$$


$$R(y) = 5 - \frac{3}{y}, r(y) = 0$$

$$\begin{aligned} V &= \pi \int_1^4 \left(5 - \frac{3}{y} \right)^2 dy \\ &= \pi \int_1^4 \left(25 + \frac{9}{y^2} - \frac{30}{y} \right) dy \\ &= \pi \left[25y - \frac{9}{y} - 30 \ln y \right]_1^4 \\ &= \pi \left[\left(100 - \frac{9}{4} - 30 \ln 4 \right) - (25 - 9) \right] \\ &= \pi \left(\frac{327}{4} - 30 \ln 4 \right) \approx 126.17 \end{aligned}$$




$$25. R(x) = \frac{1}{\sqrt{x+1}}, r(x) = 0$$

$$\begin{aligned} V &= \pi \int_0^4 \left(\frac{1}{\sqrt{x+1}} \right)^2 dx \\ &= \pi \int_0^4 \frac{1}{x+1} dx \\ &= \pi [\ln|x+1|]_0^4 \\ &= \pi \ln 5 \end{aligned}$$




$$26. R(x) = x\sqrt{4-x^2}, r(x) = 0$$

$$\begin{aligned} V &= 2\pi \int_0^2 (x\sqrt{4-x^2})^2 dx \\ &= 2\pi \int_0^2 (4x^2 - x^4) dx \\ &= 2\pi \left[\frac{4x^3}{3} - \frac{x^5}{5} \right]_0^2 \\ &= 2\pi \left[\frac{32}{3} - \frac{32}{5} \right] = \frac{128\pi}{15} \end{aligned}$$




$$27. R(x) = \frac{1}{x}, r(x) = 0$$

$$\begin{aligned} V &= \pi \int_1^3 \left(\frac{1}{x} \right)^2 dx \\ &= \pi \left[-\frac{1}{x} \right]_1^3 \\ &= \pi \left[-\frac{1}{3} + 1 \right] = \frac{2}{3}\pi \end{aligned}$$




$$28. R(x) = \frac{2}{x+1}, r(x) = 0$$

$$\begin{aligned} V &= \pi \int_0^6 \left(\frac{2}{x+1} \right)^2 dx \\ &= 4\pi \int_0^6 (x+1)^{-2} dx \\ &= 4\pi \left[\frac{-1}{x+1} \right]_0^6 \\ &= 4\pi \left[-\frac{1}{7} + 1 \right] = \frac{24\pi}{7} \end{aligned}$$




$$29. R(x) = e^{-x}, r(x) = 0$$

$$\begin{aligned} V &= \pi \int_0^1 (e^{-x})^2 dx \\ &= \pi \int_0^1 e^{-2x} dx \\ &= \left[-\frac{\pi}{2} e^{-2x} \right]_0^1 \\ &= \frac{\pi}{2} (1 - e^{-2}) \approx 1.358 \end{aligned}$$



$$30. R(x) = e^{x/4}, r(x) = 0$$

$$\begin{aligned} V &= \pi \int_0^6 (e^{x/4})^2 dx \\ &= \pi \int_0^6 e^{x/2} dx \\ &= \pi [2e^{x/2}]_0^6 \\ &= \pi (2e^3 - 2) \approx 119.92 \end{aligned}$$



$$31. \quad x^2 + 1 = -x^2 + 2x + 5$$


$$2x^2 - 2x - 4 = 0$$

$$x^2 - x - 2 = 0$$

$$(x - 2)(x + 1) = 0$$

The curves intersect at $(-1, 2)$ and $(2, 5)$.

$$\begin{aligned} V &= \pi \int_{-1}^2 \left[(5 + 2x - x^2)^2 - (x^2 + 1)^2 \right] dx + \pi \int_2^5 \left[(x^2 + 1)^2 - (5 + 2x - x^2)^2 \right] dx \\ &= \pi \int_{-1}^2 (-4x^3 - 8x^2 + 20x + 24) dx + \pi \int_2^5 (4x^3 + 8x^2 - 20x - 24) dx \\ &= \pi \left[-x^4 - \frac{8}{3}x^3 + 10x^2 + 24x \right]_{-1}^2 + \pi \left[x^4 + \frac{8}{3}x^3 - 10x^2 - 24x \right]_2^5 \\ &= \pi \frac{152}{3} + \pi \frac{125}{3} = \frac{277\pi}{3} \end{aligned}$$



$$32. \quad \sqrt{x} = -\frac{1}{2}x + 4$$


$$x = \frac{1}{4}x^2 - 4x + 16$$

$$0 = x^2 - 20x + 64$$

$$0 = (x - 4)(x - 16)$$


The curves intersect at $(4, 2)$. (Note $x = 16$ is an extraneous root.)

$$\begin{aligned} V &= \pi \int_0^4 \left[\left(4 - \frac{1}{2}x\right)^2 - (\sqrt{x})^2 \right] dx + \pi \int_4^8 \left[(\sqrt{x})^2 - \left(4 - \frac{1}{2}x\right)^2 \right] dx \\ &= \pi \int_0^4 \left(\frac{x^2}{4} - 5x + 16 \right) dx + \pi \int_4^8 \left(-\frac{x^2}{4} + 5x - 16 \right) dx \\ &= \pi \left[\frac{x^3}{12} - \frac{5x^2}{2} + 16x \right]_0^4 + \pi \left[-\frac{x^3}{12} + \frac{5x^2}{2} - 16x \right]_4^8 \\ &= \frac{88}{3}\pi + \frac{56}{3}\pi = 48\pi \end{aligned}$$



$$33. \quad y = 6 - 3x \Rightarrow x = \frac{1}{3}(6 - y)$$


$$\begin{aligned} V &= \pi \int_0^6 \left[\frac{1}{3}(6 - y) \right]^2 dy \\ &= \frac{\pi}{9} \int_0^6 [36 - 12y + y^2] dy \\ &= \frac{\pi}{9} \left[36y - 6y^2 + \frac{y^3}{3} \right]_0^6 \\ &= \frac{\pi}{9} \left[216 - 216 + \frac{216}{3} \right] \\ &= 8\pi \end{aligned}$$



$$34. \quad y = 9 - x^2, y = 0, x = 2, x = 3$$


$$x = \sqrt{9 - y}$$

$$\begin{aligned} V &= \pi \int_0^5 \left[(\sqrt{9 - y})^2 - 2^2 \right] dy \\ &= \pi \int_0^5 (5 - y) dy \\ &= \pi \left[5y - \frac{y^2}{2} \right]_0^5 = \pi \left(25 - \frac{25}{2} \right) = \frac{25\pi}{2} \end{aligned}$$




$$\begin{aligned}
 35. V &= \pi \int_0^{\pi} (\sin x)^2 dx \\
 &= \pi \int_0^{\pi} \frac{1 - \cos 2x}{2} dx \\
 &= \frac{\pi}{2} \left[x - \frac{1}{2} \sin 2x \right]_0^{\pi} = \frac{\pi}{2} [\pi] = \frac{\pi^2}{2}
 \end{aligned}$$

Numerical approximation: 4.9348



$$\begin{aligned}
 36. V &= \pi \int_0^{\pi/4} \cos^2 2x dx \\
 &= \pi \int_0^{\pi/4} \frac{1 + \cos 4x}{2} dx \\
 &= \frac{\pi}{2} \left[x + \frac{\sin 4x}{4} \right]_0^{\pi/4} \\
 &= \frac{\pi}{2} \left[\frac{\pi}{4} \right] = \frac{\pi^2}{8}
 \end{aligned}$$

Numerical approximation: 1.2337



$$\begin{aligned}
 37. V &= \pi \int_1^2 (e^{x-1})^2 dx \\
 &= \pi \int_1^2 e^{2x-2} dx \\
 &= \frac{\pi}{2} e^{2x-2} \Big|_1^2 \\
 &= \frac{\pi}{2} (e^2 - 1)
 \end{aligned}$$

Numerical approximation: 10.0359

$$\begin{aligned}
 38. V &= \pi \int_{-1}^2 [e^{x/2} + e^{-x/2}]^2 dx \\
 &= \pi \int_{-1}^2 [e^x + e^{-x} + 2] dx \\
 &= \pi [e^x - e^{-x} + 2x]_{-1}^2 \\
 &= \pi [(e^2 - e^{-2} + 4) - (e^{-1} - e - 2)] \\
 &= \pi (e^2 + e + 6 - e^{-2} - e^{-1})
 \end{aligned}$$


Numerical approximation: 49.0218

$$39. V = \pi \int_0^2 [e^{-x^2}]^2 dx \approx 1.9686$$

$$40. V = \pi \int_1^3 [\ln x]^2 dx \approx 3.2332$$

$$41. V = \pi \int_0^5 [2 \arctan(0.2x)]^2 dx \approx 15.4115$$

$$\begin{aligned}
 42. x^2 &= \sqrt{2x} \\
 x^4 &= 2x \\
 x^3 &= 2 \\
 x &= 2^{1/3} \approx 1.2599 \\
 V &= \pi \int_0^{2^{1/3}} [(\sqrt{2x})^2 - (x^2)^2] dx \approx 2.9922
 \end{aligned}$$



$$43. V = \pi \int_0^1 y^2 dy = \pi \frac{y^3}{3} \Big|_0^1 = \frac{\pi}{3}$$

$$\begin{aligned}
 44. V &= \pi \int_0^1 [1^2 - (1-y)^2] dy \\
 &= \pi \int_0^1 [2y - y^2] dy \\
 &= \pi \left[y^2 - \frac{y^3}{3} \right]_0^1 \\
 &= \pi \left(1 - \frac{1}{3} \right) = \frac{2}{3} \pi
 \end{aligned}$$

$$\begin{aligned}
 45. V &= \pi \int_0^1 (x^2 - x^4) dx \\
 &= \pi \left[\frac{x^3}{3} - \frac{x^5}{5} \right]_0^1 \\
 &= \pi \left(\frac{1}{3} - \frac{1}{5} \right) \\
 &= \frac{2\pi}{15}
 \end{aligned}$$

$$\begin{aligned}
 46. V &= \pi \int_0^1 [(1-x^2)^2 - (1-x)^2] dx \\
 &= \pi \int_0^1 [1 - 2x^2 + x^4 - 1 + 2x - x^2] dx \\
 &= \pi \int_0^1 [2x - 3x^2 + x^4] dx \\
 &= \pi \left[x^2 - x^3 + \frac{x^5}{5} \right]_0^1 \\
 &= \pi \left(\frac{1}{5} \right) = \frac{\pi}{5}
 \end{aligned}$$


$$\begin{aligned}
 47. V &= \pi \int_0^1 (1-y) dy \\
 &= \pi \left[y - \frac{y^2}{2} \right]_0^1 = \pi \left(1 - \frac{1}{2} \right) = \frac{\pi}{2}
 \end{aligned}$$

$$\begin{aligned}
 48. V &= \pi \int_0^1 (1 - \sqrt{y})^2 dy \\
 &= \pi \int_0^1 (1 - 2\sqrt{y} + y) dy \\
 &= \pi \left[y - \frac{4}{3}y^{3/2} + \frac{y^2}{2} \right]_0^1 \\
 &= \pi \left(1 - \frac{4}{3} + \frac{1}{2} \right) \\
 &= \frac{\pi}{6}
 \end{aligned}$$


$$\begin{aligned}
 49. V &= \pi \int_0^1 (y - y^2) dy \\
 &= \pi \left[\frac{y^2}{2} - \frac{y^3}{3} \right]_0^1 = \pi \left(\frac{1}{2} - \frac{1}{3} \right) = \frac{\pi}{6}
 \end{aligned}$$

$$\begin{aligned}
 50. V &= \pi \int_0^1 [(1-y)^2 - (1-\sqrt{y})^2] dy \\
 &= \pi \int_0^1 [1 - 2y + y^2 - 1 + 2\sqrt{y} - y] dy \\
 &= \pi \int_0^1 [2\sqrt{y} - 3y + y^2] dy \\
 &= \pi \left[\frac{4}{3}y^{3/2} - \frac{3y^2}{2} + \frac{y^3}{3} \right]_0^1 \\
 &= \pi \left(\frac{4}{3} - \frac{3}{2} + \frac{1}{3} \right) \\
 &= \frac{\pi}{6}
 \end{aligned}$$



51. $\pi \int_0^{\pi/2} \sin^2 x \, dx$ represents the volume of the solid generated by revolving the region bounded by $y = \sin x$, $y = 0$, $x = 0$, $x = \pi/2$ about the x -axis.



52. $\pi \int_2^4 y^4 \, dy$ represents the volume of the solid generated by revolving the region bounded by $x = y^2$, $x = 0$, $y = 2$, and $y = 4$ about the y -axis.



53.



The volumes are the same because the solid has been translated horizontally. $(4x - x^2 = 4 - (x-2)^2)$

54. (a) Matches (ii) because the axis of rotation is vertical, and this is the washer method.
 (b) Matches (iv) because the axis of rotation is horizontal, and this is the washer method.
 (c) Matches (i) because the axis of rotation is horizontal.
 (d) Matches (iii) because the axis of rotation is vertical.

$$55. V = \pi \int_0^4 (\sqrt{x})^2 dx = \pi \int_0^4 x dx = \left[\frac{\pi x^2}{2} \right]_0^4 = 8\pi$$

Let $0 < c < 4$ and set


$$\pi \int_0^c x dx = \left[\frac{\pi x^2}{2} \right]_0^c = \frac{\pi c^2}{2} = 4\pi.$$

$$c^2 = 8$$

$$c = \sqrt{8} = 2\sqrt{2}$$

So, when $x = 2\sqrt{2}$, the solid is divided into two parts of equal volume.

$$\begin{aligned} 57. V &= \pi \int_{-\sqrt{R^2-r^2}}^{\sqrt{R^2-r^2}} \left[(\sqrt{R^2-x^2})^2 - r^2 \right] dx \\ &= 2\pi \int_0^{\sqrt{R^2-r^2}} (R^2 - r^2 - x^2) dx \\ &= 2\pi \left[(R^2 - r^2)x - \frac{x^3}{3} \right]_0^{\sqrt{R^2-r^2}} \\ &= 2\pi \left[(R^2 - r^2)^{3/2} - \frac{(R^2 - r^2)^{3/2}}{3} \right] = \frac{4}{3}\pi (R^2 - r^2)^{3/2} \end{aligned}$$



58. Let $R = 6$ in the previous Exercise.

$$\frac{4}{3}\pi (36 - r^2)^{3/2} = \frac{1}{2}\left(\frac{4}{3}\right)\pi (6)^3$$

$$(36 - r^2)^{3/2} = 108$$

$$36 - r^2 = (108)^{2/3}$$

$$r^2 = 36 - 108^{2/3}$$



$$r = \sqrt{36 - 108^{2/3}} \approx 3.65$$

$$60. R(x) = \sqrt{r^2 - x^2}, r(x) = 0$$

$$\begin{aligned} V &= \pi \int_{-r}^r (r^2 - x^2) dx \\ &= 2\pi \int_0^r (r^2 - x^2) dx \\ &= 2\pi \left[r^2x - \frac{1}{3}x^3 \right]_0^r \\ &= 2\pi \left(r^3 - \frac{1}{3}r^3 \right) = \frac{4}{3}\pi r^3 \end{aligned}$$


$$59. R(x) = \frac{r}{h}x, r(x) = 0$$

$$V = \pi \int_0^h \frac{r^2}{h^2}x^2 dx = \left[\frac{r^2\pi}{3h^2}x^3 \right]_0^h = \frac{r^2\pi}{3h^2}h^3 = \frac{1}{3}\pi r^2h$$





$$61. x = r - \frac{r}{H}y = r\left(1 - \frac{y}{H}\right), R(y) = r\left(1 - \frac{y}{H}\right), r(y) = 0$$

$$\begin{aligned} V &= \pi \int_0^h \left[r\left(1 - \frac{y}{H}\right) \right]^2 dy = \pi r^2 \int_0^h \left(1 - \frac{2}{H}y + \frac{1}{H^2}y^2 \right) dy \\ &= \pi r^2 \left[y - \frac{1}{H}y^2 + \frac{1}{3H^2}y^3 \right]_0^h \\ &= \pi r^2 \left(h - \frac{h^2}{H} + \frac{h^3}{3H^2} \right) = \pi r^2 h \left(1 - \frac{h}{H} + \frac{h^2}{3H^2} \right) \end{aligned}$$



$$62. x = \sqrt{r^2 - y^2}, R(y) = \sqrt{r^2 - y^2}, r(y) = 0$$


$$\begin{aligned} V &= \pi \int_h^r (\sqrt{r^2 - y^2})^2 dy \\ &= \pi \int_h^r (r^2 - y^2) dy \\ &= \pi \left[r^2 y - \frac{y^3}{3} \right]_h^r \\ &= \pi \left[\left(r^3 - \frac{r^3}{3} \right) - \left(r^2 h - \frac{h^3}{3} \right) \right] \\ &= \pi \left(\frac{2r^3}{3} - r^2 h + \frac{h^3}{3} \right) = \frac{\pi}{3} (2r^3 - 3r^2 h + h^3) \end{aligned}$$



$$V = \pi \int_0^2 \left(\frac{1}{8} x^2 \sqrt{2-x} \right)^2 dx = \frac{\pi}{64} \int_0^2 x^4 (2-x) dx = \frac{\pi}{64} \left[\frac{2x^5}{5} - \frac{x^6}{6} \right]_0^2 = \frac{\pi}{30} \text{ m}^3$$


$$64. y = \begin{cases} \sqrt{0.1x^3 - 2.2x^2 + 10.9x + 22.2}, & 0 \leq x \leq 11.5 \\ 2.95, & 11.5 < x \leq 15 \end{cases}$$

$$\begin{aligned} V &= \pi \int_0^{11.5} \left(\sqrt{0.1x^3 - 2.2x^2 + 10.9x + 22.2} \right)^2 dx + \pi \int_{11.5}^{15} 2.95^2 dx \\ &= \pi \left[\frac{0.1x^4}{4} - \frac{2.2x^3}{3} + \frac{10.9x^2}{2} + 22.2x \right]_0^{11.5} + \pi [2.95^2 x]_{11.5}^{15} \\ &\approx 1031.9016 \text{ cubic centimeters} \end{aligned}$$




$$65. (a) R(x) = \frac{3}{5}\sqrt{25 - x^2}, r(x) = 0$$

$$\begin{aligned} V &= \frac{9\pi}{25} \int_{-5}^5 (25 - x^2) dx \\ &= \frac{18\pi}{25} \int_0^5 (25 - x^2) dx \\ &= \frac{18\pi}{25} \left[25x - \frac{x^3}{3} \right]_0^5 \\ &= 60\pi \end{aligned}$$



$$(b) R(y) = \frac{5}{3}\sqrt{9 - y^2}, r(y) = 0, x \geq 0$$

$$\begin{aligned} V &= \frac{25\pi}{9} \int_0^3 (9 - y^2) dy \\ &= \frac{25\pi}{9} \left[9y - \frac{y^3}{3} \right]_0^3 \\ &= 50\pi \end{aligned}$$



$$66. \text{ Total volume: } V = \frac{4\pi(50)^3}{3} = \frac{500,000\pi}{3} \text{ ft}^3$$

Volume of water in the tank:

$$\pi \int_{-y_0}^{y_0} (\sqrt{2500 - y^2})^2 dy = \pi \int_{-y_0}^{y_0} (2500 - y^2) dy = \pi \left[2500y - \frac{y^3}{3} \right]_{-y_0}^{y_0} = \pi \left(2500y_0 - \frac{y_0^3}{3} + \frac{250,000}{3} \right)$$

When the tank is one-fourth of its capacity:

$$\frac{1}{4} \left(\frac{500,000\pi}{3} \right) = \pi \left(2500y_0 - \frac{y_0^3}{3} + \frac{250,000}{3} \right)$$


$$125,000 = 7500y_0 - y_0^3 + 250,000$$

$$y_0^3 - 7500y_0 - 125,000 = 0$$

$$y_0 \approx -17.36$$

$$\text{Depth: } -17.36 - (-50) = 32.64 \text{ feet}$$

When the tank is three-fourths of its capacity the depth is $100 - 32.64 = 67.36$ feet.



67. (a) First find where $y = b$ intersects the parabola:

$$b = 4 - \frac{x^2}{4}$$

$$x^2 = 16 - 4b = 4(4 - b)$$

$$x = 2\sqrt{4 - b}$$


$$V = \int_0^{2\sqrt{4-b}} \pi \left[4 - \frac{x^2}{4} - b \right]^2 dx + \int_{2\sqrt{4-b}}^4 \pi \left[b - 4 + \frac{x^2}{4} \right]^2 dx$$

$$= \int_0^4 \pi \left[4 - \frac{x^2}{4} - b \right]^2 dx$$


$$= \pi \int_0^4 \left[\frac{x^4}{16} - 2x^2 + \frac{bx^2}{2} + b^2 - 8b + 16 \right] dx$$

$$= \pi \left[\frac{x^5}{80} - \frac{2x^3}{3} + \frac{bx^3}{6} + b^2x - 8bx + 16x \right]_0^4$$

$$= \pi \left(\frac{64}{5} - \frac{128}{3} + \frac{32}{3}b + 4b^2 - 32b + 64 \right) = \pi \left(4b^2 - \frac{64}{3}b + \frac{512}{15} \right)$$



(b) Graph of $V(b) = \pi\left(4b^2 - \frac{64}{3}b + \frac{512}{15}\right)$




Minimum volume is 17.87 for $b = 2.67$.

(c) $V'(b) = \pi\left(8b - \frac{64}{3}\right) = 0 \Rightarrow b = \frac{64/3}{8} = \frac{8}{3} = 2\frac{2}{3}$

$V''(b) = 8\pi > 0 \Rightarrow b = \frac{8}{3}$ is a relative minimum.


68. (a) $\pi \int_0^h r^2 dx$ (ii)

is the volume of a right circular cylinder with radius r and height h .




(b) $\pi \int_{-b}^b \left(a\sqrt{1 - \frac{x^2}{b^2}}\right)^2 dx$ (iv)

is the volume of an ellipsoid with axes $2a$ and $2b$.




(c) $\pi \int_{-r}^r \left(\sqrt{r^2 - x^2}\right)^2 dx$ (iii)

is the volume of a sphere with radius r .




(d) $\pi \int_0^h \left(\frac{rx}{h}\right)^2 dx$ (i)

is the volume of a right circular cone with the radius of the base as r and height h .



(e) $\pi \int_{-r}^r \left[\left(R + \sqrt{r^2 - x^2}\right)^2 - \left(R - \sqrt{r^2 - x^2}\right)^2 \right] dx$ (v)

is the volume of a torus with the radius of its circular cross section as r and the distance from the axis of the torus to the center of its cross section as R .




69. Let $A_1(x)$ and $A_2(x)$ equal the areas of the cross sections of the two solids for $a \leq x \leq b$. Because $A_1(x) = A_2(x)$, you have

$$V_1 = \int_a^b A_1(x) dx = \int_a^b A_2(x) dx = V_2.$$

So, the volumes are the same.


70.



$$\text{Base of cross section} = (x + 1) - (x^2 - 1) = 2 + x - x^2$$


$$(a) \quad A(x) = b^2 = (2 + x - x^2)^2 = 4 + 4x - 3x^2 - 2x^3 + x^4$$

$$V = \int_{-1}^2 (4 + 4x - 3x^2 - 2x^3 + x^4) dx = \left[4x + 2x^2 - x^3 - \frac{1}{2}x^4 + \frac{1}{5}x^5 \right]_{-1}^2 = \frac{81}{10}$$




$$(b) \quad A(x) = bh = (2 + x - x^2)1$$

$$V = \int_{-1}^2 (2 + x - x^2) dx = \left[2x + \frac{x^2}{2} - \frac{x^3}{3} \right]_{-1}^2 = \frac{9}{2}$$




71.



$$\text{Base of cross section} = 2\sqrt{4 - x^2}$$


$$(a) \quad A(x) = b^2 = (2\sqrt{4 - x^2})^2$$

$$\begin{aligned} V &= \int_{-2}^2 4(4 - x^2) dx \\ &= 4 \left[4x - \frac{x^3}{3} \right]_{-2}^2 \\ &= \frac{128}{3} \end{aligned}$$




$$(b) \quad A(x) = \frac{1}{2}bh = \frac{1}{2}(2\sqrt{4 - x^2})(\sqrt{3}\sqrt{4 - x^2}) = \sqrt{3}(4 - x^2)$$

$$\begin{aligned} V &= \sqrt{3} \int_{-2}^2 (4 - x^2) dx \\ &= \sqrt{3} \left[4x - \frac{x^3}{3} \right]_{-2}^2 \\ &= \frac{32\sqrt{3}}{3} \end{aligned}$$




$$(c) A(x) = \frac{1}{2}\pi r^2 = \frac{\pi}{2}(\sqrt{4-x^2})^2 = \frac{\pi}{2}(4-x^2)$$

$$\begin{aligned} V &= \frac{\pi}{2} \int_{-2}^2 (4-x^2) dx \\ &= \frac{\pi}{2} \left[4x - \frac{x^3}{3} \right]_{-2}^2 \\ &= \frac{16\pi}{3} \end{aligned}$$




$$(d) A(x) = \frac{1}{2}bh = \frac{1}{2}(2\sqrt{4-x^2})(\sqrt{4-x^2}) = 4-x^2$$

$$\begin{aligned} V &= \int_{-2}^2 (4-x^2) dx \\ &= \left[4x - \frac{x^3}{3} \right]_{-2}^2 \\ &= \frac{32}{3} \end{aligned}$$



72. (a) When $a = 1$: $|x| + |y| = 1$ represents a square.

When $a = 2$: $|x|^2 + |y|^2 = 1$ represents a circle.



$$(b) |y| = (1 - |x|^a)^{1/a}$$


$$A = 2 \int_{-1}^1 (1 - |x|^a)^{1/a} dx = 4 \int_0^1 (1 - x^a)^{1/a} dx$$

Solve the equation for $|y|$, as shown below. Then form n slices, each of whose area is approximated by the integral below. Finally, sum the volumes of these n slices.

$$|y| = (1 - |x|^a)^{1/a}$$

$$A = 2 \int_{-1}^1 (1 - |x|^a)^{1/a} dx = 4 \int_0^1 (1 - x^a)^{1/a} dx$$

73. For each axis of revolution, find the respective volume of each region.



$$x\text{-axis: } V = \pi \int_0^3 [9^2 - (x^2)^2] dx = \pi \int_0^3 (81 - x^4) dx = \pi \left[81x - \frac{1}{5}x^5 \right]_0^3 = \frac{972}{5}\pi = 194.4\pi$$


$$y\text{-axis: } V = \pi \int_0^9 (\sqrt{y})^2 dy = \pi \int_0^9 y dy = \pi \left[\frac{1}{2}y^2 \right]_0^9 = \frac{81}{2}\pi = 40.5\pi$$

$$\begin{aligned} x=3: V &= \pi \int_0^9 [(3)^2 - (3 - \sqrt{y})^2] dy = \pi \int_0^9 [9 - (9 - 6\sqrt{y} + y)] dy \\ &= \pi \int_0^9 (6\sqrt{y} - y) dy = \pi \left[4y^{3/2} - \frac{1}{2}y^2 \right]_0^9 = \frac{135}{2}\pi \approx 67.5\pi \end{aligned}$$

The resulting solids about each axis from least to greatest are y -axis, $x = 3$, and x -axis.

So, the answer is C.

74.



Because $y = 4$ is the axis of revolution, integrate with respect to x and the limits of integration are $x = 0$ and $x = \frac{3}{4}$.

$$R(x) = 4 - 4x^2 \text{ and } r(x) = 4 - 3x$$

$$V = \int_0^{3/4} \left[(4 - x^2)^2 - (4 - 3x)^2 \right] dx$$

So, the answer is B.

75. (a) The graphs intersect at $(0, 2)$, $(0.81, 1.4)$, and $(1.76, 2.72)$.

$$\begin{aligned} A &= \int_0^{0.81} [f(x) - g(x)] dx + \int_{0.81}^{1.76} [g(x) - f(x)] dx \\ &= \int_0^{0.81} [x^4 - 4x^2 + 2x + \sin(\pi x)] dx + \int_{0.81}^{1.76} [-\sin(\pi x) - x^4 + 4x^2 - 2x] dx \\ &= \left[\frac{1}{5}x^5 - \frac{4}{3}x^3 + x^2 - \frac{1}{\pi} \cos(\pi x) \right]_0^{0.81} + \left[\frac{1}{\pi} \cos(\pi x) - \frac{1}{5}x^5 + \frac{4}{3}x^3 - x^2 \right]_{0.81}^{1.76} \\ &\approx 0.5988 + 1.3066 \\ &\approx 1.905 \end{aligned}$$

$$\begin{aligned} \text{(b) Base of cross section: } g(x) - f(x) &= [2 - \sin(\pi x)] - (x^4 - 4x^2 + 2x + 2) \\ &= -\sin(\pi x) - x^4 + 4x^2 - 2x \end{aligned}$$

$$\begin{aligned} V &= \int_{0.807}^{1.757} [-\sin(\pi x) - x^4 + 4x^2 - 2x]^2 dx \\ &= 2.222 \end{aligned}$$

So, the volume of the solid is 2.222 units³.

$$\text{(c) } h(x) = g(x) - f(x)$$

$$\frac{dh}{dx} = g'(x) - f'(x) = [-\pi \cos(\pi x)] - (4x^3 - 8x + 2) = -\pi \cos(\pi x) - 4x^3 + 8x - 2$$

$$\text{So, } h'(1.2) = -\pi \cos(\pi \cdot 1.2) - 4(1.2)^3 + 8(1.2) - 2 \approx 3.230.$$