



UNSW
A U S T R A L I A



UNIVERSITY OF NEW SOUTH WALES

SCHOOL OF MATHEMATICS AND STATISTICS

Assignment

Number Theory

Author:
Adam J. Gray

Student Number:
3329798

Question 1

Part a

Use the character table given in lectures for \mathbb{Z}_5 , extended to a Dirichlet character, to evaluate

$$\sum_{i=1}^4 \chi_i(n) \overline{\chi_i(b)}, \quad \text{for each } b \in \mathbb{U}_5.$$

Part b

Use the results of (a) to prove, in detail, that there are infinitely many primes congruent to 1 mod 5, 2 mod 5, and 3 mod 5 and 4 mod 5.

Solution

For this question we use the following character table:

	1	3	4	2
χ_1	1	1	1	1
χ_2	1	-1	1	-1
χ_3	1	i	-1	-i
χ_4	1	-i	-1	i

Part a

$$\sum_{i=1}^4 \chi_i(n) \overline{\chi_i(b)} = \begin{cases} 0 & \text{if } n \not\equiv b \pmod{5} \\ 4 & \text{if } n \equiv b \pmod{5} \end{cases}$$

This follows immediately from the orthogonality relation proved in lectures.

Part b

Firstly see that

$$\begin{aligned} L(s, \chi_i) &= \sum_{n=1}^{\infty} \frac{\chi_i(n)}{n^s} \\ &= \prod_{p \text{ prime}} \left(1 - \frac{\chi_i(p)}{p^s} \right)^{-1} \end{aligned}$$

and so

$$\begin{aligned}
 \log L(s, \chi_i) &= - \sum_{n=1}^{\infty} \log \left(1 - \frac{\chi_i(p)}{p^s} \right) \\
 &= \sum_{p \text{ prime}} \frac{\chi_i(p)}{p^s} + \frac{1}{2} \left(\frac{\chi_i(p)}{p^s} \right)^2 + \frac{1}{3} \left(\frac{\chi_i(p)}{p^s} \right)^3 + \dots \\
 &= \sum_{p \text{ prime}} \frac{\chi_i(p)}{p^s} + R_i(s)
 \end{aligned}$$

where

$$R_i(s) = \sum_{p \text{ prime}} \frac{1}{2} \left(\frac{\chi_i(p)}{p^s} \right)^2 + \frac{1}{3} \left(\frac{\chi_i(p)}{p^s} \right)^3 + \frac{1}{4} \left(\frac{\chi_i(p)}{p^s} \right)^4 + \dots$$

See that

$$\begin{aligned}
 |R_i(s)| &\leq \sum_{p \text{ prime}} \frac{1}{2} \left(\frac{|\chi_i(p)|}{p^s} \right)^2 + \frac{1}{3} \left(\frac{|\chi_i(p)|}{p^s} \right)^3 + \frac{1}{4} \left(\frac{|\chi_i(p)|}{p^s} \right)^4 + \dots \\
 &\leq \sum_{p \text{ prime}} \frac{1}{2} \left(\frac{1}{p^s} \right)^2 + \frac{1}{2} \left(\frac{1}{p^s} \right)^3 + \frac{1}{2} \left(\frac{1}{p^s} \right)^4 + \dots \\
 &\text{by the geometric sum formula} \\
 &= \frac{1}{2} \sum_{p \text{ prime}} \frac{1}{p^s(p^s - 1)} \\
 &< \frac{1}{2} \sum_{p \text{ prime}} \frac{1}{p(p - 1)} \\
 &< \frac{1}{2} \sum_{n=1}^{\infty} \frac{1}{n(n - 1)} \\
 &= \frac{1}{2}
 \end{aligned}$$

and so $R_i(s)$ is bounded as $s \rightarrow 1^+$.

Now see that

$$L(s, \chi_1) = \sum_{b=1}^4 \sum_{n=0}^{\infty} \frac{1}{(5n+b)^s} \rightarrow \infty \text{ as } s \rightarrow 1^+.$$

For χ_2 see that

$$\begin{aligned}
 \Re(L(s, \chi_2)) &= \underbrace{\frac{1}{1^s} - \frac{1}{2^s} - \frac{1}{3^s} + \frac{1}{4^s}}_{> \frac{1}{6}} + \underbrace{\frac{1}{6^s} - \frac{1}{7^s} + \frac{1}{8^s} - \frac{1}{9^s}}_{> 0} + \underbrace{\dots}_{> 0} \\
 &> \frac{1}{6}
 \end{aligned}$$

and

$$\Re(L(s, \chi_2)) = \frac{1}{1^s} - \underbrace{\left(\frac{1}{2^s} + \frac{1}{3^s} - \frac{1}{4^s} - \frac{1}{6^s}\right)}_{>0} - \underbrace{\left(\frac{1}{7^s} + \frac{1}{8^s} - \frac{1}{9^s} - \frac{1}{11^s}\right)}_{>0} - \underbrace{\cdots}_{>0} < 1$$

so $\frac{1}{6} < \Re(L(s, \chi_2)) < 1$ for all $s > 1$.

For χ_3 see that

$$\Re(L(s, \chi_3)) = \underbrace{\frac{1}{1^s} - \frac{1}{4^s}}_{>\frac{3}{4}} + \underbrace{\frac{1}{6^s} + \frac{1}{9^s}}_{>0} + \underbrace{\cdots}_{>0} > \frac{3}{4}$$

and

$$\Re(L(s, \chi_3)) = \frac{1}{1^s} - \underbrace{\left(\frac{1}{4^s} - \frac{1}{6^s}\right)}_{>0} - \underbrace{\left(\frac{1}{9} - \frac{1}{11}\right)}_{>0} - \underbrace{\cdots}_{>0} < 1$$

so $\frac{3}{4} < \Re(L(s, \chi_3)) < 1$ for $s > 0$. We can also see that

$$\Re(L(s, \chi_4)) = \Re(L(s, \chi_3))$$

and so $\frac{3}{4} < \Re(L(s, \chi_4)) < 1$ for $s > 0$.

Now see that (by applying the result of part a)

$$\begin{aligned} \sum_{i=1}^4 \chi_i(b) \log(L(s, \chi_i)) &= \sum_{p \text{ prime}} \frac{\sum_{i=1}^4 \chi_i(b) \chi_i(p)}{p^s} + \underbrace{\sum_{i=1}^4 \chi_i(b) R_i(s)}_{\text{bounded as } s \rightarrow 1^+} \\ &= 4 \sum_{\substack{p \text{ prime} \\ p \equiv b \pmod{5}}} \frac{1}{p^s} + \underbrace{\sum_{i=1}^4 \chi_i(b) R_i(s)}_{\text{bounded as } s \rightarrow 1^+} . \end{aligned}$$

Now as only $\log(L(s, \chi_1))$ is unbounded as $s \rightarrow 1^+$ then

$$\sum_{\substack{p \text{ prime} \\ p \equiv b \pmod{5}}} \frac{1}{p^s} \rightarrow \infty \text{ as } s \rightarrow 1^+$$

which implies that there are infinitely many primes $p \equiv b \pmod{5}$ for $b \in \{1, 2, 3, 4\}$.