

Spatial Data and Analysis in R

A PRISM Workshop

Adam Lauretig

The Ohio State University



Outline

Introduction

Spatial Data Prep

Spatial Autocorrelation

Regression

Spatial Regression

Discussion

Why Are We Here?

- ▶ Tobler's first law of geography:

Why Are We Here?

- ▶ Tobler's first law of geography:
- ▶ “Everything is related to everything else, but near things are more related than distant things”

Why Are We Here?

- ▶ Tobler's first law of geography:
- ▶ “Everything is related to everything else, but near things are more related than distant things”
- ▶ We want to quantify how the spatial relationship between our observations affect our inferences

A Caveat

- ▶ There are entire disciplines which study these issues (one of them is downstairs)

A Caveat

- ▶ There are entire disciplines which study these issues (one of them is downstairs)
- ▶ I will be introducing *spatial statistics* with a touch of *GIS*

A Caveat

- ▶ There are entire disciplines which study these issues (one of them is downstairs)
- ▶ I will be introducing *spatial statistics* with a touch of *GIS*
- ▶ I will not be discussing *GIS* in depth, nor will I discuss remote sensing *at all*

What are Spatial Data?

- ▶ Information (attributes) associated with a location

What are Spatial Data?


- ▶ Information (attributes) associated with a location
- ▶ Many kinds of spatial data: Points, Lines, Polygons, Raster data

What are Spatial Data?

- ▶ Information (attributes) associated with a location
- ▶ Many kinds of spatial data: Points, Lines, Polygons, Raster data
- ▶ Today, we are working with polygon data


Prepping our data

- Spatial data come in *shapefiles* which are really mini-databases




Prepping our data

- ▶ Spatial data come in *shapefiles* which are really mini-databases
- ▶ ORDBMS - Linking spatial and attribute data



Prepping our data


- ▶ Spatial data come in *shapefiles* which are really mini-databases
- ▶ ORDBMS - Linking spatial and attribute data
- ▶ Six parts, all combine to create a map to represent data




Loading Our data



Percent Black



AIDs Rate per 1000 people



Measuring Spatial Autocorrelation

- ▶ What is spatial autocorrelation?

Measuring Spatial Autocorrelation

- ▶ What is spatial autocorrelation?
- ▶ Observations with more similar values tend to occur more closely together

Measuring Spatial Autocorrelation

- ▶ What is spatial autocorrelation?
- ▶ Observations with more similar values tend to occur more closely together
- ▶ Most common test: Moran's I

Formula for Moran's I

$$I = \frac{n}{\sum_{i=1}^n \sum_{j=1}^n w_{ij}} \frac{\sum_{i=1}^n \sum_{j=1}^n w_{ij} (x_i - \bar{x})(x_j - \bar{x})}{\sum_{i=1}^n (x_i - \bar{x})^2},$$

where w_{ij} is the weight between observation i and j

The Weights Matrix W

- ▶ In Moran's I , there was this thing w_{ij}

The Weights Matrix W

- ▶ In Moran's I , there was this thing w_{ij}
- ▶ This is the *weights matrix*

The Weights Matrix W

- ▶ In Moran's I , there was this thing w_{ij}
- ▶ This is the *weights matrix*
- ▶ It allows us to measure the effect neighboring observations j have on our observation of interest i

The Weights Matrix W

- ▶ In Moran's I , there was this thing w_{ij}
- ▶ This is the *weights matrix*
- ▶ It allows us to measure the effect neighboring observations j have on our observation of interest i
- ▶ Can be specified in a variety of ways, the simplest of which is binary ("contiguity"): 1 if observations share a boundary, 0 if they do not

The Weights Matrix W

- ▶ In Moran's I , there was this thing w_{ij}
- ▶ This is the *weights matrix*
- ▶ It allows us to measure the effect neighboring observations j have on our observation of interest i
- ▶ Can be specified in a variety of ways, the simplest of which is binary ("contiguity"): 1 if observations share a boundary, 0 if they do not
- ▶ The default in R is "row standardized," where $w_{ij} = \frac{1}{\sum_j}$

Creating a weights matrix in *R*

```
library(rgdal)
library(spdep)
library(sp)
library(spatstat)
file_path <- "/Users/adamlauretig/data/Prism_presentation/NYAIDS_data"
ny <- readOGR(dsn = file_path,
             layer = "NYAIDS", verbose=FALSE)
nygal <- poly2nb(ny) #Create the neighborhood object
nyQ1.gal <- nb2listw(nygal, zero.policy=T) #Create the weights object
```

Running the Moran's I

Moran I statistic	Expectation	Variance	p-value
0.68	-0.01	0.00	0.00

Where are these clusters?

- ▶ We can calculate this using a *Local Indicator of Spatial Autocorrelation* (LISA)


Where are these clusters?

- ▶ We can calculate this using a *Local Indicator of Spatial Autocorrelation* (LISA)
- ▶ Measure how similar a value is compared to neighboring values

Where are these clusters?


- ▶ We can calculate this using a *Local Indicator of Spatial Autocorrelation* (LISA)
- ▶ Measure how similar a value is compared to neighboring values
- ▶ While the Moran's I detects clustering, the LISA detects *clusters*

Where are these clusters?



Where are these clusters?

LISA Map AIDS Rate; weights: Q1



What About Regression with Spatial Data?

- ▶ Remember, a key regression assumption is that our observations are IID

What About Regression with Spatial Data?

- ▶ Remember, a key regression assumption is that our observations are IID
- ▶ If there is spatial autocorrelation in our data, we have fewer independent data points than we initially supposed

What About Regression with Spatial Data?

- ▶ Remember, a key regression assumption is that our observations are IID
- ▶ If there is spatial autocorrelation in our data, we have fewer independent data points than we initially supposed
- ▶ This shrinks our standard errors, making variables appear significant

What About Regression with Spatial Data?

- ▶ Remember, a key regression assumption is that our observations are IID
- ▶ If there is spatial autocorrelation in our data, we have fewer independent data points than we initially supposed
- ▶ This shrinks our standard errors, making variables appear significant
- ▶ Once we adjust our variance-covariance matrix, previously significant covariates might lose their significance

What About Regression with Spatial Data?

- ▶ Remember, a key regression assumption is that our observations are IID
- ▶ If there is spatial autocorrelation in our data, we have fewer independent data points than we initially supposed
- ▶ This shrinks our standard errors, making variables appear significant
- ▶ Once we adjust our variance-covariance matrix, previously significant covariates might lose their significance
- ▶ Additionally, with spatial autocorrelation, our coefficients may be biased

What About Regression with Spatial Data?

- ▶ Remember, a key regression assumption is that our observations are IID
- ▶ If there is spatial autocorrelation in our data, we have fewer independent data points than we initially supposed
- ▶ This shrinks our standard errors, making variables appear significant
- ▶ Once we adjust our variance-covariance matrix, previously significant covariates might lose their significance
- ▶ Additionally, with spatial autocorrelation, our coefficients may be biased
- ▶ Two ways of handling this: Spatial Error Models, and Spatial Autoregressive models

The Math: Spatial Error Model

- ▶ Normally: $y_i = x_i\beta + e_i$, where $e = I(Y - X\beta)$

The Math: Spatial Error Model

- ▶ Normally: $y_i = x_i\beta + e_i$, where $e = I(Y - X\beta)$
- ▶ But, we want to model spatial dependence in the residuals

The Math: Spatial Error Model

- ▶ Normally: $y_i = x_i\beta + e_i$, where $e = I(Y - X\beta)$
- ▶ But, we want to model spatial dependence in the residuals
- ▶ $e_i = \sum_{j=1}^n w_{ij} + e_j + \varepsilon_i$, where $w_{ij} = 0$

The Math: Spatial Error Model

- ▶ Normally: $y_i = x_i\beta + e_i$, where $e = I(Y - X\beta)$
- ▶ But, we want to model spatial dependence in the residuals
- ▶ $e_i = \sum_{j=1}^n w_{ij} + e_j + \varepsilon_i$, where $w_{ii} = 0$
- ▶ Basically, we regress the error e_i on the surrounding errors

The Math: Spatial Error Model

- ▶ Normally: $y_i = x_i\beta + e_i$, where $e = I(Y - X\beta)$
- ▶ But, we want to model spatial dependence in the residuals
- ▶ $e_i = \sum_{j=1}^n w_{ij} + e_j + \varepsilon_i$, where $w_{ii} = 0$
- ▶ Basically, we regress the error e_i on the surrounding errors
- ▶ We wind up with $e = (I - W)(Y - X\beta)$

The Math: Spatial Error Model

- ▶ Normally: $y_i = x_i\beta + e_i$, where $e = I(Y - X\beta)$
- ▶ But, we want to model spatial dependence in the residuals
- ▶ $e_i = \sum_{j=1}^n w_{ij} + e_j + \varepsilon_i$, where $w_{ii} = 0$
- ▶ Basically, we regress the error e_i on the surrounding errors
- ▶ We wind up with $e = (I - W)(Y - X\beta)$
- ▶ ε is the residual of residuals, with $\sum_{\varepsilon} = \sigma^2 I$

The Math: Spatial Error Model

- ▶ Normally: $y_i = x_i\beta + e_i$, where $e = I(Y - X\beta)$
- ▶ But, we want to model spatial dependence in the residuals
- ▶ $e_i = \sum_{j=1}^n w_{ij} + e_j + \varepsilon_i$, where $w_{ii} = 0$
- ▶ Basically, we regress the error e_i on the surrounding errors
- ▶ We wind up with $e = (I - W)(Y - X\beta)$
- ▶ ε is the residual of residuals, with $\sum \varepsilon = \sigma^2 I$
- ▶ The full model: $y_i = x_i\beta + \sum_{j=1}^n w_{ij}e_j + \varepsilon_i$

The Math: Spatial Autoregressive Model

- ▶ Option 2: the Spatial autoregressive model

The Math: Spatial Autoregressive Model

- ▶ Option 2: the Spatial autoregressive model
- ▶ Instead of lagging the error term, lag y , the DV

The Math: Spatial Autoregressive Model

- ▶ Option 2: the Spatial autoregressive model
- ▶ Instead of lagging the error term, lag y , the DV
- ▶ $y_i = x_i\beta + \sum_{j=1}^n w_{ij}y_j + \varepsilon_i$

The Math: Spatial Autoregressive Model

- ▶ Option 2: the Spatial autoregressive model
- ▶ Instead of lagging the error term, lag y , the DV
- ▶ $y_i = x_i\beta + \sum_{j=1}^n w_{ij}y_j + \varepsilon_i$
- ▶ SAR vs. SEM


Plain OLS

	Model 1
(Intercept)	-3.05*** (0.80)
PctWht	-0.01*** (0.00)
PctHisp	0.02*** (0.00)
Gini	7.97*** (0.71)
PctHSEd	0.02** (0.01)
PctFemHH	0.01 (0.01)
R ²	0.57
Adj. R ²	0.55
Num. obs.	174
RMSE	0.54

*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

Table: Statistical models

Did we model out our autocorrelation?




Did we model out our spatial autocorrelation?

Moran I statistic	Expectation	Variance	p-value
0.63	-0.01	0.00	0.00

No! This statistic didn't change much from before we ran our regression: .68 vs .63

Where is the residual autocorrelation?

LISA Map AIDS Rate; weights: Q1



Can we Model this Spatial Autocorrelation?

- ▶ One option is to use fixed effects

Can we Model this Spatial Autocorrelation?

- ▶ One option is to use fixed effects
- ▶ But this removes something interesting – the spatial relationship

Can we Model this Spatial Autocorrelation?

- ▶ One option is to use fixed effects
- ▶ But this removes something interesting – the spatial relationship
- ▶ Another option is to explicitly model the spatial relationship

Spatial Autoregressive Model

	Model 1
(Intercept)	-1.52* (0.63)
PctWht	-0.01*** (0.00)
PctHisp	0.01** (0.00)
Gini	4.48*** (0.64)
PctHSEd	0.01 (0.01)
PctFemHH	0.00 (0.01)
ρ	0.54*** (0.06)
Num. obs.	174
Parameters	8
Log Likelihood	-101.93
AIC (Linear model)	285.72
AIC (Spatial model)	219.85
LR test: statistic	67.86
LR test: p-value	0.00

*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

Table: SAR Model output

Spatial Error Model


	Model 1
(Intercept)	-0.28 (0.88)
PctWht	-0.01** (0.00)
PctHisp	0.01** (0.00)
Gini	4.25*** (0.90)
PctHSEd	0.01 (0.01)
PctFemHH	0.01 (0.01)
λ	0.66*** (0.06)
Num. obs.	174
Parameters	8
Log Likelihood	-107.85
AIC (Linear model)	285.72
AIC (Spatial model)	231.70
LR test: statistic	56.01
LR test: p-value	0.00

*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$

Table: SEM model output

Comparing Findings - OLS

Comparing Regression Results



Did These Resolve our Spatial Autocorrelation?

Moran I statistic	Expectation	Variance	p-value
-0.10	-0.01	0.00	0.08

YES!

Conditional Autoregressive Model (CAR)

- ▶ Model the spatial relationship according to a Markov Random Field (MRF)

Conditional Autoregressive Model (CAR)

- ▶ Model the spatial relationship according to a Markov Random Field (MRF)
- ▶ Each observation depends only on values at neighboring locations, not global values

Conditional Autoregressive Model (CAR)

- ▶ Model the spatial relationship according to a Markov Random Field (MRF)
- ▶ Each observation depends only on values at neighboring locations, not global values
- ▶ Written out:

Conditional Autoregressive Model (CAR)

- ▶ Model the spatial relationship according to a Markov Random Field (MRF)
- ▶ Each observation depends only on values at neighboring locations, not global values
- ▶ Written out:
- ▶ c_{ij} is the weight matrix

Conditional Autoregressive Model (CAR)

- ▶ Model the spatial relationship according to a Markov Random Field (MRF)
- ▶ Each observation depends only on values at neighboring locations, not global values
- ▶ Written out:
- ▶ c_{ij} is the weight matrix
- ▶

$$E(y_i|y_{-i}) = x_i\beta + \sum_{j=1}^n c_{ij}[y_j - x_j\beta]$$

Conditional Autoregressive Model (CAR)

- ▶ Model the spatial relationship according to a Markov Random Field (MRF)
- ▶ Each observation depends only on values at neighboring locations, not global values
- ▶ Written out:
- ▶ c_{ij} is the weight matrix

$$E(y_i|y_{-i}) = x_i\beta + \sum_{j=1}^n c_{ij}[y_j - x_j\beta]$$

$$\text{var}(y_i|y_{-i}) = \sigma_i^2$$

Conditional Autoregressive Model (CAR) (cont'd)

- ▶ This defines a joint multivariate normal distribution with variance

Conditional Autoregressive Model (CAR) (cont'd)

- ▶ This defines a joint multivariate normal distribution with variance



$$\Sigma_Y = (I - C)^{-1} \sum_C$$

Conditional Autoregressive Model (CAR) (cont'd)

- ▶ This defines a joint multivariate normal distribution with variance

- ▶
$$\Sigma_Y = (I - C)^{-1} \sum_C$$

- ▶
$$\Sigma_{CAR} = \sigma^2 (I - C)^{-1} V_C$$

Conditional Autoregressive Model (CAR) (cont'd)

- ▶ This defines a joint multivariate normal distribution with variance

- ▶
$$\Sigma_Y = (I - C)^{-1} \sum_C$$

- ▶
$$\Sigma_{CAR} = \sigma^2 (I - C)^{-1} V_C$$

- ▶
$$= \sigma^2 V_{CAR}$$

SAR vs. CAR

► SAR:

$$y_i \sim N(0, (I - W)^{-1} D^{\sim} (I - W')^{-1})$$

$$\Sigma_{SAR} = \sigma^2 (I - W)^{-1} V_{\epsilon} (I - W')^{-1}$$

SAR vs. CAR

► SAR:

$$y_i \sim N(0, (I - W)^{-1} D (I - W')^{-1})$$

$$\Sigma_{SAR} = \sigma^2 (I - W)^{-1} V_\epsilon (I - W')^{-1}$$

► CAR:

$$y_i \sim N(0, (I - C)^{-1} D)$$

$$\Sigma_{CAR} = \sigma^2 (I - C)^{-1} V_C$$

SAR vs. CAR

- ▶ CAR only requires the variance to estimate, SAR requires the full variance-covariance matrix

SAR vs. CAR

- ▶ CAR only requires the variance to estimate, SAR requires the full variance-covariance matrix
- ▶ CAR is computationally less intensive

SAR vs. CAR

- ▶ CAR only requires the variance to estimate, SAR requires the full variance-covariance matrix
- ▶ CAR is computationally less intensive
- ▶ CAR: local/first order dependency

SAR vs. CAR

- ▶ CAR only requires the variance to estimate, SAR requires the full variance-covariance matrix
- ▶ CAR is computationally less intensive
- ▶ CAR: local/first order dependency
- ▶ CAR: fit with MCMC – can fit GLMs

SAR vs. CAR

- ▶ CAR only requires the variance to estimate, SAR requires the full variance-covariance matrix
- ▶ CAR is computationally less intensive
- ▶ CAR: local/first order dependency
- ▶ CAR: fit with MCMC – can fit GLMs
- ▶ SAR: global/higher order dependency

SAR vs. CAR

- ▶ CAR only requires the variance to estimate, SAR requires the full variance-covariance matrix
- ▶ CAR is computationally less intensive
- ▶ CAR: local/first order dependency
- ▶ CAR: fit with MCMC – can fit GLMs
- ▶ SAR: global/higher order dependency
- ▶ SAR: fit with MLE

Fitting a CAR (1)

	Median	2.5%	97.5%	n.sample	% accept	n.effective	Geweke.diag
(Intercept)	-0.60	-2.35	1.08	18000.00	100.00	2417.70	0.50
PctWht	-0.01	-0.02	-0.01	18000.00	100.00	2973.70	1.40
PctHispanic	0.01	0.01	0.02	18000.00	100.00	2592.00	-0.80
Gini	3.98	2.10	5.97	18000.00	100.00	1628.90	-0.30
PctHSEd	0.02	0.00	0.03	18000.00	100.00	3252.90	-0.90
PctFemHH	0.01	-0.01	0.03	18000.00	100.00	2006.10	0.20
nu2	0.01	0.00	0.06	18000.00	100.00	1599.50	0.70
tau2	0.50	0.34	0.65	18000.00	100.00	3907.90	-0.80
rho	0.81	0.61	0.94	18000.00	44.10	5196.60	0.90

Table: uninformative priors

Fitting a CAR (2)

	Median	2.5%	97.5%	n.sample	% accept	n.effective	Geweke.diag
(Intercept)	-0.81	-2.48	0.86	18000.00	100.00	11033.20	1.40
PctWht	-0.01	-0.02	-0.01	18000.00	100.00	13600.70	0.20
PctHispanic	0.01	0.01	0.02	18000.00	100.00	13780.40	0.30
Gini	4.03	2.12	5.93	18000.00	100.00	7508.30	-2.00
PctHSEd	0.02	0.00	0.03	18000.00	100.00	15248.50	-0.70
PctFemHH	0.01	-0.01	0.03	18000.00	100.00	10575.60	1.30
nu2	0.08	0.06	0.13	18000.00	100.00	10732.90	-1.30
tau2	0.25	0.16	0.38	18000.00	100.00	10628.20	2.70
rho	0.91	0.75	0.98	18000.00	43.80	10169.70	0.50

Table: weakly informative priors

Geographically Weighted Regression

- ▶ A non-parametric method for inference with non-stationary spatial data

Geographically Weighted Regression

- ▶ A non-parametric method for inference with non-stationary spatial data
- ▶ Non-stationarity: spatial variation is not constant across the data

Geographically Weighted Regression

- ▶ A non-parametric method for inference with non-stationary spatial data
- ▶ Non-stationarity: spatial variation is not constant across the data
- ▶ Use kernel regression to model local spatial variation

Geographically Weighted Regression


- ▶ A non-parametric method for inference with non-stationary spatial data
- ▶ Non-stationarity: spatial variation is not constant across the data
- ▶ Use kernel regression to model local spatial variation
- ▶ A weighted moving window regression:

$$Y(x) = \alpha(u_i, v_i) + \sum_k \beta_k(u_i, v_i) x_k + e_i$$

Geographically Weighted Regression (Cont'd)

- ▶ A Kernel as a way of weighting observations which are closer together

The concept of a kernel




- × regression point
 - data point
- w_{ij} is the weight of data point j at regression point i
 d_{ij} is the distance between regression point i and data point j

Geographically Weighted Regression (Cont'd)

- ▶ A Kernel as a way of weighting observations which are closer together
- ▶ Usually, fit adaptively – different “sizes” depending on the density of the data


The concept of a kernel



- x regression point w_{ij} is the weight of data point j at regression point i
 • data point d_{ij} is the distance between regression point i and data point j

Fitting a Geographically Weighted Regression

- Usually, we represent the results from a GWR visually



Geographically Weighted Regression (Cont'd)

- ▶ GWR usually more of an exploratory tool than a hypothesis-testing tool

Geographically Weighted Regression (Cont'd)

- ▶ GWR usually more of an exploratory tool than a hypothesis-testing tool
- ▶ Danger: overfitting the data

Geographically Weighted Regression (Cont'd)

- ▶ GWR usually more of an exploratory tool than a hypothesis-testing tool
- ▶ Danger: overfitting the data
- ▶ Extensions: GWR + Ridge regression, GWR GLMs, GWPCA

Workflow for Spatial Regression

- ▶ In addition to normal EDA, do some ESDA (Exploratory Spatial Data Analysis), mapping out variables

Workflow for Spatial Regression

- ▶ In addition to normal EDA, do some ESDA (Exploratory Spatial Data Analysis), mapping out variables
- ▶ Check for spatial autocorrelation

Workflow for Spatial Regression

- ▶ In addition to normal EDA, do some ESDA (Exploratory Spatial Data Analysis), mapping out variables
- ▶ Check for spatial autocorrelation
- ▶ Run your normal regression, with the variables you think are necessary

Workflow for Spatial Regression

- ▶ In addition to normal EDA, do some ESDA (Exploratory Spatial Data Analysis), mapping out variables
- ▶ Check for spatial autocorrelation
- ▶ Run your normal regression, with the variables you think are necessary
- ▶ Check once more for spatial autocorrelation, in your residuals

Workflow for Spatial Regression

- ▶ In addition to normal EDA, do some ESDA (Exploratory Spatial Data Analysis), mapping out variables
- ▶ Check for spatial autocorrelation
- ▶ Run your normal regression, with the variables you think are necessary
- ▶ Check once more for spatial autocorrelation, in your residuals
- ▶ If there's still autocorrelation, run a spatial model

Workflow for Spatial Regression

- ▶ In addition to normal EDA, do some ESDA (Exploratory Spatial Data Analysis), mapping out variables
- ▶ Check for spatial autocorrelation
- ▶ Run your normal regression, with the variables you think are necessary
- ▶ Check once more for spatial autocorrelation, in your residuals
- ▶ If there's still autocorrelation, run a spatial model
- ▶ One final check for autocorrelation in your residuals

Where Can we Apply These Methods in Political Science

- ▶ Voting and Political Behavior patterns (Data available at the Census Tract level (or less))

Where Can we Apply These Methods in Political Science

- ▶ Voting and Political Behavior patterns (Data available at the Census Tract level (or less))
- ▶ Agricultural/industrial data (economic output)

Where Can we Apply These Methods in Political Science

- ▶ Voting and Political Behavior patterns (Data available at the Census Tract level (or less))
- ▶ Agricultural/industrial data (economic output)
- ▶ Conflict/Political Violence data (ex: ACLED)

Where Can we Apply These Methods in Political Science

- ▶ Voting and Political Behavior patterns (Data available at the Census Tract level (or less))
- ▶ Agricultural/industrial data (economic output)
- ▶ Conflict/Political Violence data (ex: ACLED)
- ▶ Anything with an address or lon/lat coordinates can be georeferenced

Other tools

- ▶ Spatial/Spatio-temporal scan statistics

Other tools

- ▶ Spatial/Spatio-temporal scan statistics
- ▶ Kernel Density Estimation

Other tools

- ▶ Spatial/Spatio-temporal scan statistics
- ▶ Kernel Density Estimation
- ▶ Kriging/Geostatistics

Other tools

- ▶ Spatial/Spatio-temporal scan statistics
- ▶ Kernel Density Estimation
- ▶ Kriging/Geostatistics
- ▶ Spatio-temporal approaches

Resources

- ▶ Yuri Zhukov's Spatial Workshop:
<http://www.people.fas.harvard.edu/~zhukov/spatial.html>

Resources

- ▶ Yuri Zhukov's Spatial Workshop:
<http://www.people.fas.harvard.edu/~zhukov/spatial.html>
- ▶ Brunsdon, Chris, and Lex Comber. *An introduction to R for spatial analysis & mapping*. Sage, 2015.

Resources

- ▶ Yuri Zhukov's Spatial Workshop:
<http://www.people.fas.harvard.edu/~zhukov/spatial.html>
- ▶ Brunsdon, Chris, and Lex Comber. *An introduction to R for spatial analysis & mapping*. Sage, 2015.
- ▶ Bivand, Roger S., and Edzer J. Pebesma. *Applied Spatial Data Analysis with R*. Springer, 2013.

Resources

- ▶ Yuri Zhukov's Spatial Workshop:
<http://www.people.fas.harvard.edu/~zhukov/spatial.html>
- ▶ Brunsdon, Chris, and Lex Comber. *An introduction to R for spatial analysis & mapping*. Sage, 2015.
- ▶ Bivand, Roger S., and Edzer J. Pebesma. *Applied Spatial Data Analysis with R*. Springer, 2013.
- ▶ Waller, Lance A., and Carol A. Gotway. *Applied spatial statistics for public health data*. John Wiley & Sons, 2004.

Resources

- ▶ Yuri Zhukov's Spatial Workshop:
<http://www.people.fas.harvard.edu/~zhukov/spatial.html>
- ▶ Brunsdon, Chris, and Lex Comber. *An introduction to R for spatial analysis & mapping*. Sage, 2015.
- ▶ Bivand, Roger S., and Edzer J. Pebesma. *Applied Spatial Data Analysis with R*. Springer, 2013.
- ▶ Waller, Lance A., and Carol A. Gotway. *Applied spatial statistics for public health data*. John Wiley & Sons, 2004.
- ▶ Cressie, Noel. *Statistics for spatial data*. John Wiley & Sons, 1993.

Acknowledgements

Thank you to Jan Box-Steffensmeier for this opportunity, and to Elisabeth Root, whose code I drew heavily from.

References

