

This Notebook was developed in February 2017 on the basis of a Notebook written in 2011

Last updating March 8th 2017

Construction of the G2 Lie algebra

Pietro G. Frè

University of Torino, Physics Department, Via. P. Giuria 1, 10125, Torino Italy

Embassy of Italy in the Russian Federation, Denezhny Pereulok, 5, Moscow, Russia

Instructions

Group Theory of $G_{2,2}$

In an orthonormal basis the $g_{2(2)}$ positive roots α (rutte) read :

$$\alpha_1 = (1, 0) ;$$

$$\alpha_2 = \left(-\frac{3}{2}, \frac{\sqrt{3}}{2} \right) ;$$

$$\alpha_3 = \alpha_1 + \alpha_2 = \left(-\frac{1}{2}, \frac{\sqrt{3}}{2} \right) ;$$

$$\alpha_4 = 2\alpha_1 + \alpha_2 = \left(\frac{1}{2}, \frac{\sqrt{3}}{2} \right) ;$$

$$\alpha_5 = 3\alpha_1 + \alpha_2 = \left(\frac{3}{2}, \frac{\sqrt{3}}{2} \right) ;$$

$$\alpha_6 = 3\alpha_1 + 2\alpha_2 = (0, \sqrt{3}) ;$$

Instructions for the user

After running this Notebook you have at your disposal the explicit form of the g_2 Lie algebra in the fundamental 7-dimensional representation and in the maximally split real section where all the generators are completely real.

The available objects are the following ones

1. The array **rutte** contains the six positive roots enumerated as above
2. The array **Steppig2** contains the step operators associated with the positive roots and enumerated in the same order as in **rutte**.
3. The array **MSteppig2** contains the step operators associated with the negative roots enumerated in the same order as their opposites in **rutte**.
4. The array **Cartanig2** contains the Cartan generators in an orthonormal basis.
5. The array **TTsolv** contains the 8 generators of the Borel solvable Lie algebra enumerated as follows $= \{\mathcal{H}_1, \mathcal{H}_2, \mathcal{E}^{\alpha_1}, \dots, \mathcal{E}^{\alpha_6}\}$

Construction of the fundamental

representation in the standard Weyl basis