

Exploring a lower resolution physics grid in CAM-SE-CSLAM

A.R. Herrington^{1*}, P.H. Lauritzen², S. Goldhaber², Mark A. Taylor³, K.A. Reed¹

¹School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, New York

²National Center for Atmospheric Research, Boulder, Colorado, USA

³Sandia National Laboratories, Albuquerque, New Mexico, USA

Key Points:

-
-
-

*Stony Brook, New York

Corresponding author: Adam R. Herrington, adam.herrington@stonybrook.edu

Abstract

1 Introduction

Global atmospheric models fundamentally consist of two components. The dynamical core (*dynamics*), which numerically integrate the adiabatic equations of motion and tracer advection, and the physical parameterizations (*physics*), which compute the effects of diabatic and subgrid-scale processes (e.g., radiative transfer and moist convection) on the grid-scale. More out of convenience than anything else, the physics are evaluated on the dynamics grid, i.e., the physics grid and dynamics grid coincide. From linear stability and accuracy analysis of numerical methods, it is a common result that the shortest simulated wavelengths are not accurately represented by the dynamical core. Additionally, simulated downscale cascades result in an unrealistic collection of energy and/or enstrophy near the truncation scale, which may be observed from kinetic energy spectra in model simulations [Skamarock, 2011]. Some form of dissipation must be incorporated into models to mitigate these numerical artifacts near the grid scale [Jablonowski and Williamson, 2011]. This numerical dissipation has no physical analogy [although see Grinstein *et al.*, 2007], and the grid-scale is therefore contaminated by numerous un-physical processes. The under-resolved nature of the grid-scale led Lander and Hoskins [1997] to speculate whether the physics should be evaluated on a grid that is more reflective of the scales actually resolved by the dynamical core.

Experimentation with different physics grid resolutions have so far been limited to models employing the spectral transform method [Lander and Hoskins, 1997; Williamson, 1999; Wedi, 2014]. Lander and Hoskins [1997] argue that passing under-resolved states to the physics may be especially problematic in spectral transform models, since the physics are evaluated on a latitude-longitude transform grid, and contains more degrees of freedom than the spectral representation to prevent aliasing of quadratic quantities. However, Lander and Hoskins [1997] indicate that the spectral truncation of the physics tendencies damps errors that may result from passing an under-resolved state to the physics, although the extent to which these errors may still be present in the model was not addressed.

Another class of spectral-transform models evaluate the quadratic terms using semi-Lagrangian methods, which are implicitly diffusive, relaxing constraints on the resolution of the transform grid. Wedi [2014] experimented with different transform grid resolutions and concluded that the standard high resolution quadratic grid actually improves forecast skill over the use of a lower-resolution transform grid. They suggest that increasing the resolution of the transform grid simulates a kind of sub-grid variability on the spectral state, which is thought to be under-represented in global atmospheric models [Shutts, 2005]. This is in principle the purpose of “super-parameterization,” in which a cloud resolving model is embedded in each grid cell to simulate the requisite subgrid variability, and improves both diurnal and sub-seasonal variability in the model [Randall *et al.*, 2003].

After the physics tendencies are transformed into spectral space, the tendencies may be truncated at any particular wave number. Williamson [1999] conducted a pair of convergence tests using a global spectral transform model; a conventional convergence test and one in which the spectral truncation of the physics tendencies is held fixed and the resolution of the dynamical core increased. In contrast to the realistic weather forecasts of Wedi [2014], Williamson [1999] run their model to equilibrium in an idealized climate configuration. When the physics and dynamics resolutions increase together, as in more typical convergence studies, the strength of the Hadley Cell increases monotonically with resolution. This sensitivity of Hadley Cell strength to horizontal resolution is a common result of global models at hydrostatic resolutions [see Herrington and Reed, 2017, and references therein]. But with the truncation wave number of physics tendencies held fixed, the Hadley Cell showed very little sensitivity to dynamical core resolution, resembling the

solution for which the dynamics truncation wave number is equal to that of the lower resolution physics.

Herrington and Reed [2017] speculate that the results of *Williamson* [1999] indicate that the scales of motion resolved by the dynamical core are aliased to the lower resolution physics. It may be worth considering that if the resolution of the dynamics is reduced in response to a coarser physics grid, then the dynamics may be no better resolved on the coarser physics grid, compared with the conventional method of evaluating the physics and dynamics at the same resolution. The results of *Williamson* [1999]; *Wedi* [2014] do not provide evidence that a lower resolution physics grid reduces computational errors in spectral transform models, but this was seldom discussed in either study.

Global spectral transform models, while remarkably efficient at small processor counts, do not scale well on massively parallel systems. High-order Galerkin methods are becoming increasingly popular in climate and weather applications due to their high-parallel efficiency, high-processor efficiency, high-order accuracy (for smooth problems), and geometric flexibility facilitating mesh-refinement applications. High resolution climate simulations with NCAR’s Community Atmosphere Model [CAM; *Neale et al.*, 2012] are typically performed using a continuous Galerkin dynamical core referred to as CAM-SE [CAM Spectral Elements; *Taylor et al.*, 2008; *Dennis et al.*, 2012; *Lauritzen et al.*, 2018]. CAM-SE may be optionally coupled to a conservative, semi-Lagrangian tracer advection scheme for accelerated multi-tracer transport [CAM-SE-CSLAM; *Lauritzen et al.*, 2017]. Tracer advection then evolves on an entirely separate, finite-volume grid which contains the same degrees of freedom as CAM-SE’s quadrature node grid.

Element-based Galerkin methods are susceptible to grid-imprinting, and may need be considered when contemplating a particular physics grid [*Herrington et al.*, 2018, hereafter referred to as HL18]. Grid imprinting manifests at the element boundaries, since the global basis is least smooth (C^0 ; all derivatives are discontinuous) for quadrature nodes lying on the element boundaries, and the gradients (e.g., the pressure gradient) are systematically tighter producing local extremes. Through computing the physics tendencies at the nodal points, element boundary extrema is also observed in the physics tendencies.

HL18 has shown that through evaluating the physics on the finite-volume tracer advection grid in CAM-SE-CSLAM, element boundary noise is substantially reduced, although still problematic in regions of steep terrain, at low latitudes. Through integrating CAM-SE’s basis functions over the control volumes of the finite-volume grid, element boundary extrema is additionally weighted by the C^∞ solutions (i.e., the basis representation is infinitely smooth and all derivatives are continuous) of the element interior, and the state is smoother. Additionally, in defining an area averaged state, the finite-volume physics grid is made consistent with assumptions inherent to the physics, and is more appropriate for coupling to other model components (e.g., the land model), which is typically performed using finite-volume based mapping algorithms.

The finite-volume grid of HL18 is found through dividing the elements of CAM-SE’s gnomonic cubed-sphere grid with equally spaced, equi-angular coordinate lines parallel to the element boundaries, such that there are 3×3 control volumes per element (hereafter referred to as *pg3*). While a 3×3 physics grid was chosen in order to have the same degrees of freedom as the dynamical core, the control volumes encompass a region of the element in which their proximity to the element boundaries are not equal. Therefore, not every control volume in an element has the same smoothness properties. This may be avoided through defining a physics grid in which the elements are instead divided into 2×2 control volumes (hereafter referred to as *pg2*). The control volumes of the *pg2* grid all have the same proximity to the element boundaries, and should mitigate the element boundary noise that remains in the *pg3* grid, and shown in HL18.

In this study, we test the hypothesis that the coarser, *pg2* physics grid is effective at reducing spurious noise at element boundaries, particularly over regions of rough topography. In addition, the recent trend towards running models at ever higher resolutions is an almost prohibitive computational burden. As the physics makes up over half of the computational cost in CAM-SE [Lauritzen *et al.*, 2018], the improvement in computational performance using a coarser resolution physics grid is potentially significant. However, any advantages of using a coarser physics grid need be weighed against any potential reduction in simulation quality, e.g., possible aliasing of the resolved scales of motion by the coarser grid, as suggested by the results of Williamson [1999]. Section 2 described the implementation of the *pg2* grid into CAM-SE-CSLAM. Section 3 provides the results of a hierarchy of model configurations to identify any changes in grid imprinting, or in the overall solution, compared with the *pg3* configuration. Section 4 provides a discussion of the results and conclusions.

2 Methods

Separating dynamics, tracer and physics grids introduces the added complexity of having to map the state from dynamics and tracer grids to the physics grid; and mapping physics tracer tendencies back to the tracer grid and physics tendencies needed by the dynamical core to the dynamics grid. The dynamics grid refers to the Gauss-Lobatto-Legendre (GLL) quadrature nodes by the spectral-element method to solve the momentum equations for the momentum vector (u, v) , thermodynamics equation for temperature (T) , continuity equation for dry air (M) , and continuity equations for water vapor and condensates thermodynamically active [see, e.g., Lauritzen *et al.*, 2018, for details]. By tracer grid we refer to the *pg3* grid on which CSLAM performs tracer transport of water vapor, condensates and other tracers. The GLL value for water vapor and condensates is overwritten by the CSLAM values every physics time-step so that the spectral-element advection of water species does not become decoupled from the the CSLAM advection of the same water species. Mapping velocity components, dry air mass and temperature from the GLL grid to the *pg2* grid is done by using the internal degree 3 Lagrange basis functions in CAM-SE [as described in Herrington *et al.*, 2018, for *pg3*; exactly the same methods can be used for *pg2*].

As compared to the *pg3* configuration, the extra complication of the *pg2* setup is that tracer state needs to be mapped from the tracer grid to the physics grid and tracer tendencies need to be mapped from the physics grid to CSLAM grid. In order to describe the algorithm some notation needs to be introduced.

The mapping algorithm is applied to each element Ω (with spherical area $\Delta\Omega$) so without loss of generality consider one element. Let $\Delta A_k^{(pg)}$ and $\Delta A_\ell^{(nc)}$ be the spherical area of the physics grid grid cell $A_k^{(pg)}$ and CSLAM control volume $A_\ell^{(nc)}$, respectively. The physics grid cells and CSLAM cells respectively span the element without gaps or overlaps

$$\bigcup_{k=1}^{pg^2} A_k^{(pg)} = \Omega \text{ and } A_k^{(pg)} \cap A_\ell^{(pg)} = \emptyset \quad \forall k \neq \ell, \quad (1)$$

$$\bigcup_{k=1}^{nc^2} A_k^{(nc)} = \Omega \text{ and } A_k^{(nc)} \cap A_\ell^{(nc)} = \emptyset \quad \forall k \neq \ell. \quad (2)$$

The overlap areas between the k -th physics grid cell and CSLAM cells is denoted

$$A_{k\ell} = A_k^{(pg)} \cap A_\ell^{(nc)}, \quad (3)$$

so that

$$A_k^{(pg)} = \bigcup_{\ell=1}^{nc^2} A_{k\ell}. \quad (4)$$

This overlap grid is also referred to as an exchange grid.

2.1 Mapping tracers from CSLAM to pg

For mapping tracer state from the CSLAM grid to any physics grid can be done using existing CSLAM technology, i.e. do a high-order shape-preserving reconstruction of mixing ratio and dry air mass inside each CSLAM control volume and integrate those reconstruction functions over the overlap areas [Lauritzen *et al.*, 2010; Nair and Lauritzen, 2010]. This algorithm retains the properties of CSLAM: inherent mass-conservation, mixing ratio shape-preservation and linear-correlation preservation.

In mathematical terms the remapping is given by

$$\Delta M_{\ell}^{(pg)} \Delta A_{\ell} = \sum_{k=1}^{nc^2} \Delta M_{k\ell}^{(nc)} \Delta A_{k\ell}, \quad (5)$$

$$\Delta M_{\ell}^{(pg)} m_{\ell}^{(pg)} \Delta A_{\ell} = \frac{1}{\Delta M_{\ell}^{(pg)}} \sum_{k=1}^{nc^2} [\Delta M m]_{k\ell}^{(nc)} \Delta A_{k\ell}, \quad (6)$$

where

$$\Delta M_{k\ell}^{(nc)} = \frac{1}{\Delta A_{k\ell}} \int_{A_{k\ell}} \Delta M(x, y) dA. \quad (7)$$

$$[\Delta M m]_{k\ell}^{(nc)} = \frac{1}{\Delta A_{k\ell}} \int_{A_{k\ell}} [\Delta M m](x, y) dA. \quad (8)$$

The tendencies from the parameterizations are computed on the physics grid. The tracer tendency in physics grid cell k is denoted $f_k^{(pg)}$. The problem is how to map $f_k^{(pg)}$ to the CSLAM control volumes $f_{\ell}^{(nc)}$ satisfying the following constraints:

1. Local mass-conservation

$$f_k^{(pg)} \Delta p_k^{(pg)} = \cup_{\ell=1}^{nc^2} \Delta A_{k\ell} \Delta p_{\ell}^{(nc)} f_{\ell}^{(nc)}, \quad (9)$$

where $\Delta p_k^{(pg)}$ is the pressure level thickness in physics grid cell k and similarly for $\Delta p_{\ell}^{(nc)}$.

2. **Shape-preservation in mixing ratio:** The forcing on the CSLAM grid should not produce a value smaller than the new physics grid mixing ratio, $m_k^{(pg)} + \Delta t f_k^{(pg)}$ or a value smaller than the existing CSLAM mixing ratios over the overlap areas $m_{k\ell}^{(nc)}$

$$m_k^{(min)} = \min \left(m_k^{(pg)} + \Delta t f_k^{(pg)}, \left\{ m_{k\ell}^{(nc)} | \ell = 1, nc^2 \right\} \right), \quad (10)$$

where Δt is the physics time-step. Similarly for maxima

$$m_k^{(max)} = \max \left(m_k^{(pg)} + \Delta t f_k^{(pg)}, \left\{ m_{k\ell}^{(nc)} | \ell = 1, nc^2 \right\} \right), \quad (11)$$

3. **Linear correlation preservation:** The physics forcing must not disrupt linear tracer correlation between species on the CSLAM grid [see, e.g., Lauritzen and Thuburn, 2012].
4. **Consistency:** A constant mixing ratio tendency, $cnst$, on the physics grid, $f_k^{(pg)} = cnst \forall k$, must result in the same (constant) forcing on the CSLAM grid, $f_{\ell}^{(nc)} = f_k^{(pg)} = cnst \forall \ell$.

To motivate the algorithm that will simultaneously satisfy 1-4 it is informative to discuss how ‘standard’ mapping algorithms will violate one or more of the constraints.

- Conservative remapping:
- Interpolation:

some text about how challenging it is to satisfy 1-3 simultaneously




Figure 1. Indice notation for the (a) $pg2$ grid, (b), $pg3$ grid and (c) their exchange grid. **Peter - do you think you will use this figure?**




Figure 2. Make captions stand-alone while being concise

2.2 Algorithm

In the $pg2$ configuration, mapping the fields to and from the quadrature grid and $pg2$ grid is identical to that described in HL18. As discussed above, in mapping to the physics grid, CAM-SE's Lagrange basis functions are integrated over the $pg2$ control volumes to provide the physics with a volume averaged state. The procedure is accurate to machine precision, conserves thermal energy and dry air mass, and is consistent (i.e., the mapping preserves a constant). The reverse mapping, from the physics grid to the quadrature grid, is done using a tensor-product Lagrange interpolation (see Appendix A in HL18). The Lagrange interpolation is consistent, conserves dry air mass (**Peter, is this true?**), but does not conserve thermal energy. Errors arising from the lack of energy conservation were estimated to be small; about two orders of magnitude less than the energy dissipation due to the dynamical core alone.

The semi-Lagrangian advection of tracers in our $pg2$ configuration is solved on the CSLAM grid. Preserving linear correlations in mapping to and from the CSLAM and $pg2$ grids requires additional considerations; one such problem is depicted schematically in Figure 2. Consider a single element of CSLAM control volumes, containing only a single cell with mixing ratio 1.0, and 0.0 everywhere else (m_l ; Figure 2a). Assume that the mixing ratios mapped to the $pg2$ grid (m_k ; Figure 2b) result in a negative tracer tendency from the physics (f_k ; Figure 2c). The non-zero values of the tendencies for $pg2$ areas overlapping CSLAM grid cells originally containing a value of zero (f_{kl} ; Figure 2d), are driven negative by the mapped tendency (Figure 2e). Preserving linear correlations is

therefore difficult since the mapping between CSLAM and *pg2* is not reversible. Simulations performed without addressing this artifact were found to be in serious error.

Describe algorithm here

Peter - I think the results of the terminator tests should be mentioned here. We could just put in a sentence saying it passes. But I'm assuming that if we don't use the algorithm that weights the tendency by the amount of available mixing ratio, it will fail. If that's the case, we could just do a two panel plot showing the iCLy at day 15 for with and without the algorithm.

2.2.1 Model Configurations

Two model configurations using CESM2.1 (<https://doi.org/10.5065/D67H1H0V>) are chosen to carry out the objectives discussed in Section 1. To test the hypothesis, that the *pg2* grid reduces spurious grid-noise over mountainous regions, a Held-Suarez configuration [*FHS94* compset; *Held and Suarez*, 1994] modified to include real world topography is analyzed. HL18 indicate that this configuration tends to have more grid-noise over steep terrain than in a more complex configuration using CAM, version 6 moist physics [CAM6;], and is therefore a conservative choice for evaluating any change in grid imprinting between *pg3* and *pg2*.

To understand whether the resolved scales of motion are influenced by the use of a coarser resolution physics grid, a suite of aqua-planet simulations [*Neale and Hoskins*, 2000; *Medeiros et al.*, 2016] are carried out over a range of spectral-element grid resolutions, using CAM6 physics (*QPC6* compset). An aqua-planet is an ocean covered planet in perpetual equinox, with fixed, zonally-symmetric sea surface temperatures idealized after present day Earth. In CAM, there is a strong sensitivity of solutions to the physics time-step [Δt_{phys} ; *Williamson and Olson*, 2003; *Williamson*, 2013; *Herrington and Reed*, 2018] and it is unclear how to choose Δt_{phys} across resolutions.

Here, a scaling for Δt_{phys} across resolutions is proposed, based on results of the moist bubble test [*Herrington and Reed*, 2018] using CAM-SE-CSLAM and detailed in Appendix A: . The scaling is linear in grid-spacing and analogous to a CFL criterion,

$$\Delta t_{phys} = \Delta t_{phys,0} \times \frac{N_e}{N_{e,0}} s, \quad (12)$$

where $\Delta t_{phys,0}$ is taken to be the standard 1800s used in CAM-SE-CSLAM at low resolution, $N_{e,0} = 30$ (equivalent to a dynamics grid-spacing of 111.2km). N_e refers to the horizontal resolution of the grid; each of the six sides of the cubed-sphere is divided into $N_e \times N_e$ elements. Throughout the paper, grid resolutions are abbreviated with an *ne* followed by the quantity N_e , e.g., *ne30*.

The only other parameters varied across resolutions are the dynamics time-step, Δt_{dyn} , and explicit numerical dissipation. Δt_{dyn} is set according to the CFL criterion. The spectral element method is not implicitly dissipative, so fourth-order hyper-viscosity operators are applied to the state to suppress numerical artifacts. The scaling of the hyper-viscosity coefficients, ν , across resolutions is defined as,

$$\nu_T = \nu_{vor} = 0.30 \times \left(\frac{30}{N_e} 1.1 \times 10^5 \right)^3 \frac{m^4}{s}, \quad (13)$$

$$\nu_p = \nu_{div} = 0.751 \times \left(\frac{30}{N_e} 1.1 \times 10^5 \right)^3 \frac{m^4}{s}, \quad (14)$$

where subscripts *T*, *vor*, *p*, *div* refer to state variables the operators are applied to, temperature, vorticity, pressure and divergence, respectively. No explicit dissipation of moisture is required since CSLAM numerics are implicitly diffusive.




Figure 3. Mean ω at two model levels in the middle troposphere, in a Held-Suarez configuration outfitted with real world topography. (Left) *ne30pg2* (Middle) *ne30pg3* and (Right) *ne30pg3* with the divergence damping coefficient increased by an order of magnitude. The ω fields are computed a two-year simulation. The data are presented on a raster plot in order to identify individual grid cells

3 Results

3.1 Held-Suarez with Topography

Flow over topography can result in significant grid imprinting using the spectral element method [Lauritzen *et al.*, 2015; Herrington *et al.*, 2018]. Figure 3 shows the results of the Held-Suarez with topography simulations. The middle panel is the vertical pressure velocity, ω , averaged over two years, over the Andes and Himalayan region at two different levels in the mid-troposphere, using the *ne30pg3* grid. The fields are displayed as a raster plot on the physics grid, so that individual extrema, which characterize the flow over the Andes between about $10^\circ - 20^\circ$ S, and the Himalayas between $20^\circ - 30^\circ$ N, may be identified as spurious.

As discussed in Herrington *et al.* [2018], grid imprinting over mountainous terrain tends to occur in regions of weak gravitational stability, causing extrema to extend through the full depth of the troposphere as resolved updrafts and downdrafts. Thus, grid imprinting over mountains may be alleviated through increasing the divergence damping in the model. Figure 3 (right panel) repeats the *ne30pg3* simulation through increasing ν_{div} by an order of magnitude. The spurious noise over the Andes and the Himalayas are damped, and grid point extrema tend to diffuse into neighboring grid cells. The wavenumber-power spectrum of the kinetic energy arising from divergent modes is provided in Figure 4, confirming that divergent modes are significantly damped at higher wavenumbers (greater than 30), by about an order of magnitude relative to the default *ne30pg3* simulation.




Figure 4. Kinetic energy power spectrum arising from divergent modes in *ne30pg3*, *ne30pg2* and *ne30pg3* with the divergence damping coefficient increased by an order of magnitude, in the Held-Suarez with topography simulations. Spectra computed from five months of six-hourly winds.

The ω field of the *ne30pg2* simulation is provided in Figure 3 (left panel). Grid cell extrema over the Andes is less prevalent than in the *ne30pg3* simulation, as seen by the reduction in large magnitude ω (e.g., red grid cells). The spurious oscillations at the foot of the Himalayas appear to have been entirely eliminated. This improvement in grid imprinting is due to the consistent numerical properties of the control volumes in the *pg2* configuration discussed in Section 1. The divergent modes are marginally damped relative to *ne30pg3* for wavenumbers greater than about 50, but are an order of magnitude greater than the enhanced divergence damping *ne30pg3* run (Figure 4).

3.2 Aqua-planets

When the physics and dynamics grids are of a different resolution, it is not clear which grid determines the resolvable scales of motion. This may be tested through comparing *ne30pg2*, which has a physics grid spacing, $\Delta x_{phys} = 166.8\text{km}$, 1.5 times greater than the dynamics grid spacing, $\Delta x_{dyn} = 111.2\text{km}$, to a simulation where both are equal to the physics grid resolution, $\Delta x_{dyn} = \Delta x_{phys} = 166.8\text{km}$ (*ne20pg3*), and a simulation where both are equal to the dynamics resolution, $\Delta x_{dyn} = \Delta x_{phys} = 111.2\text{km}$ (*ne30pg3*; see Table 1). The resolvable scales in the *ne30pg2* solution is expected to be bounded by the *ne30pg3* and *ne20pg3* solutions. Although according to equation 12, Δt_{phys} in the *ne20* simulation is different than that of the *ne30* runs, here it is set to the *ne30* value (1800s) in order to reduce the differences between the configurations, and because lower resolution runs aren't very sensitive to Δt_{phys} (Figure A.3).

Table 1. Δx and Δt for the physics and dynamics in the low resolution simulations

Grid name	Δx_{dyn}	Δt_{dyn}	Δx_{phys}	Δt_{phys}
ne20pg3	166.8km	300s	166.8km	1800s
ne30pg2	111.2km	300s	166.8km	1800s
ne30pg3	111.2km	300s	111.2km	1800s

The Tropical regions are sensitive to horizontal resolution in global atmospheric models primarily due to the large resolution sensitivity of resolved updrafts and downdrafts at hydrostatic scales [Herrington and Reed, 2018]. Figure ?? shows a snapshot of the ω field in the Inter-Tropical Convergence Zone (ITCZ) in the pressure-longitude plane, in the three simulations. ω is overlain with the $\pm 10K/day$ contours of the physics temperature tendencies (white), and a contour outlining the region where parameterized deep convection is active (white). A large regions of the ITCZ is comprised of upward ω balancing the compensating subsidence warming from parameterized deep convection. Large magnitude ω are comprised of resolved updrafts driven by the large heating rates due to stratiform cloud formation, primarily in the middle-to-upper troposphere due to detrainment of moisture by the deep convection scheme [Zhang and McFarlane, 1995]. Downdrafts tend to occur in the lower portion of the Troposphere, due to re-evaporation of condensates produced by the overlying stratiform cloud.

Figure ?? indicates that the resolved updrafts have a characteristic length scale associated with them, which tend to be roughly seven times the grid spacing, and therefore the length scale changes with model resolution [Herrington and Reed, 2018]. This is consistent with the *ne20pg3* transect, indicating wider updrafts compared with the *ne30pg3* transect. A scale analysis of the Poisson equation [Jeevanjee and Romps, 2016] modified for hydrostatic scales shows that the ratio of ω scales due to the Archimedean buoyancy scales, at two different resolutions is,

$$\frac{\omega_1}{\omega_2} = \frac{D_2}{D_1}, \quad (15)$$

where D is the characteristic buoyancy length scale of the resolution, and it is assumed that the magnitude of the buoyancy and the vertical scale of the buoyancy is unchanged or compensating across the two resolutions. This relationship is robust in a simple moist bubble configuration in CAM-SE [Herrington and Reed, 2018] and CAM-SE-CSLAM (Appendix A:), even though the scaling is derived from the dry anelastic equations.

The probability density function (PDF) of upward ω everywhere in the simulations is shown in Figure 5b. Since the component of ω due to buoyancy is determined by the physics temperature tendencies mapped to the GLL grid, $f_T^{(gll)}$, ω is shown on the GLL grid, ω_{gll} . From the figure It is clear that larger magnitude ω_{gll} is more frequent in *ne30pg2*, as compared with *ne20pg3*, and is actually more similar to the *ne30pg3* distribution.

Figure 5a shows the wave-number power spectrum of the temperature tendencies by the physics (hereafter referred to as *forcing*), after mapping to the GLL grid, in the middle-to-upper troposphere where stratiform heating is common. The variance is larger for higher wavenumbers in *ne30pg3* and *ne30pg2*, as compared with *ne20pg3*, consistent with the larger D inferred from Figure ?. A larger D in *ne20pg3* is consistent with equation 15, in that the frequency of large magnitude ω is lower, compared with the *ne30pg2* and *ne30pg3* simulations.

The *ne30pg2* spectra is remarkably similar to the *ne30pg3* spectra (Figure 5a), consistent with their similar distribution of large magnitude ω in Figure 5b. These figures




Figure 5. (Left) Wavenumber-power spectrum of the temperature tendencies from the moist physics, near the 369 hPa level, (Middle) probability density distribution and (Right) the scaled probability density distribution of upward ω everywhere in the model. The scaled distributions are scaled to *ne30pg3* using Δx_{phys} .

suggest that D is very similar between *ne30pg2* and *ne30pg3*. This is further illustrated through scaling the PDF's,

$$P(\omega_s) = \alpha \times P(\omega/\alpha), \quad (16)$$

where $P(\omega_s)$ is the PDF of the scaled ω , ω_s , and α is the ratio of the ω to ω of the target grid resolution. If one assumes D is linear in Δx , then from equation 15, $\alpha = \Delta x_{target}/\Delta x$, where Δx_{target} is the grid spacing of the target resolution. The target resolution is taken here to be Δx of the *ne30pg3* grid.

If D is in fact determined by the physics grid spacing, Δx_{phys} , then one sets $\Delta x = \Delta x_{phys}$ in α for the *ne30pg2* simulation. This scaled PDF, however, severely overestimates the frequency of upward ω in *ne30pg3* (Figure 5c). It is clear from the similarity of the un-scaled PDF's that D is determined by the dynamics grid spacing, Δx_{dyn} . In contrast, the scaled *ne20pg3* PDF agree's quite well with the *ne30pg3* distribution, suggesting that the scale parameter α explains the difference in vertical motion between the two simulations.

The authors have determined that there are two reasons D is determined by the dynamics grid, and not the physics grid, in *ne30pg2*. The first reason, is that the hyperviscosity coefficients are a function of Δx_{dyn} , and therefore the same in the *pg2* and *pg3* simulations. The fourth-order hyperviscosity is rather scale selective, targeting near grid-scale features more so than, e.g., a second-order operator. The difference in Δx_{phys} between *pg2* and *pg3* are small enough that the hyperviscosity renders this distinction somewhat ambiguous. Through increasing ν in *ne30pg2* to *ne20* values, the variance of forcing on the GLL grid is reduced at higher wave-numbers compared to the standard *ne30pg2* run (not shown), indicating that the length scale of the resolvable features, i.e., D , are greater.

The second reason D is determined by Δx_{dyn} , is that high-order mapping of the physics tendencies from *pg2* to the higher-resolution *GLL* and *CSLAM* grids helps to reconstruct scales that are not present on the *pg2* grid. The left panel of Figure 6a shows a close-up of the wavenumber power spectrum of the forcing on the physics grid (dotted) and the *GLL* grid (solid). In *ne30pg3*, the variances are similar, and even damped at higher wavenumbers (larger than 65) on the *GLL* grid compared to the physics grid. Through using high-order mapping in *ne30pg2*, the variance on the *GLL* grid is actually larger than on the *pg2* grid, matching the variance of the *ne30pg3* forcing on the *GLL* grid, mentioned earlier in reference to Figure 5a. Since the adiabatic dynamics is prognosed on the *GLL* grid, the similar forcing variance on the *GLL* grid in *ne30pg2* and *ne30pg3* is consistent with the similar PDFs of ω on the *GLL* grid in the two simulations




Figure 6. (Left) Wavenumber-power spectrum of the temperature tendencies from the moist physics, near the 369 hPa level, and (right) probability density distribution of upward ω , everywhere in the model, for three year-long aqua-planet simulations. Solid lines refer to values of on the dynamics grid, and dashed lines, the values on the physics grid. See text for details regarding the three simulations.

(see Figure 6, right panel). Repeating the *ne30pg2* simulation, but using low-order mapping, i.e., piecewise constant mapping from *pg2* to *CSLAM* and bilinear mapping from *pg2* to *GLL*, the forcing variance on the *GLL* grid is similar, and even slightly less at high wavenumbers than on the *pg2* grid. Following suit, the frequency of large magnitude ω on the *GLL* grid in the low-order run is less compared to the default *ne30pg2* simulation (Figure 6, right panel).

Through using the low-order mapping in *ne30pg2*, and by increasing ν to *ne20* values, the simulation more closely resembles the *ne20pg3* run (Figure 5). In this case, D is more accurately determined by Δx_{phys} , since the scaled PDF matches the *ne30pg3* simulation quite well. Therefore, at low resolution, our default *ne30pg2* configuration does not indicate that the scales of motion are aliased to the resolution of the coarser resolution physics grid, as they more closely resemble the *ne30pg3* solution.

3.2.1 High Resolution

The physics time-step, Δt_{phys} , used for each grid is scaled by the dynamics time-step to prevent time truncation errors at higher resolutions [Herrington and Reed, 2018].

3.2.2 Across Resolutions

4 Conclusions

Mitigating grid-imprinting through increasing the divergence damping coefficient an order of magnitude greater than is required for numerical stability is not ideal from a model development perspective. The hyper-viscosity coefficients are one of the only a handful of free-parameters in CAM-SE to tune the kinetic energy spectrum to match observations [Skamarock et al., 2014; Lauritzen et al., 2018].




Figure 7. As in Figure 5, but for the high resolution simulations. Asterisks indicate that the physics time-step in these simulations are $\Delta t_{phys} = 675s$, which is larger than those used in the default *ne120* grid of $\Delta t_{phys} = 450s$ (see Table ??).




Figure 8. Year long mean values, averaged over $\pm 10^\circ$ latitude.

A: Defining Δt_{phys} across resolutions

Herrington and Reed [2018] developed a moist bubble test, which indicate that time-truncation errors are large at high resolution (roughly $50km$ and less), and may provide insight on a reasonable scaling of Δt_{phys} across resolutions in more complex configurations. In the test a set of non-rotating simulations are initialized with a super-saturated thermal bubble, and the grid spacing and bubble radius are simultaneously reduced by the same factor in each run through varying the planetary radius. The test was designed to mimic the reduction in buoyancy length scales that occur when the model resolution is increased in more complex configurations [*Hack et al.*, 2006; *Herrington and Reed*, 2018].

The moist bubble test is performed with CAM-SE-CSLAM and coupled to the simple condensation routine of *Kessler* [1969] across five different resolutions (pertaining to the *ne30*, *ne40*, *ne60*, *ne80*, and *ne120* grids). The results are expressed as the minimum ω throughout each one day simulation, and shown in Figure A.2. Two sets of simulations are performed with both *pg3* and *pg2*, one with Δt_{phys} determined by equation 12, and an equivalent set of simulations with $\Delta t_{phys} = \Delta t_{phys,0}$ for all resolutions.

Since the diameters of the bubbles, D , are set proportional to Δx_{dyn} , *Herrington and Reed* [2018] has shown that ω converges to the scaling of equation 15 in the limit of small Δt_{phys} , where small Δt_{phys} is defined as $\Delta t_{phys} = \Delta t_{dyn}$, where Δt_{dyn} is the CFL limiting time-step. Equation 15 is overlain as grey lines in Figure A.2, with *ne30* being the reference resolution. The solutions using Δt_{phys} from equation 12 follow the scaling, whereas fixing $\Delta t_{phys} = 1800s$ across resolutions damps the solution relative to the an-




Figure A.1. The magnitude of ω in the *pg3* solutions are systematically larger than the *pg2* solutions, which is primarily a result of the damping effect of integrating the basis functions over a larger control volume.

Figure A.2.

analytical solution, progressively more so at higher resolutions. If Δt_{phys} is too large, the solution has non-negligible error, which is avoided through scaling Δt_{phys} according to equation 12.

It is not clear if the results of the idealized test extend to the results of more complex configurations. To get a handle on whether the test is useful for understanding more realistic configurations, four aqua-planet simulations are performed with CAM6 physics. A pair of *ne30pg2* simulations, one in which Δt_{phys} is set to the appropriate value from equation 12 (1800s), and one where it is set to the Δt_{phys} corresponding to the *ne20* resolution (2700s). Similarly, a pair of *ne120pg2* simulations are performed, one with Δt_{phys} set to the value from equation 12 (450s), one with Δt_{phys} set to the *ne80* value (625s).

Figure A.3 shows the PDFs of ω from a year of six-hourly data in the simulations. At lower resolution, Δt_{phys} has only a very small effect on the solution, near the tail-end of the distributions (Figure A.3a). At high-resolution, values of ω less than about 3 Pa/s are more frequent in the small Δt_{phys} run, with the discrepancy growing more for larger magnitudes of ω (Figure A.3b). These results are similar to the aqua-planet results in *Herrington and Reed* [2018] using a prior version of CAM physics, version 5, and show that solutions are more sensitive to Δt_{phys} at higher-resolution. The progressively larger errors with increasing resolution also manifests in the moist bubble tests, indicating that truncation errors arising from large Δt_{phys} do exist in more complex configurations.




Figure A.3. Probability density distribution of upward ω everywhere in the model in the aqua-planets using the *ne30pg2* grid (Left) and the *ne120pg2* grid (Right). Figure computed for one year of 6-hourly data. The different colors indicate the physics time-steps used in the runs.

References

- Dennis, J. M., J. Edwards, K. J. Evans, O. Guba, P. H. Lauritzen, A. A. Mirin, A. St-Cyr, M. A. Taylor, and P. H. Worley (2012), CAM-SE: A scalable spectral element dynamical core for the Community Atmosphere Model, *Int. J. High. Perform. C.*, 26(1), 74–89, doi:10.1177/1094342011428142.
- Grinstein, F. F., L. G. Margolin, and W. J. Rider (Eds.) (2007), *Implicit Large Eddy Simulation: Computing turbulent fluid dynamics*, Cambridge University Press.
- Hack, J., M. Caron, G. Danabasoglu, K. W. Oleson, C. Bitz, and J. Truesdale (2006), Ccsn-cam3 climate simulation sensitivity to changes in horizontal resolution, *J. Climate*, 19(1), 2269–2289.
- Held, I. M., and M. J. Suarez (1994), A proposal for the intercomparison of the dynamical cores of atmospheric general circulation models, *Bull. Am. Meteorol. Soc.*, 73, 1825–1830.
- Herrington, A., and K. Reed (2018), An idealized test of the response of the community atmosphere model to near-grid-scale forcing across hydrostatic resolutions, *J. Adv. Model. Earth Syst.*, 10(2), 560–575.
- Herrington, A., P. Lauritzen, M. A. Taylor, S. Goldhaber, B. E. Eaton, J. Bacmeister, K. Reed, and P. Ullrich (2018), Physics-dynamics coupling with element-based high-order galerkin methods: quasi equal-area physics grid, *Mon. Wea. Rev.*, doi:10.1175/MWR-D-18-0136.1.
- Herrington, A. R., and K. A. Reed (2017), An explanation for the sensitivity of the mean state of the community atmosphere model to horizontal resolution on aquaplanets, *J. Climate*, 30(13), 4781–4797, doi:10.1175/jcli-d-16-0069.1.
- Jablonowski, C., and D. L. Williamson (2011), The pros and cons of diffusion, filters and fixers in atmospheric general circulation models., in: P.H. Lauritzen, R.D. Nair, C. Jablonowski, M. Taylor (Eds.), *Numerical techniques for global atmospheric models, Lecture Notes in Computational Science and Engineering, Springer, 2010, in press.*, 80.
- Jeevanjee, N., and D. M. Romps (2016), Effective buoyancy at the surface and aloft, *Quart. J. Roy. Meteor. Soc.*, 142(695), 811–820.
- Kessler, E. (1969), On the distribution and continuity of water substance in atmospheric circulations, *Meteorol. Monogr.*, 10(32), 88.

- Lander, J., and B. Hoskins (1997), Believable scales and parameterizations in a spectral transform model, *Mon. Wea. Rev.*, *125*, 292–303., doi:10.1175/1520-0493.
- Lauritzen, P., and J. Thuburn (2012), Evaluating advection/transport schemes using inter-related tracers, scatter plots and numerical mixing diagnostics, *Quart. J. Roy. Met. Soc.*, *138*(665), 906–918, doi:10.1002/qj.986.
- Lauritzen, P. H., R. D. Nair, and P. A. Ullrich (2010), A conservative semi-Lagrangian multi-tracer transport scheme (CSLAM) on the cubed-sphere grid, *J. Comput. Phys.*, *229*, 1401–1424, doi:10.1016/j.jcp.2009.10.036.
- Lauritzen, P. H., J. T. Bacmeister, P. F. Callaghan, and M. A. Taylor (2015), Ncar global model topography generation software for unstructured grids, *Geoscientific Model Development Discussions*, *8*(6), 4623–4651, doi:10.5194/gmdd-8-4623-2015.
- Lauritzen, P. H., M. A. Taylor, J. Overfelt, P. A. Ullrich, R. D. Nair, S. Goldhaber, and R. Kelly (2017), CAM-SE-CSLAM: Consistent coupling of a conservative semi-lagrangian finite-volume method with spectral element dynamics, *Mon. Wea. Rev.*, *145*(3), 833–855, doi:10.1175/MWR-D-16-0258.1.
- Lauritzen, P. H., R. Nair, A. Herrington, P. Callaghan, S. Goldhaber, J. Dennis, J. T. Bacmeister, B. Eaton, C. Zarzycki, M. A. Taylor, A. Gettelman, R. Neale, B. Dobbins, K. Reed, and T. Dubos (2018), NCAR CESM2.0 release of CAM-SE: A reformulation of the spectral-element dynamical core in dry-mass vertical coordinates with comprehensive treatment of condensates and energy, *J. Adv. Model. Earth Syst.*, doi:10.1029/2017MS001257.
- Medeiros, B., D. L. Williamson, and J. G. Olson (2016), Reference aquaplanet climate in the community atmosphere model, version 5, *J. Adv. Model. Earth Syst.*, *8*(1), 406–424, doi:10.1002/2015MS000593.
- Nair, R. D., and P. H. Lauritzen (2010), A class of deformational flow test cases for linear transport problems on the sphere, *J. Comput. Phys.*, *229*, 8868–8887, doi:10.1016/j.jcp.2010.08.014.
- Neale, R. B., and B. J. Hoskins (2000), A standard test for agcms including their physical parametrizations: I: the proposal, *Atmos. Sci. Lett.*, *1*(2), 101–107, doi:10.1006/asle.2000.0022.
- Neale, R. B., C.-C. Chen, A. Gettelman, P. H. Lauritzen, S. Park, D. L. Williamson, A. J. Conley, R. Garcia, D. Kinnison, J.-F. Lamarque, D. Marsh, M. Mills, A. K. Smith, S. Tilmes, F. Vitt, P. Cameron-Smith, W. D. Collins, M. J. Iacono, R. C. Easter, S. J. Ghan, X. Liu, P. J. Rasch, and M. A. Taylor (2012), Description of the NCAR Community Atmosphere Model (CAM 5.0), *NCAR Technical Note NCAR/TN-486+STR*, National Center of Atmospheric Research.
- Randall, D., M. Khairoutdinov, A. Arakawa, and W. Grabowski (2003), Breaking the cloud parameterization deadlock, *Bulletin of the American Meteorological Society*, *84*(11), 1547–1564.
- Shutts, G. (2005), A kinetic energy backscatter algorithm for use in ensemble prediction systems, *Quart. J. Roy. Meteorol. Soc.*, *131*, 3079–3102.
- Skamarock, W. (2011), Kinetic energy spectra and model filters, in: P.H. Lauritzen, R.D. Nair, C. Jablonowski, M. Taylor (Eds.), *Numerical techniques for global atmospheric models*, *Lecture Notes in Computational Science and Engineering*, Springer, 80.
- Skamarock, W. C., S.-H. Park, J. B. Klemp, and C. Snyder (2014), Atmospheric kinetic energy spectra from global high-resolution nonhydrostatic simulations, *Journal of the Atmospheric Sciences*, *71*(11), 4369–4381, doi:10.1175/JAS-D-14-0114.1.
- Taylor, M., J. Edwards, and A. St-Cyr (2008), Petascale atmospheric models for the community climate system model: new developments and evaluation of scalable dynamical cores, *J. Phys.: Conf. Ser.*, *125*, doi:10.1088/1742-6596/125/1/012023.
- Wedi, N. P. (2014), Increasing horizontal resolution in numerical weather prediction and climate simulations: illusion or panacea?, *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, *372*(2018), doi:10.1098/rsta.2013.0289.

- Williamson, D. L. (1999), Convergence of atmospheric simulations with increasing horizontal resolution and fixed forcing scales, *Tellus A*, 51, 663–673, doi:10.1034/j.1600-0870.1999.00009.x.
- Williamson, D. L. (2013), The effect of time steps and time-scales on parametrization suites, *Quart. J. Roy. Meteor. Soc.*, 139(671), 548–560, doi:10.1002/qj.1992.
- Williamson, D. L., and J. G. Olson (2003), Dependence of aqua-planet simulations on time step, *Q. J. R. Meteorol. Soc.*, 129(591), 2049–2064.
- Zhang, G., and N. McFarlane (1995), Sensitivity of climate simulations to the parameterization of cumulus convection in the canadian climate center general-circulation model, *ATMOSPHERE-OCEAN*, 33(3), 407–446.