

# Parameterized convection, grid-scale clouds and resolution sensitivity in the Community Atmosphere Model

Adam R. Herrington\* & Kevin A. Reed

School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY 11794

\*Correspondence to: [adam.herrington@stonybrook.edu](mailto:adam.herrington@stonybrook.edu)

## This paper describes...

**Key Words:** Climate models, physical parameterizations, resolution sensitivity

*Received ...*

## 1. Introduction

An increasing number of Atmospheric General Circulation Models (AGCMs) are being developed to maximize efficiency on massively parallel systems, permitting regionally-refined high-resolution, or even globally high-resolution weather ( $\Delta x \leq 5$  km) and climate ( $\Delta x \leq 50$  km) simulations (Satoh *et al.* 2008; Skamarock *et al.* 2012; Zängl *et al.* 2014; Harris *et al.* 2016; Ullrich *et al.* 2017; Lauritzen *et al.* 2018). These models are built using unstructured meshes that while allows for substantial grid flexibility, would require physical parameterizations (*physics*) that behave consistently as the truncation scale of the model changes with different grid resolutions, referred to as scale-aware physics. The most common approach towards developing scale-aware physics is through the lens of limited area, large-eddy simulations (e.g., Plant and Craig 2008; Arakawa and Wu 2013; Song and Zhang 2018). Through subsequently filtering large-eddy solutions to lower-resolution grids, a relationship between first- and higher-order moments (Germano 1992) may be understood and ultimately parameterized as a function of grid resolution. While this approach is likely necessary for developing scale-aware physics, it is not sufficient. The equations of motions have inherent scale dependencies, with the properties of dynamical modes a function of native grid resolution (Orlanski 1981; Weisman *et al.* 1997; Pauluis and Garner 2006; Jeevanjee and Romps 2016). Scale-aware physics should also recognize these native grid dependencies.

The sensitivity of the Community Atmosphere Model (CAM; Neale *et al.* 2012), and its predecessor, the Community Climate Model (CCM) to resolution (*resolution* refers to *horizontal resolution*, hereafter) is well documented through convergence studies (Kiehl and Williamson 1991; Williamson *et al.* 1995; Williamson 2008; Rauscher *et al.* 2013; Zarzycki *et al.* 2014; Herrington and Reed 2017). CAM/CCM is a well supported climate model, but despite thirty years of continual model development, there are robust sensitivities to resolution that have persisted in all versions of the model. This study argues that a unifying cause, the inherent scale sensitivities of the underlying dynamical equations, can explain the robust responses to resolution that occur in CAM/CCM, since it is difficult to conceive that inevitable responses to native grid resolution could be ignored in the pursuit of scale-aware physics.

In CAM/CCM, the atmosphere progressively dries with increasing resolution, seen through a reduction in simulated total precipitable water (Kiehl and Williamson 1991; Williamson *et al.* 1995; Williamson 2008; Rauscher *et al.* 2013; Zarzycki *et al.* 2014; Herrington and Reed 2017), which typically, but not always (see Williamson *et al.* 1995; Zarzycki *et al.* 2014), coincides with a reduction in cloud cover. Kiehl and Williamson (1991) and Williamson *et al.* (1995) suggested that the drying of the atmosphere is due to greater magnitude resolved vertical velocities with increasing resolution, with greater subsiding motion increasing the export of dry air from the upper troposphere. This mechanism is consistent with an analysis of moisture budgets in CAM, version 4 (CAM4; Neale *et al.* 2010) across multiple resolutions (Yang *et al.* 2014; Herrington and Reed 2017).

It is well known that the magnitude of vertical velocities increase with resolution in atmospheric models. While the cause of this sensitivity has been established for large-eddy simulations (see Jeevanjee 2017, and references therein), only recently has the vertical velocity field in AGCMs and their sensitivity to resolution received attention (Donner *et al.* 2016; O'Brien *et al.* 2016), albeit with conflicting explanations (Rauscher *et al.* 2016; Herrington and Reed 2018). To generalize the relationship between vertical velocity and resolution, let  $\alpha$  refer to the ratio of  $W_0$ , the vertical velocity scale of some reference grid spacing  $\Delta x_0$ , to  $W$ , the vertical velocity scale of any  $\Delta x$ . A power law for  $\alpha^{-1}$  in  $\Delta x$  is then,

$$\alpha^{-1} = \frac{W}{W_0} = \left( \frac{\Delta x}{\Delta x_0} \right)^n, \quad (1)$$

where  $n$  is the power law exponent.

Rauscher *et al.* (2016) derive an estimate  $n = b - 1$  by combining a scale analysis of the continuity equation with a power law representation  $\Delta x^{2b}$  of the second-order structure function of the horizontal wind. Strictly speaking,  $\Delta x$  here refers to the distance between two points for which the velocity increment is computed in the structure function, but with this distance set to the model grid-spacing. Observations indicate that  $b = \frac{1}{3}$  for scales less than about 1000 km (Cho *et al.* 1999), which according to the Weiner–Khinchin theorem  $-(2b + 1) = -\frac{5}{3}$  is equal to the slope of the kinetic energy spectrum, and supported by observations of mesoscale flow (Nastrom and Gage 1985). Rauscher *et al.* (2016)

argue that the  $-\frac{5}{3}$  slope being common in both observations and models provides an emergent constraint for  $b = \frac{1}{3}$  and  $n = -\frac{2}{3}$ .

In large-eddy simulations, the sensitivity of vertical velocities to resolution is adequately explained by a scale analysis of the dynamical equations (Weisman *et al.* 1997; Pauluis and Garner 2006; Jeevanjee and Romps 2016). For hydrostatic scales relevant to AGCMs, a scale analysis of the Poisson equation gives  $W \propto D^{-1}$ , where  $D$  is the horizontal scale of a buoyancy perturbation driving vertical motion (Herrington and Reed 2018). In CAM aqua-planet simulations, the largest source of buoyancy is from grid-scale cloud formation, whose horizontal extents are set by the effective resolution of the model (i.e., some multiple of  $\Delta x$ ), indicating  $n = -1$  (Herrington and Reed 2018). Herrington and Reed (2017) has shown that the  $n = -1$  scaling does not explain the behavior of CAM4 in a convergence experiment, but follow-up studies (Herrington and Reed 2018; Herrington *et al.* 2019) indicate that the inadequacy of the  $n = -1$  scaling is not definitive, due to time-truncation errors associated with fixing the physics time-step ( $\Delta t_{phys}$ ) across resolutions in that study.

Another robust response of the CAM/CCM lineage to resolution is an increase in grid-scale precipitation rates at the expense of parameterized convective precipitation rates. Grid-scale precipitation is formed from grid-scale clouds, but is often referred to in the literature as stratiform precipitation and stratiform clouds, and used hereafter. The behavior of the difference precipitation routines with resolution is shown in Figure 1, which is a bar-graph of the climatological, global mean stratiform and convective precipitation rates in prior CAM/CCM convergence studies. The propensity for precipitation rates to shift from the convection scheme to the stratiform scheme with resolution has been documented in other AGCMs as well (Pope and Stratton 2002; Terai *et al.* 2018), but none have given a satisfactory explanation for this sensitivity. The studies of Kiehl and Williamson (1991), Williamson *et al.* (1995) and Williamson (2013) indicate that the tendency to reduce  $\Delta t_{phys}$  with resolution would by itself reduce the convective precipitation rates, however Figure 1 (top row) indicates that convergence studies with fixed  $\Delta t_{phys}$  still show a reduction in convective precipitation rates with resolution.

In this study, a convergence experiment using CAM, version 6 (CAM6; [https://ncar.github.io/CAM/doc/build/html/users\\_guide/index.html](https://ncar.github.io/CAM/doc/build/html/users_guide/index.html)) is carried out and analyzed in detail. It is shown that the resolution sensitivity of vertical velocities are well described with  $n = -1$  in equation (1), provided  $\Delta t_{phys}$  is defined in a way that avoids large truncation errors across resolutions. The reduction in convective precipitation rates with resolution in CAM6 is shown to result from the greater magnitude subsiding motion, creating a more stable atmosphere in which the criterion for parameterized convection occurs less often. The feedback of the resolved vertical motion on the physics indicates that the root cause of resolution sensitivity in CAM arises from the sensitivity of resolved dynamical modes to native grid resolution. Section 2 describes CAM6 and details the convergence experiment. Section 3 contains a thorough analysis of the CAM6 simulations and Section 4 provides some discussion and conclusions.

## 2. Methods

### 2.1. Dynamical Core

This study uses the spectral-element dynamical core option of Community Atmosphere Model (CAM-SE; Dennis *et al.* 2012), coupled with a mass conserving, semi-Lagrangian advection method for accelerated multi-tracer transport (CSLAM; Lauritzen *et al.* 2017), and dry-mass vertical coordinate with comprehensive treatment of moisture and energy (Lauritzen *et al.* 2018). The dry



**Figure 1.** Bar-graph of the convective (solid) and stratiform (white) climatological precipitation rates in prior CAM/CCM convergence studies. Each window contains a single convergence study, with identical x-axis; the approximate grid resolution. Colors indicate the model configuration; January ensemble (black) and aqua-planet configurations with SST profiles *QOBS* (blue) and *CNTL* (red) after Neale and Hoskins (2000). Studies included in this figure are Kiehl and Williamson (1991) (CCM1), Williamson *et al.* (1995) (CCM2), Williamson (2008) (CAM3), Rauscher *et al.* (2013); Zarzycki *et al.* (2014); Herrington and Reed (2017) (CAM4), Zarzycki *et al.* (2014) (CAM5) and this study (CAM6). CCM2\* refers to the modified parameter experiment of Williamson *et al.* (1995), where parameters vary with resolution to reduce the dependence of cloud fraction on resolution.

dynamics are solved using the high-order, momentum, mass and energy conserving spectral element method (Taylor and Fournier 2010), with the elements defined by a cubed-sphere grid. The notation for the horizontal grid resolution is an ‘ne’ followed by the number of elements making up an edge of one cubed-sphere face, e.g., *ne30*. Hyper-viscous  $\nabla^4$  explicit numerical dissipation is applied to temperature, dry pressure thickness, rotational and divergent winds (Lauritzen *et al.* 2018). CSLAM tracer transport uses a finite volume grid constructed from the cubed-sphere of elements, and contains the same degrees of freedom as the dry dynamics.

### 2.2. Physical Parameterizations

The physics are evaluated on the finite-volume CSLAM grid, and the tendencies mapped back to the spectral element grid. The coupled system, referred to as CAM-SE-CSLAM, conserves energy, mass and preserves linear correlations between two reactive species to within machine precision (Herrington *et al.* 2018). A coarser physics grid, containing  $\frac{5}{9}$  fewer degrees of freedom than the dynamical core grid is also available as part of the CAM-SE-CSLAM package (Herrington *et al.* 2019). This lower-resolution physics grid is used in this study, but only as a member of a perturbed parameter ensemble and not in the default convergence experiment. The dynamics time-step is subcycled within a longer physics time-step  $\Delta t_{phys}$ , and the temperature and momentum increments from the physics are divided by the number of subcycles and added to the dynamical core at the beginning of each subcycle. The full moisture increment from the physics is applied only at the start of the first subcycle to conserve tracer mass (*ftype* = 2 option in Lauritzen and Williamson 2019).

The simulations use the CAM6 physics package. The Cloud Layers Unified by Binormals (CLUBB Golaz *et al.* 2002; Bogenschutz *et al.* 2013) is an assumed probability distribution function (PDF) high-order closure model that handles shallow convection, planetary boundary layer mixing and cloud

macrophysics. The macrophysics are coupled with a two-moment bulk cloud microphysics scheme with prognostic precipitation (Gettelman *et al.* 2015), and microphysics are coupled with a three mode Modular Aerosol Model (Liu *et al.* 2012). The combined macrophysics/microphysics routines generate stratiform precipitation from stratiform clouds. Deep convection is parameterized using a quasi-equilibrium mass flux scheme (Zhang and McFarlane 1995) and an entraining plume model (referred to as the dilute convective available potential energy, or *dilute CAPE* hereafter; Raymond and Blyth 1992; Neale *et al.* 2008) is used as a convective trigger (convection occurs if dilute CAPE  $\geq 70$  J/kg), and for closing the mass fluxes in the cloud ensemble. The deep convection scheme also parameterizes convective momentum transport (Richter and Rasch 2008).

### 2.3. Experimental Design

The convergence experiment is performed in an aqua-planet configuration (Neale and Hoskins 2000; Medeiros *et al.* 2016), an all ocean planet with fixed, zonally symmetric sea surface temperatures modeled after present day Earth (*QOBS* in Neale and Hoskins 2000). The aqua-planets are in a perpetual equinox, and aerosols are largely absent from the simulations. Each simulation is ran for one simulated year. Six different horizontal grids are used in this study, which are provided in Table 1. In addition to the six simulations used in the convergence experiment, an ensemble of 24 simulations containing different model parameters (e.g., using the lower resolution physics grid) and across different resolutions are ran for year in order to increase confidence in the assessment of resolution sensitivity in this study. All analyses exclude the first month of the simulations, and are computed on their native grids unless otherwise stated.

The horizontal hyper-viscosity operators  $\nu$  vary with resolution after Herrington *et al.* (2019), also provided in Table 1. The values of  $\nu$  are a factor 2.5 greater for divergence damping and are not shown.  $\Delta t_{phys}$  is chosen to scale with resolution, in proportion to the grid spacing,

$$\Delta t_{phys} = \Delta t_{phys,0} \times \frac{n_{e,0}}{n_e} s, \quad (2)$$

where  $\Delta t_{phys,0}$  is taken to be the standard 1800 s used in CAM-SE-CSLAM for the standard climate resolution,  $n_{e,0} = 30$  (equivalent to an average equatorial grid spacing  $\Delta x = 111.2$  km). This scaling was chosen to avoid large time-truncation errors in a rising moist bubble test (Appendix A in Herrington *et al.* 2019), and it is understood that this choice of  $\Delta t_{phys}$  will likely lead to greater resolution sensitivity (Williamson 2008). The convective time-scale in the deep convection scheme is fixed at 3600 s in all simulations.

## 3. Results


Table 1 provides globally averaged, climatological metrics for the CAM6 simulations which are typically published in CAM/CCM convergence studies. Total precipitable water, total cloud fraction and deep convective precipitation rate decreases, while stratiform precipitation increases, monotonically with resolution (also shown in Figure 1). Resolution sensitivity in CAM6 is similar to all prior versions of the model.

### 3.1. Vertical Velocities and Resolution

The PDF of negative, or upward vertical pressure velocities  $\omega$  in the aqua-planets is shown in Figure 2a. The magnitude of upward  $\omega$  increases monotonically with resolution, with positive, or downward  $\omega$  behaving similarly (not shown). This monotonic



**Figure 2.** Probability density distribution of the upward vertical pressure velocities  $\omega$  computed everywhere in the model from six-hourly output over the entirety of the year-long simulations. (a) Values on their native grid (solid) and values remapped to the  $ne20$  grid (dotted), (b) values on their native grid, scaled to the  $ne120$  resolution.



**Figure 3.** Components of the climatological, global mean vertical pressure velocity, (a)  $\langle f_u \rangle \langle \omega_u \rangle$  and (b)  $\langle f_d \rangle \langle \omega_d \rangle$ . Grey crosses are for the 24 member perturbed parameter ensemble.

increase in the magnitude of  $\omega$  is evident even after remapping all the model output to a common grid ( $ne20$ ; dotted curves in Figure 2a).

The PDF's may be scaled to the highest-resolution resolution grid through  $P(\omega)_s = \alpha P(\omega/\alpha)$ , where  $\alpha$  is the scale factor from equation 1, and setting  $\Delta x_0$  to the  $ne120$  grid-spacing. Figure 2b shows the scaled PDF's using  $n = -1$  in equation 1. The scaled PDF's all collapse onto the high-resolution reference, indicating that the power-law with  $n = -1$  explains to first-order the variation in vertical velocity with resolution in the aqua-planet simulations.

Changes to the vertical velocity field can be further understood through decomposing the mass weighted vertical mean  $\omega$  into upward and downward components,

$$\langle \omega \rangle = \langle f_u \rangle \langle \omega_u \rangle + \langle f_d \rangle \langle \omega_d \rangle, \quad (3)$$

where  $\langle f_x \rangle$  and  $\langle \omega_x \rangle$  refers to the vertical mass fraction  $\left( \frac{\int dp_x}{\int dp} \right)$  and the  $x$  component of the mass weighted vertical mean of  $\omega$   $\left( \frac{\int \omega_x dp_x}{\int dp_x} \right)$ , respectively, subscript  $u$  refers to upward motion and  $d$ , downward motion.


The global mean, climatological components  $\langle f_u \rangle \langle \omega_u \rangle$  and  $\langle f_d \rangle \langle \omega_d \rangle$  are provided in Figure 3a,b for the aqua-planet simulations. The magnitude of both  $\langle f_u \rangle \langle \omega_u \rangle$  and  $\langle f_d \rangle \langle \omega_d \rangle$  increase monotonically with resolution, and are equal and opposite, which is a requirement of mass conservation in the model and a convenient check of the calculation. While  $\langle f_d \rangle$  is about 25% larger than  $\langle f_u \rangle$  in all simulations, the vertical mass fractions varies by only few percent with resolution, and so the monotonic behavior of  $\langle f_x \rangle \langle \omega_x \rangle$  with resolution is primarily from variations in  $\langle \omega_x \rangle$  (not shown).

### 3.2. Vertical Velocities and Deep Convective Precipitation

The large increase in magnitude of the upward and downward vertical velocities with resolution may be expected to impact

Table 1. Experimental design and global mean climatologies.

Variable	ne20	ne30	ne40	ne60	ne80	ne120
$\Delta x$ (km)	166.8	111.2	83.4	55.6	41.7	27.8
$\nu$ ( $m^4/s$ )	$1.5 \times 10^{15}$	$4.0 \times 10^{14}$	$1.5 \times 10^{14}$	$4.0 \times 10^{13}$	$1.5 \times 10^{13}$	$4.0 \times 10^{12}$
$\Delta t_{phys}$ (s)	2700	1800	1350	900	675	450
Total Cloud Fraction	0.844	0.835	0.824	0.810	0.804	0.800
Total Precipitable Water (mm)	23.31	23.01	22.62	22.25	21.93	21.72
Convective Precipitation (mm/day)	1.91	1.83	1.68	1.47	1.29	1.08
Stratiform Precipitation (mm/day)	1.26	1.42	1.60	1.85	2.05	2.22




**Figure 4.** Scatter plot of global mean, climatological  $\langle f_d \rangle \langle \omega_d \rangle$  and  $FREQZM$ , and the fitted linear regression which has a Pearson's R-value = 0.99, using all 27 simulations. Grey crosses are for the 24 member perturbed parameter ensemble runs.

the behavior of other model components. Curiously, there is an excellent negative correlation (Pearson's R-value = 0.99,  $N = 27$ ) between the global mean, climatological  $\langle f_d \rangle \langle \omega_d \rangle$  and a measure of the activity of the [Zhang and McFarlane \(1995\)](#) deep convection scheme (referred to as the *ZM scheme* hereafter), global mean, climatological  $FREQZM$  (Figure 4). At any given grid-point and time-step,  $FREQZM$  is a binary variable: one if the ZM scheme is active, zero if it is not. Time mean  $FREQZM$  is therefore the fraction of the model time that the ZM scheme is triggered, i.e., dilute CAPE exceeds  $\geq 70$  J/kg. The relationship indicates that model simulations with greater subsidence also have less convective activity.

To further understand this relationship, a logistic regression between  $\langle f_d \rangle \langle \omega_d \rangle$  and  $FREQZM$  is performed for each grid column within each of the simulations. Logistic regression uses an iterative method to fit a continuous variable predictor,  $x$  to a binary predictand  $p$  through the exponential ([Wilks 2011](#)),

$$p = \frac{\exp[b_0 + b_1 x]}{1 + \exp[b_0 + b_1 x]}, \quad (4)$$

where  $b_0$  and  $b_1$  are the shape parameters of the exponential. The predictor is then the instantaneous  $\langle f_d \rangle \langle \omega_d \rangle$  of a grid column, and the predictand the binary  $FREQZM$ . The assumption is then that subsidence is the independent variable, which the authors believe is reasonable since the environment of subsiding regions is generally stable, and the ZM scheme is modulated by a stability calculation, dilute CAPE. Grid column regressions that are statistically significant at the 95% level using a log-likelihood test ([Wilks 2011](#)) are retained for analysis. Since the aqua-planets have zonally symmetric boundary conditions, there is a zonally



**Figure 5.** Zonal mean (a) R-values and (b) sensitivity parameter in the logistic regression, (c) climatological surface latent heat fluxes and (d) drizzle fraction.

varying structure in the goodness of fit (R-value) and parameter  $b_1$  (hereafter referred to as the sensitivity parameter; Figure 5a,b).

The zonal mean R-values indicate the greatest goodness of fit in the  $\pm 10^\circ$  latitude band, hereafter referred to as the deep tropics. Figure 5c shows the climatological, zonal-mean latent heat flux in the simulations, which is expected to contribute positively to the component of dilute CAPE associated with the thermodynamic state of boundary layer parcels ([Zhang 2002](#)). In the deep tropics, latent heat fluxes are small, and the sensitivity parameter is large and negative (Figure 5b), consistent with the idea that subsiding motion actively depresses dilute CAPE and the activity of the ZM scheme in the simulations. The sensitivity parameter becomes less negative in the deep tropics with resolution, likely due to the greater magnitude  $\langle f_d \rangle \langle \omega_d \rangle$  with resolution, which requires a lower sensitivity parameter to predict the binary  $FREQZM$ . The R-values generally decrease with resolution indicating that there is degradation in the relationship with resolution.

Table 2 shows the fractional contribution of the deep tropics to the climatological, global mean change in convective precipitation with resolution. This is a reflection of changes in the partitioning of the Intertropical Convergence Zone (ITCZ) from convective to stratiform precipitation with resolution. The table indicates that a majority (60 – 70%) of the reduction in convective precipitation with resolution is from changes within the deep tropics (except



Table 2. Fractional contribution of latitude bands  $\pm 10^\circ$  and  $\pm 15^\circ$  to changes in global mean precipitation with resolution. The grid headers refer to differences with respect to the next lowest grid resolution, e.g.,  $ne30 = ne30 - ne20$ ,  $ne40 = ne40 - ne30$ , etc..., and computed through remapping all data to the  $ne20$  grid.

Variable	$ne30$	$ne40$	$ne60$	$ne80$	$ne120$
$\pm 10^\circ$ (17.6% of global area)					
Convective Precipitation	-0.58	0.62	0.66	0.72	0.70
Stratiform Precipitation	0.55	0.63	0.69	0.67	0.41
$\pm 15^\circ$ (25.8% of global area)					
Convective Precipitation	0.22	0.75	0.73	0.79	0.72
Stratiform Precipitation	0.46	0.64	0.71	0.70	0.49

in going from  $ne20$  to  $ne30$ , where deep tropical convective precipitation rates increase with resolution due to an extremely wide double-ITCZ in the  $ne20$  simulation that spans well outside of  $\pm 10^\circ$  latitude). Expanding the latitude boundaries marginally to  $\pm 15^\circ$ , roughly 75% of the changes in convective precipitation with resolution occur in this region (again, ignoring  $ne30 - ne20$ ; Table 2). Taken together, the region with the largest change in convective precipitation with resolution is also the region where the logistic regression indicates that subsiding motion is most skillful at depressing the activity of the convection scheme.

To estimate the dilute CAPE values associated with subsiding motion in the deep tropics, temperature and moisture profiles are conditionally sampled depending on whether  $\langle \omega \rangle$ , the mass-weighted vertical integral of  $\omega$ , is positive or negative, indicating predominantly subsiding or ascending grid columns. The time mean temperature and moisture profiles of subsiding and ascending regions are then used to compute the dilute CAPE used in the ZM scheme, offline.

Figure 6a shows the dilute CAPE values associated with mean conditions for ascending, descending and all grid columns in the deep tropics, with resolution. The ascending regions are associated with larger values of dilute CAPE ( $> 180$  J/kg) relative to subsiding regions ( $< 110$  J/kg), and the dilute CAPE value computed for mean conditions over the entire deep tropics decreases monotonically with resolution, consistent with the reduction in convective activity with resolution. The space-time weights associated with ascending and descending grid columns in the deep tropics vary drastically with resolution (Figure 6b). The subsiding (ascending) space-time weights change from 0.32 (0.68) at  $ne20$ , monotonically increasing (decreasing) with resolution to 0.51 (0.49) in  $ne120$ . The increasing occurrence of stable, subsiding grid columns with resolution results in a reduction in CAPE for the entire deep tropics, verified by the similar CAPE values derived through taking the weighted sum of the ascending/descending CAPE values (grey crosses in Figure 6a).

Figure 7 shows the time mean temperature and specific humidity profiles of subsiding grid cells in the deep tropics, expressed as anomalies from the mean temperature and specific humidity of the entire deep tropics. The mean profiles of subsiding regions have an anomalous warming layer in the 600 – 800 hPa layer and an anomalous moisture deficit throughout the entire column. This warming and drying patterns is consistent with the effects of subsidence, whose motion produces adiabatic warming and advection of drier, upper-troposphere air downward. Both warming and drying the environment oppose the growth of dilute CAPE through reducing parcel buoyancy; warming the environment relative to the parcel temperature reduces parcel buoyancy (Zhang 2002), and mixing drier environmental air into a rising air parcel reduces the moisture available to warm the parcel through latent heating (Raymond and Blyth 1992).

An experiment is carried out to unravel the contributions of warming and drying to dilute CAPE values of subsiding grid columns. Through computing dilute CAPE using the mean




Figure 6. (a) Dilute CAPE computed from time mean temperature and moisture profiles of ascending (red), subsiding (blue) and all grid columns (black) in the deep tropics ( $\pm 10^\circ$  latitude), and (b) space-time weights of ascending (red) and descending (blue) grid columns in the deep tropics. (c) Dilute CAPE computed for ascending/descending grid columns, but using the mean temperature profile for the entire deep tropics, (d) Same as (c) but using the mean moisture profile for the entire deep tropics. Grey crosses in (a) are dilute CAPE derived from the sum of the products of space-time weights with the dilute CAPE values of ascending/descending grid columns.




Figure 7. Time mean (a) temperature and (b) specific humidity profiles of subsiding grid cells in the deep tropics ( $\pm 10^\circ$  latitude) in the convergence experiment, presented as anomalies from the mean temperature and specific humidity of the entire deep tropics in each simulation.

specific humidity of subsiding regions in the deep tropics, but setting the temperature profile to the mean profile for the entire deep tropics, one can isolate the role of changes in moisture in subsiding regions on the dilute CAPE calculation, while doing the reverse isolates the influence of changes in temperature. Figure 7c shows the influence of moisture of subsiding/ascending regions on dilute CAPE, and Figure 7d the influence of temperature. The first order cause for the reduction of dilute CAPE is clearly due to the reduction (increase) in moisture in subsiding (ascending) region, while temperature has a smaller, second order influence on the dilute CAPE changes. This analysis indicates that the drying of the deep tropics with resolution manifests through an increase in occurrence of subsiding motion with resolution, which reduces the frequency the ZM scheme is active through reducing the frequency the dilute CAPE threshold for convection occurs.

### 3.3. Vertical Velocities and Shallow Convective Precipitation

AGCMs are known to suffer from producing too much light rain, or drizzle, relative to observations (Dai 2006). Figure 5d shows the fraction of ZM precipitation that is less  $\leq 5$  mm/day in the simulations, which stubbornly persists at 70% in the region






Figure 8



**Figure 9.** (a) Dilute CAPE computed from time mean temperature and moisture profiles of ascending (red), subsiding (blue) and all grid columns (black) in the subtropics ( $\pm 10^\circ - 30^\circ$ ) latitude band, and (b) space-time weights of ascending (red) and descending (blue) grid columns in the subtropics. Grey crosses in (a) are dilute CAPE derived from the sum of the products of space-time weights with the dilute CAPE values of ascending/descending grid columns.

between  $10^\circ - 20^\circ$  latitude at all resolutions. Figure ?? shows a snapshot of  $\omega$  in the longitude-pressure plane at  $\sim 18^\circ$  latitude in the *ne30* simulations, and overlain by an isoline delineating where the ZM mass fluxes are fairly active. The ZM mass fluxes rarely extend above the 800 hPa level, and often occurs with abundant subsiding motion aloft. In this subtropical region, there are opposing influences on dilute CAPE, a global maximum in latent heat fluxes (Figure 5c) which increases dilute CAPE from below, and subsidence, opposing dilute CAPE from above. It seems plausible that these two opposing influences on stability translates into small values of dilute CAPE, but still large enough to trigger the ZM scheme. The ZM scheme, properly a deep convection scheme, is operating as a shallow convection scheme in the subtropics and produces drizzle.

Figure 8 shows the dilute CAPE values computed from mean temperature and moisture of subsiding and ascending regions in the  $\pm(10^\circ - 30^\circ)$  latitude bands, hereafter referred to as the subtropics. The spread in dilute CAPE between ascending and descending grid columns is much smaller than for the deep tropics (Figure 6a), the dilute CAPE computed from the mean temperature and moisture over the entire subtropics varies much less with resolution compared with the deep tropics ( $\sim 20$  J/kg across all resolutions, compared with  $\sim 80$  J/kg in the deep tropics). Unlike the deep tropics, the reduction in dilute CAPE with resolution is not from an increase in occurrence of subsiding grid columns, since the space-time weights of ascending/descending motion are generally invariant with resolution (Figure 8b). The decline in dilute CAPE with resolution in the subtropics is a result of a greater magnitude of downward motion, but this sensitivity is not sufficient to suppress the occurrence of drizzling in the subtropics.



**Figure 10.** Precipitation rates vs. upward moisture flux at the 850 hPa level. Solid lines refer to median stratiform (green), convective (blue) and total (magenta) precipitation rates conditional on bins of the moisture flux, and shaded regions refer to the conditional interquartile ranges.

### 3.4. Vertical Velocities and Stratiform Precipitation

In contrast to the impact of vertical motion on the ZM scheme, stratiform precipitation rates are more intuitively connected to vertical velocities. Rauscher *et al.* (2016) proposed that total precipitation rates in models ( $R$ , sum of convective and stratiform precipitation rates) are determined by the upward moisture flux through cloud base,

$$R \approx -\frac{1}{g\rho_w}\omega^+q^+ \quad (5)$$

where  $\omega^+$  and  $q^+$  are the upward vertical pressure velocity and specific humidity at cloud base, respectively, and  $\rho_w$  is the density of rainwater.

Through approximating the cloud base as the 850 hPa level, equation 5 was found to be a good approximation to total precipitation rates in a regional model (Rauscher *et al.* 2016) and in the CAM-SE AGCM with CAM5 physics (O'Brien *et al.* 2016). Figure 9 shows the median convective, stratiform and total precipitation rates conditioned on the 850 hPa moisture flux using 1 mm/day bins, and in the CAM6 *ne30* simulation. The figure shows that the moisture flux is a good approximation to the median total precipitation rate, and that this increase in precipitation rates with moisture flux is primarily due to stratiform precipitation rates.

The increase in stratiform precipitation rates with resolution is likely a due to the increase in magnitude of  $\omega$  with resolution. To unravel the contributions of  $\omega$  and  $q$  at the 850 hPa level,  $\omega_{850}$  and  $q_{850}$  to the increase in climatological stratiform precipitation with resolution, the area averaged, climatological stratiform precipitation rate  $\overline{R}_s$  is decomposed into a time mean magnitude term  $M$  and space-time frequency term  $f$ ,

$$\overline{R}_s = \sum_i \sum_j f(\omega_i, q_j) M(\omega_i, q_j) \quad (6)$$

after Terai *et al.* (2018), and the subscript 850 is dropped from  $\omega$  and  $q$  for brevity.

Table 2 indicates that the  $\pm 15^\circ$  latitude band accounts for most of the change in global mean stratiform precipitation


with resolution. Figure 10 is a plot of the terms  $M(\omega_i, q_j)$  and  $f(\omega_i, q_j) \times M(\omega_i, q_j)$  for all resolutions, and in the  $\pm 15^\circ$  latitude region. The plots are computed using 6-hourly instantaneous output of  $\omega_i$  and  $q_j$ , with 0.05 Pa/s and 0.4 g/kg bins. Values in  $(\omega_i, q_j)$  space are only shown for bins with  $f \geq 1 \times 10^{-5}$ , a reasonable cut-off to a bins' contribution to  $R_s$ . Larger  $\omega$  values correspond to larger magnitude stratiform precipitation rates, which have larger space-time frequencies  $f(\omega_i, q_j)$  at higher resolutions, explaining the increase in  $R_s$  with resolution in Table 2. Changes in the space-time frequency of the  $q$  field contributes comparatively less to changes in stratiform precipitation.

## Acknowledgement

Funding support for this work was in part provided by the U.S. Department of Energy Office of Science (DE-SC0019459) and the National Science Foundation (AGS1648629).

## References

- Arakawa A, Wu CM. 2013. A unified representation of deep moist convection in numerical modeling of the atmosphere. part i. *Journal of the Atmospheric Sciences* **70**(7): 1977–1992.
- Bogenschutz PA, Gettelman A, Morrison H, Larson VE, Craig C, Schanen DP. 2013. Higher-order turbulence closure and its impact on climate simulations in the community atmosphere model. *Journal of Climate* **26**(23): 9655–9676.
- Cho JY, Zhu Y, Newell RE, Anderson BE, Barrick JD, Gregory GL, Sachse GW, Carroll MA, Albercook GM. 1999. Horizontal wavenumber spectra of winds, temperature, and trace gases during the pacific exploratory missions: 1. climatology. *Journal of Geophysical Research: Atmospheres* **104**(D5): 5697–5716.
- Dai A. 2006. Precipitation characteristics in eighteen coupled climate models. *Journal of Climate* **19**(18): 4605–4630.
- Dennis JM, Edwards J, Evans KJ, Guba O, Lauritzen PH, Mirin AA, St-Cyr A, Taylor MA, Worley PH. 2012. CAM-SE: A scalable spectral element dynamical core for the Community Atmosphere Model. *Int. J. High. Perform. C.* **26**(1): 74–89, doi:10.1177/1094342011428142, URL <http://hpc.sagepub.com/content/26/1/74.abstract>.
- Donner LJ, O'Brien TA, Rieger D, Vogel B, Cooke WF. 2016. Are atmospheric updrafts a key to unlocking climate forcing and sensitivity? *Atmospheric Chemistry and Physics* **16**(20): 12 983–12 992.
- Germano M. 1992. Turbulence: the filtering approach. *Journal of Fluid Mechanics* **238**: 325–336.
- Gettelman A, Morrison H, Santos S, Bogenschutz P, Caldwell P. 2015. Advanced two-moment bulk microphysics for global models. part ii: Global model solutions and aerosol–cloud interactions. *Journal of Climate* **28**(3): 1288–1307.
- Golaz JC, Larson VE, Cotton WR. 2002. A pdf-based model for boundary layer clouds. part i: Method and model description. *Journal of the Atmospheric Sciences* **59**(24): 3540–3551, doi:10.1175/1520-0469(2002)059(3540:apbmfb)2.0.co;2.
- Harris LM, Lin SJ, Tu C. 2016. High-resolution climate simulations using gfdl hiram with a stretched global grid. *Journal of Climate* **29**(11): 4293–4314, doi:10.1175/jcli-d-15-0389.1.
- Herrington A, Lauritzen P, Taylor MA, Goldhaber S, Eaton BE, Bacmeister J, Reed K, Ullrich P. 2018. Physics-dynamics coupling with element-based high-order galerkin methods: quasi equal-area physics grid. *Mon. Wea. Rev.* **47**: 69–84, doi:10.1175/MWR-D-18-0136.1.
- Herrington A, Reed K. 2018. An idealized test of the response of the community atmosphere model to near-grid-scale forcing across hydrostatic resolutions. *J. Adv. Model. Earth Syst.* **10**(2): 560–575.
- Herrington AR, Lauritzen PH, Reed KA, Goldhaber S, Eaton BE. 2019. Exploring a lower resolution physics grid in cam-se-cslam. *Journal of Advances in Modeling Earth Systems* **11**.
- Herrington AR, Reed KA. 2017. An explanation for the sensitivity of the mean state of the community atmosphere model to horizontal resolution on aquaplanets. *J. Climate* **30**(13): 4781–4797, doi:10.1175/jcli-d-16-0069.1, URL <http://dx.doi.org/10.1175/jcli-d-16-0069.1>.
- Jeevanjee N. 2017. Vertical velocity in the gray zone. *Journal of Advances in Modeling Earth Systems* **9**(6): 2304–2316, doi:10.1002/2017MS001059.
- Jeevanjee N, Romps DM. 2016. Effective buoyancy at the surface and aloft. *Quart. J. Roy. Meteor. Soc.* **142**(695): 811–820.



**Figure 11.** Decomposition of the climatological stratiform precipitation rates, averaged over the  $\pm 15^\circ$  latitude band into  $\omega_{850}$  and  $q_{850}$  environmental conditions. Left column shows the time mean magnitude term  $M(\omega_i, q_j)$  and the right column is the magnitude term multiplied by the space-time frequency term  $f(\omega_i, q_j) \times M(\omega_i, q_j)$ . Integrals over  $f \times M$  gives the climatological, area averaged stratiform precipitation rate. Panel labels denote the grid resolution of the model run.



- Kiehl J, Williamson D. 1991. Dependence of cloud amount on horizontal resolution in the national center for atmospheric research community climate model. *Journal of Geophysical Research: Atmospheres* **96**(D6): 10955–10980.
- Lauritzen PH, Nair R, Herrington A, Callaghan P, Goldhaber S, Dennis J, Bacmeister JT, Eaton B, Zarzycki C, Taylor MA, Gettelman A, Neale R, Dobbins B, Reed K, Dubos T. 2018. NCAR CESM2.0 release of CAM-SE: A reformulation of the spectral-element dynamical core in dry-mass vertical coordinates with comprehensive treatment of condensates and energy. *J. Adv. Model. Earth Syst.* doi:10.1029/2017MS001257.
- Lauritzen PH, Taylor MA, Overfelt J, Ullrich PA, Nair RD, Goldhaber S, Kelly R. 2017. CAM-SE-CSLAM: Consistent coupling of a conservative semi-lagrangian finite-volume method with spectral element dynamics. *Mon. Wea. Rev.* **145**(3): 833–855, doi:10.1175/MWR-D-16-0258.1.
- Lauritzen PH, Williamson DL. 2019. A total energy error analysis of dynamical cores and physics-dynamics coupling in the community atmosphere model (cam). *J. Adv. Model. Earth Syst.* doi:10.1029/2018MS001549.
- Liu X, Easter RC, Ghan SJ, Zaveri R, Rasch P, Shi X, Lamarque JF, Gettelman A, Morrison H, Vitt F, *et al.* 2012. Toward a minimal representation of aerosols in climate models: Description and evaluation in the community atmosphere model cam5. *Geoscientific Model Development* **5**(3): 709.
- Medeiros B, Williamson DL, Olson JG. 2016. Reference aquaplanet climate in the community atmosphere model, version 5. *J. Adv. Model. Earth Syst.* **8**(1): 406–424, doi:10.1002/2015MS000593.
- Nastrom GD, Gage KS. 1985. A climatology of atmospheric wavenumber spectra of wind and temperature observed by commercial aircraft. *J. Atmos. Sci.* **42**: 950–960.
- Neale RB, Chen CC, Gettelman A, Lauritzen PH, Park S, Williamson DL, Conley AJ, Garcia R, Kinnison D, Lamarque JF, Marsh D, Mills M, Smith AK, Tilmes S, Vitt F, Cameron-Smith P, Collins WD, Iacono MJ, Easter RC, Ghan SJ, Liu X, Rasch PJ, Taylor MA. 2012. Description of the NCAR Community Atmosphere Model (CAM 5.0). NCAR Technical Note NCAR/TN-486+STR, National Center of Atmospheric Research.
- Neale RB, Chen CC, Gettelman A, Lauritzen PH, Park S, Williamson DL, Conley AJ, Garcia R, Kinnison D, Lamarque JF, Marsh D, Mills M, Smith AK, Tilmes S, Vitt F, Cameron-Smith P, Collins WD, Iacono MJ, Easter RC, Ghan SJ, Liu X, Rasch PJ, Taylor MA. 2010. Description of the NCAR Community Atmosphere Model (CAM 4.0). NCAR Technical Note, National Center of Atmospheric Research.
- Neale RB, Hoskins BJ. 2000. A standard test for agcms including their physical parametrizations: I: the proposal. *Atmos. Sci. Lett* **1**(2): 101–107, doi:10.1006/asle.2000.0022.
- Neale RB, Richter JH, Jochum M. 2008. The impact of convection on ENSO: From a delayed oscillator to a series of events. *J. Climate* **21**: 5904–5924.
- O'Brien TA, Collins WD, Kashinath K, Rübel O, Byna S, Gu J, Krishnan H, Ullrich PA. 2016. Resolution dependence of precipitation statistical fidelity in hindcast simulations. *J. Adv. Model. Earth Syst.* **8**(2): 976–990, doi:10.1002/2016ms000671, URL <http://dx.doi.org/10.1002/2016ms000671>.
- Orlanski I. 1981. The quasi-hydrostatic approximation. *J. Atmos. Sci.* **38**: 572–582, doi:10.1175/1520-0469(1981)038<0572:TQHA>2.0.CO;2, URL [http://dx.doi.org/10.1175/1520-0469\(1981\)038<0572:TQHA>2.0.CO;2](http://dx.doi.org/10.1175/1520-0469(1981)038<0572:TQHA>2.0.CO;2).
- Pauluis O, Garner S. 2006. Sensitivity of radiative–convective equilibrium simulations to horizontal resolution. *J. Atmos. Sci.* **63**(7): 1910–1923.
- Plant RS, Craig GC. 2008. A stochastic parameterization for deep convection based on equilibrium statistics. *J. Atmos. Sci.* **65**: 87–105, doi:10.1175/2007JAS2263.1, URL <http://dx.doi.org/10.1175/2007JAS2263.1>.
- Pope V, Stratton R. 2002. The processes governing horizontal resolution sensitivity in a climate model. *Climate Dynamics* **19**(3–4): 211–236.
- Rauscher SA, O'Brien TA, Piani C, Coppola E, Giorgi F, Collins WD, Lawston PM. 2016. A multimodel intercomparison of resolution effects on precipitation: simulations and theory. *Climate Dynamics* **47**(7–8): 2205–2218, doi:10.1007/s00382-015-2959-5.
- Rauscher SA, Ringler TD, Skamarock WC, Mirin AA. 2013. Exploring a global multiresolution modeling approach using aquaplanet simulations. *Journal of Climate* **26**(8): 2432–2452, doi:10.1175/jcli-d-12-00154.1.
- Raymond DJ, Blyth AM. 1992. Extension of the stochastic mixing model to cumulonimbus clouds. *Journal of the Atmospheric Sciences* **49**(21): 1968–1983, doi:10.1175/1520-0469(1992)049<1968:eotmmm>2.0.co;2.
- Richter JH, Rasch PJ. 2008. Effects of convective momentum transport on the atmospheric circulation in the community atmosphere model, version 3. *J. Climate* **21**(7): 1487–1499.
- Satoh M, Matsuno T, Tomita H, Miura H, Nasuno T, Iga S. 2008. Nonhydrostatic icosahedral atmospheric model (NICAM) for global cloud resolving simulations. *J. Comput. Phys.* : 3486–3514.
- Skamarock WC, Klemp JB, Duda MG, Fowler L, Park SH, Ringler TD. 2012. A multi-scale nonhydrostatic atmospheric model using centroidal Voronoi tessellations and C-grid staggering. *Mon. Wea. Rev.* **240**: 3090–3105, doi:10.1175/MWR-D-11-00215.1.
- Song F, Zhang GJ. 2018. Understanding and improving the scale dependence of trigger functions for convective parameterization using cloud-resolving model data. *Journal of Climate* **31**(18): 7385–7399.
- Taylor MA, Fournier A. 2010. A compatible and conservative spectral element method on unstructured grids. *J. Comput. Phys.* **229**(17): 5879 – 5895, doi:10.1016/j.jcp.2010.04.008.
- Terai CR, Caldwell PM, Klein SA, Tang Q, Branstetter ML. 2018. The atmospheric hydrologic cycle in the acme v0. 3 model. *Climate dynamics* **50**(9–10): 3251–3279.
- Ullrich PA, Jablonowski C, Kent J, Lauritzen PH, Nair R, Reed KA, Zarzycki CM, Hall DM, Dazlich D, Heikes R, Konor C, Randall D, Dubos T, Meurdesoif Y, Chen X, Harris L, Kühnlein C, Lee V, Qaddouri A, Girard C, Giorgetta M, Reinert D, Klemp J, Park SH, Skamarock W, Miura H, Ohno T, Yoshida R, Walko R, Reinecke A, Viner K. 2017. "dcmp2016: A review of non-hydrostatic dynamical core design and intercomparison of participating models". *Geosci. Model Dev.* **10**: 4477–4509, doi:10.5194/gmd-10-4477-2017.
- Weisman ML, Skamarock WC, Klemp JB. 1997. The resolution dependence of explicitly modeled convective systems. *Monthly Weather Review* **125**(4): 527–548, doi:10.1175/1520-0493(1997)125<0527:TRDOEM>2.0.CO;2.
- Wilks DS. 2011. *Statistical methods in the atmospheric sciences*, vol. 100. Academic press.
- Williamson DL. 2008. Convergence of aqua-planet simulations with increasing resolution in the community atmospheric model, version 3. *Tellus A* **60**(5): 848–862, doi:10.1111/j.1600-0870.2008.00339.x.
- Williamson DL. 2013. The effect of time steps and time-scales on parametrization suites. *Quart. J. Roy. Meteor. Soc.* **139**(671): 548–560, doi:10.1002/qj.1992.
- Williamson DL, Kiehl JT, Hack JJ. 1995. Climate sensitivity of the near community climate model (ccm2) to horizontal resolution. *Climate Dynamics* **11**(7): 377–397, doi:10.1007/s003820050082.
- Yang Q, Leung LR, Rauscher SA, Ringler TD, Taylor MA. 2014. Atmospheric moisture budget and spatial resolution dependence of precipitation extremes in aquaplanet simulations. *Journal of Climate* **27**(10): 3565–3581, doi:10.1175/jcli-d-13-00468.1.
- Zarzycki CM, Levy MN, Jablonowski C, Overfelt JR, Taylor MA, Ullrich PA. 2014. Aquaplanet experiments using cam's variable-resolution dynamical core. *J. Climate* **27**(14): 5481–5503, doi:10.1175/JCLI-D-14-00004.1.
- Zhang G, McFarlane N. 1995. Sensitivity of climate simulations to the parameterization of cumulus convection in the canadian climate centre general circulation model. *Atmosphere-ocean* **33**(3): 407–446.
- Zhang GJ. 2002. Convective quasi-equilibrium in midlatitude continental environment and its effect on convective parameterization. *Journal of Geophysical Research: Atmospheres* **107**(D14): ACL–12.
- Zängl G, Reinert D, Rípodas P, Baldauf M. 2014. The icon (icosahedral non-hydrostatic) modelling framework of dwd and mpi-m: Description of the non-hydrostatic dynamical core. *Quarterly Journal of the Royal Meteorological Society* **141**(687): 563–579, doi:10.1002/qj.2378.