

# ALGEBRAIC DEPENDENCE AND MILNOR K-THEORY

ADAM TOPAZ

ABSTRACT. This paper shows that algebraic (in)dependence is encoded in Milnor K-theory of fields. As an application, we show that the isomorphism type of a field is determined by its Milnor K-theory, up to purely inseparable extensions, in most situations.

## CONTENTS

1. Introduction	1
2. Notation and preliminaries	2
3. The local theory	5
4. Algebraic dependence	9
5. Main results	15
References	22

## 1. INTRODUCTION

Let  $K$  be a field. The Milnor K-theory of  $K$  has a very simple definition:

$$K_*^M(K) := \frac{T_*(K^\times)}{\langle x \otimes y \mid x + y = 1 \rangle}$$

where  $T_*(K^\times)$  denotes the tensor algebra of the  $\mathbb{Z}$ -module  $K^\times$ , and the two-sided ideal  $\langle x \otimes y \mid x + y = 1 \rangle$  consists of the so-called *Steinberg relations*.

In degree one, we have the multiplicative group,  $K_1^M(K) = K^\times$ , while the ring structure of  $K_*^M(K)$  involves the additive structure of  $K$  as well. It is natural to ask whether the *field*  $K$  itself is determined (up-to isomorphism) by  $K_*^M(K)$ . This question was considered in [2, 4], focusing mostly on finitely-generated field extensions and eventually relying on the so-called *fundamental theorem of projective geometry* to reconstruct the fields in question.

In this paper, we investigate this question for fields which do not necessarily satisfy any finiteness conditions, and we obtain the following main result.

**Main Theorem.** *Let  $K$  be any field whose absolute transcendence degree is at least 5. Then the isomorphism type of  $K$  is determined, up to purely inseparable extensions, by the  $\mathbb{Q}$ -algebra  $\mathbb{Q} \otimes_{\mathbb{Z}} K_*^M(K)$ .*

---

*Date:* Last updated: January 18, 2023.

*2020 Mathematics Subject Classification.* 12G99, 12J10, 19D45.

*Key words and phrases.* Algebraic dependence, Milnor K-theory, valuation theory, anabelian geometry.

Research supported by the author's NSERC discovery grant.

See Theorem 5.4 for the precise statement. We also prove a similar *relative* result for relatively algebraically closed field extensions of sufficiently large transcendence degree in Theorem 5.3. Note that the theorem above imposes no additional restrictions on  $K$  besides the bound on the absolute transcendence degree. For example, this theorem applies to any sufficiently large *algebraically closed* field.

Since we wish to work with fields whose multiplicative group may even be *divisible*, it is important to work with  $\mathbb{Q} \otimes_{\mathbb{Z}} K_*^M(K)$  as opposed to  $K_*^M(K)$  itself. More precisely, if  $K$  is radically closed, then the quotient  $K^\times/\text{torsion}$  is already a  $\mathbb{Q}$ -vector space, and thus  $\mathbb{Q}^\times$  provides a source of indeterminacy for  $K_*^M(K)/\langle \text{torsion} \rangle$ . Namely, any  $r \in \mathbb{Q}^\times$  yields an automorphism of  $K_*^M(K)/\langle \text{torsion} \rangle$  defined by  $t \mapsto t^r$  in degree one, and such automorphisms only arise from field theory if  $r$  is a power of the characteristic of  $K$ . By tensoring  $K_*^M(K)$  with  $\mathbb{Q}$ , we can control for such indeterminacies in our main result.

Furthermore, if  $K$  is any field and  $K \rightarrow K^i$  denotes the perfection of  $K$ , then the corresponding map

$$K_*^M(K) \rightarrow K_*^M(K^i)$$

induces an *isomorphism* after tensoring with  $\mathbb{Q}$ . Thus, inseparability is an additional source of indeterminacy which must be accounted for, hence we can only expect to recover the isomorphism type of  $K$  up to purely inseparable extensions when working with  $\mathbb{Q} \otimes_{\mathbb{Z}} K_*^M(K)$ .

The technical core of this work is in recovering all the information about *algebraic dependence* from Milnor K-theory, see Theorem 4.6. Once we obtain all information about algebraic dependence, our reconstruction results will follow by applying a distant cousin of the fundamental theorem of projective geometry, due to Evans-Hrushovski [8, 9] and Gismatullin [10], based on the *group-configuration theorem*. In the case where the fields (or field extensions) in question are finitely-generated, we can instead use one of the main theorems from [4] to obtain better reconstruction results. For example, Theorem 5.7 (which uses [4, Theorem 4] in an essential way) shows that the isomorphism type of a finitely-generated field  $K$  of absolute transcendence degree  $\geq 2$  is determined by  $K_*^M(K)$  with no need to pass to inseparable extensions.

## 2. NOTATION AND PRELIMINARIES

We will primarily work with a fixed field denoted by  $K$ . In some cases we will also consider subfields of  $K$ , usually denoted  $k$ .

**2.1. Quotients of Milnor K-theory.** For a subgroup  $T$  of  $K^\times$ , we write

$$K_*^M(K|T) := \frac{K_*^M(K)}{\langle T \rangle}$$

where  $\langle T \rangle$  refers to the (two-sided) ideal of  $K_*^M(K)$  generated by  $T \subset K^\times = K_1^M(K)$ . If  $T$  is trivial, then we omit it from the notation to match the standard notation for Milnor K-theory:  $K_*^M(K) := K_*^M(K|\{1\})$ . In the case where  $T = k^\times$  for a subfield  $k$  of  $K$ , we write  $K_*^M(K|k)$  instead of  $K_*^M(K|k^\times)$ .

As usual, we will use the notation  $\{f_1, \dots, f_n\}$  to denote the product of  $f_1, \dots, f_n \in K^\times = K_1^M(K)$  in  $K_n^M(K)$ , and such elements of  $K_n^M(K)$  will be called *symbols*. A similar notation and terminology will also be used for the variants of  $K_*^M(K)$  we consider in this paper, while ensuring that the variant being considered is clear from context.

**2.2. Duality.** For a subgroup  $T$  of  $K^\times$ , we write

$$\mathcal{K}_{K|T} := \mathbb{Q} \otimes_{\mathbb{Z}} (K^\times/T).$$

If  $T = \{1\}$ , then we write  $\mathcal{K}_K$  instead of  $\mathcal{K}_{K|T}$ , and if  $T = k^\times$  for a subfield  $k$  of  $K$ , we will write  $\mathcal{K}_{K|k}$  instead of  $\mathcal{K}_{K|k^\times}$ . The operation of  $\mathcal{K}_{K|T}$  will always be written *additively*.

We will consider the dual

$$\mathcal{G}_K := \text{Hom}_{\mathbb{Z}}(K^\times, \mathbb{Q}) = \text{Hom}_{\mathbb{Q}}(\mathcal{K}_K, \mathbb{Q})$$

as a topological vector space with respect to the weak topology, where  $\mathbb{Q}$  is given the discrete topology. We have an obvious perfect pairing

$$K^\times \times \mathcal{G}_K \rightarrow \mathbb{Q}.$$

For a subspace  $\mathcal{H}$  of  $\mathcal{G}_K$ , we write  $\mathcal{H}^\perp \subset K^\times$  for the orthogonal of  $\mathcal{H}$  with respect to this pairing. For a subgroup  $T$  of  $K^\times$ , we will use the notation  $\mathcal{G}_{K|T} \subset \mathcal{G}_K$  for the orthogonal of  $T$  with respect to this pairing. As always, if  $T = k^\times$  for a subfield  $k$  of  $K$ , we write  $\mathcal{G}_{K|k}$  instead of  $\mathcal{G}_{K|k^\times}$ . When  $T = \{1\}$  is trivial, one has  $\mathcal{G}_{K|T} = \mathcal{G}_K$ , so our convention of omitting  $T$  from the notation in this case still works.

A subgroup  $T$  of an abelian group  $A$  will be called *saturated* if  $A/T$  is torsion-free. For any subgroup  $T$  of  $K^\times$ , the subspace  $\mathcal{G}_{K|T}$  is closed, and for a subspace  $\mathcal{H} \subset \mathcal{G}_K$ , the subgroup  $\mathcal{H}^\perp$  is saturated. In fact, if  $T$  is any subgroup of  $K^\times$ , then  $\mathcal{G}_{K|T}^\perp$  is the saturation of  $T$  (i.e. the smallest saturated subgroup containing  $T$ ) and if  $\mathcal{H} \subset \mathcal{G}_K$  is any subspace then  $\mathcal{G}_{K|\mathcal{H}^\perp}$  is the closure of  $\mathcal{H}$ .

The maps  $\mathcal{H} \mapsto \mathcal{H}^\perp$  and  $T \mapsto \mathcal{G}_{K|T}$  provide a one-to-one order-reversing correspondence between the closed subspaces of  $\mathcal{G}_K$  and the saturated subgroups of  $K^\times$ . We also have canonical perfect pairing

$$\mathcal{K}_{K|T} \times \mathcal{G}_{K|T} \rightarrow \mathbb{Q}$$

associated to any subgroup  $T$  of  $K^\times$ . We will say so explicitly when considering orthogonals with respect to this pairings to avoid any potential confusion with the notation  $(-)^{\perp}$  introduced above.

The base-change  $\mathbb{Q} \otimes_{\mathbb{Z}} K_*^{\text{M}}(K|T)$  will be denoted by  $\mathcal{K}_*(K|T)$ . As usual, if  $T = \{1\}$  then we write  $\mathcal{K}_*(K)$  instead of  $\mathcal{K}_*(K|\{1\})$  and if  $T = k^\times$  for a subfield  $k$  of  $K$ , then we write  $\mathcal{K}_*(K|k)$  instead of  $\mathcal{K}_*(K|k^\times)$ . Note that for any subgroup  $T$  of  $K^\times$ , one has  $\mathcal{K}_1(K|T) = \mathcal{K}_{K|T}$ .

**2.3. Alternating pairs.** Elements of  $\mathcal{G}_K$  will be considered both as  $\mathbb{Z}$ -linear maps  $K^\times \rightarrow \mathbb{Q}$  and as  $\mathbb{Q}$ -linear maps  $\mathcal{K}_K \rightarrow \mathbb{Q}$ . If  $f \in \mathcal{G}_K$  with  $T = \ker(f) \subset K^\times$ , then we may also consider  $f$  as a  $\mathbb{Z}$ -linear map  $K^\times/T \rightarrow \mathbb{Q}$  and as a  $\mathbb{Q}$ -linear map  $\mathcal{K}_{K|T} \rightarrow \mathbb{Q}$ .

A pair of elements  $f, g \in \mathcal{G}_K$  will be called an *alternating pair* provided that

$$f(x) \cdot g(y) = f(y) \cdot g(x)$$

whenever  $x, y \in K^\times$  satisfy  $x + y = 1$  in  $K$ . We denote the associated binary relation on  $\mathcal{G}_K$  by  $\mathcal{R}_K$ :

$$\mathcal{R}_K(f, g) \iff f, g \text{ are an alternating pair.}$$

In fact, for the majority of this paper we will be working with the structure consisting of the following data (associated to various subgroups  $T$  of  $K^\times$ ), which we abbreviate as  $\mathcal{A}(K|T)$ :

- (1) The  $\mathbb{Q}$ -vector space  $\mathcal{K}_{K|T}$ .
- (2) The topological  $\mathbb{Q}$ -vector space  $\mathcal{G}_{K|T}$ .

- (3) The canonical pairing  $\mathcal{K}_{K|T} \times \mathcal{G}_{K|T} \rightarrow \mathbb{Q}$ .
- (4) The restriction of the relation  $\mathcal{R}_K$  to  $\mathcal{G}_{K|T}$ .

As before, we write  $\mathcal{A}(K)$  instead of  $\mathcal{A}(K|T)$  if  $T$  is trivial and  $\mathcal{A}(K|k)$  instead of  $\mathcal{A}(K|k^\times)$  when  $k$  is a subfield of  $K$ .

Recall that the Steinberg relations in Milnor K-theory are generated by basic tensors of the form  $x \otimes y$  for  $x, y \in K^\times$  satisfying  $x + y = 1$ . Thus, the alternating condition for pairs of elements of  $\mathcal{G}_K$  can be tested using the product in Milnor K-theory, as the following fact summarizes.

**Fact 2.1.** *Let  $f, g \in \mathcal{G}_K$  be given and let  $T \subset (\mathbb{Q} \cdot f + \mathbb{Q} \cdot g)^\perp$  be any subgroup. The following are equivalent:*

- (1) *For all  $x, y \in K^\times/T = K_1^M(K|T)$  satisfying  $\{x, y\} = 0$  in  $K_2^M(K|T)$ , one has*

$$f(x) \cdot g(y) = f(y) \cdot g(x).$$

- (2) *For all  $x, y \in \mathcal{K}_{K|T} = \mathcal{K}_1(K|T)$  satisfying  $\{x, y\} = 0$  in  $\mathcal{K}_2(K|T)$ , one has*

$$f(x) \cdot g(y) = f(y) \cdot g(x).$$

- (3)  *$f, g$  are an alternating pair.*

In particular, this shows that the data  $\mathcal{A}(K|T)$  is *completely determined* (in a functorial manner) by the algebra  $\mathcal{K}_*(K|T)$ . Indeed,  $\mathcal{G}_{K|T}$  is the (weak) dual of  $\mathcal{K}_{K|T} = \mathcal{K}_1(K|T)$ , and the fact above shows that for  $f, g \in \mathcal{G}_{K|T}$  one has  $\mathcal{R}_K(f, g)$  if and only if for all  $x, y \in \mathcal{K}_1(K|T)$  such that  $\{x, y\} = 0$  in  $\mathcal{K}_2(K|T)$ , one has  $f(x) \cdot g(y) = f(y) \cdot g(x)$ .

We will borrow some notation and terminology from group theory by considering  $\mathcal{R}_K(-, -)$  as being analogous to the condition that two elements of a group commute. Namely, for a closed subspace  $\mathcal{H}$  of  $\mathcal{G}_K$  we consider the following (closed) subspaces of  $\mathcal{G}_K$ :

- (1)  $\mathcal{C}_K(\mathcal{H}) := \{f \in \mathcal{G}_K \mid \forall g \in \mathcal{H}, \mathcal{R}_K(f, g)\}$ , the  $\mathcal{R}_K$ -centralizer of  $\mathcal{H}$ .
- (2)  $\mathcal{Z}_K(\mathcal{H}) := \{f \in \mathcal{H} \mid \forall g \in \mathcal{H}, \mathcal{R}_K(f, g)\}$ , the  $\mathcal{R}_K$ -centre of  $\mathcal{H}$ .

**2.4. Valuations.** Valuations on  $K$  will only be considered up-to equivalence. Let  $v, w$  be two valuations. Our convention is that  $v \leq w$  means  $v$  is a *coarsening* of  $w$ .

Let  $v$  be a valuation of  $K$ . We shall write  $\mathcal{O}_v$  for the valuation ring of  $v$ ,  $\mathfrak{m}_v$  for the maximal ideal of  $\mathcal{O}_v$ ,  $Kv$  for the residue field of  $v$  and  $vK$  for the value group of  $v$ . The unit group  $\mathcal{O}_v^\times$  will be denoted by  $U_v$  and the principal unit group  $1 + \mathfrak{m}_v$  will be denoted by  $U_v^1$ . If  $k$  is a subfield of  $K$ , then we write  $kv$  and  $vk$  for the residue field and value group of the restriction of  $v$  to  $k$ .

We define:

$$\mathcal{I}_v := \mathcal{G}_{K|U_v}, \quad \mathcal{D}_v := \mathcal{G}_{K|U_v^1}.$$

Note that  $\mathcal{I}_v \subset \mathcal{D}_v$  and that the exact sequence

$$1 \rightarrow Kv^\times \rightarrow K^\times/U_v^1 \rightarrow vK \rightarrow 1$$

dualizes to an exact sequence

$$0 \rightarrow \mathcal{I}_v \rightarrow \mathcal{D}_v \rightarrow \mathcal{G}_{Kv} \rightarrow 0.$$

For a subgroup  $T$  of  $K^\times$ , write  $Tv$  for the image of  $T \cap U_v$  in  $Kv^\times$  and  $vT$  for the image of  $T$  in  $vK$ . We have an exact sequence

$$1 \rightarrow Kv^\times/Tv \rightarrow K^\times/(T \cdot U_v^1) \rightarrow vK/vT \rightarrow 1$$

which dualizes to an exact sequence of the form

$$0 \rightarrow \mathcal{G}_{K|T} \cap \mathcal{I}_v \rightarrow \mathcal{G}_{K|T} \cap \mathcal{D}_v \rightarrow \mathcal{G}_{Kv|Tv} \rightarrow 0.$$

In the special case where  $T = k^\times$  for a subfield  $k$  of  $K$ , our notational conventions are compatible. Namely, the natural exact sequence

$$1 \rightarrow Kv^\times / kv^\times \rightarrow K^\times / (k^\times \cdot U_v^1) \rightarrow vK / vk \rightarrow 1$$

dualizes to an exact sequence

$$0 \rightarrow \mathcal{G}_{K|k} \cap \mathcal{I}_v \rightarrow \mathcal{G}_{K|k} \cap \mathcal{D}_v \rightarrow \mathcal{G}_{Kv|kv} \rightarrow 0.$$

The subspace of  $\mathcal{K}_K$  generated by the image of  $U_v^1$  will be denoted by  $\mathcal{U}_v^1$  and the subspace generated by the image of  $U_v$  will be denoted by  $\mathcal{U}_v$ . Similarly, if  $T$  is a subgroup of  $K^\times$ , we write  $\mathcal{U}_{v|T}^1$  for the image of  $\mathcal{U}_v^1$  in  $\mathcal{K}_{K|T}$  and  $\mathcal{U}_{v|T}$  for the image of  $\mathcal{U}_v$ . As always, when  $T = k^\times$ , we write  $\mathcal{U}_{v|k}$  and  $\mathcal{U}_{v|k}^1$  instead of  $\mathcal{U}_{v|k^\times}$  and  $\mathcal{U}_{v|k^\times}^1$ .

Note that  $\mathcal{U}_v$  resp.  $\mathcal{U}_v^1$  is the orthogonal of  $\mathcal{I}_v$  resp.  $\mathcal{D}_v$  with respect to the pairing  $\mathcal{K}_K \times \mathcal{G}_K \rightarrow \mathbb{Q}$ . Similarly,  $\mathcal{U}_{v|T}$  resp.  $\mathcal{U}_{v|T}^1$  is the orthogonal of  $\mathcal{G}_{K|T} \cap \mathcal{I}_v$  resp.  $\mathcal{G}_{K|T} \cap \mathcal{D}_v$  with respect to the pairing  $\mathcal{K}_{K|T} \times \mathcal{G}_{K|T} \rightarrow \mathbb{Q}$ .

### 3. THE LOCAL THEORY

Our starting point is the following fundamental result.

**Theorem 3.1.** *Let  $K$  be any field, and let  $\mathcal{D}$  be a closed subspace of  $\mathcal{G}_K$ . The following are equivalent:*

- (1) *For all  $f, g \in \mathcal{D}$ , one has  $\mathcal{R}_K(f, g)$ .*
- (2) *There exists a valuation  $v$  of  $K$  and a closed subspace  $\mathcal{I} \subset \mathcal{D}$  of codimension  $\leq 1$ , such that  $\mathcal{D} \subset \mathcal{D}_v$  and  $\mathcal{I} \subset \mathcal{I}_v$ .*

Variants of this theorem have appeared in the works of Bogomolov [3], Bogomolov-Tschinkel [1], Efrat [6], Koenigsmann [11], Engler-Koenigsmann [7], the author [13, 14], and others, albeit primarily the Galois-theoretic context. The proof of Theorem 3.1 has now been completely formally verified using the **Lean3** interactive theorem prover [5] and its formally verified mathematics library **mathlib** [12], see [15]. We thus omit the proof, referring instead to the references above for the key ideas and to [15] for the computer-verified proof.

The power of this theorem is in the implication (1)  $\Rightarrow$  (2), while the converse is a simple consequence of the ultrametric inequality. We will need a slightly stronger variant of the “easy” direction (2)  $\Rightarrow$  (1), formulated as follows.

**Lemma 3.2.** *Suppose that  $v$  is a valuation of  $K$  and  $f, g \in \mathcal{D}_v$  are given. Let  $f_v$  and  $g_v$  denote the images of  $f$  and  $g$  in  $\mathcal{G}_{Kv}$  under the canonical map  $\mathcal{D}_v \rightarrow \mathcal{G}_{Kv}$ . Then  $\mathcal{R}_K(f, g)$  holds if and only if  $\mathcal{R}_{Kv}(f_v, g_v)$  holds.*

*Proof.* Suppose  $\mathcal{R}_K(f, g)$  holds, and let  $x, y \in Kv^\times$  satisfy  $x + y = 1$ . We may choose lifts  $\tilde{x}, \tilde{y} \in U_v$  of  $x, y$  such that  $\tilde{x} + \tilde{y} = 1$ . Thus

$$f_v(x) \cdot g_v(y) = f(\tilde{x}) \cdot g(\tilde{y}) = f(\tilde{y}) \cdot g(\tilde{x}) = f_v(y) \cdot g_v(x).$$

Conversely, suppose that  $\mathcal{R}_{Kv}(f_v, g_v)$  holds and let  $x, y \in K^\times$  be such that  $x + y = 1$ . We must show that

$$f(x) \cdot g(y) = f(y) \cdot g(x).$$

If  $v(x) > 0$  then  $f(y) = g(y) = 0$  since  $f, g \in \mathcal{D}_v$ , so the equation in question trivially holds true. The equation similarly holds true if  $v(y) > 0$ . If  $v(x) < 0$  then  $y = x \cdot (x^{-1} - 1)$  while  $v(x^{-1}) > 0$ . Since  $f(-1) = 0$  and  $f \in \mathcal{D}_v$ , it follows that

$$f(y) = f(x) + f(x^{-1} - 1) = f(x) + f(1 - x^{-1}) = f(x).$$

We similarly have  $g(y) = g(x)$ , so the equation again holds true. The equation similarly holds true if  $v(y) < 0$ .

The last case to consider is where  $v(x) = v(y) = 0$ , in which case  $x, y \in U_v$  and the values of  $f$  and  $g$  at  $x$  and  $y$  can be computed in the residue field. In other words, letting  $\bar{x}$  and  $\bar{y}$  denote the images of  $x$  and  $y$  in  $Kv^\times$ , we have  $\bar{x} + \bar{y} = 1$  so that

$$f(x) \cdot g(y) = f_v(\bar{x}) \cdot g_v(\bar{y}) = f_v(\bar{y}) \cdot g_v(\bar{x}) = f(y) \cdot g(x).$$

In any case, we see that the necessary equation does indeed hold.  $\square$

**3.1. Valuative subspaces.** A closed subspace  $\mathcal{I}$  of  $\mathcal{G}_K$  will be called *valuative* provided that  $\mathcal{I} \subset \mathcal{I}_v$  for some valuation  $v$  of  $K$ .

**Lemma 3.3.** *Suppose that  $\mathcal{I}$  is valuative. Then there exists a unique minimal valuation  $v_{\mathcal{I}}$  such that  $\mathcal{I} \subset \mathcal{I}_{v_{\mathcal{I}}}$ . The valuation  $v = v_{\mathcal{I}}$  is characterized by the following two properties:*

- (1) *One has  $\mathcal{I} \subset \mathcal{I}_v$ , or equivalently  $U_v \subset \mathcal{I}^\perp$ .*
- (2) *The subgroup  $v(\mathcal{I}^\perp)$  contains no nontrivial convex subgroups.*

*Proof.* The collection of all valuations  $w$  such that  $\mathcal{I} \subset \mathcal{I}_w$  is nonempty by assumption. Also, if  $w_i$  is a chain of such valuations, then the infimum  $w$  of the  $w_i$  satisfies

$$U_w = \bigcup_i U_{w_i}$$

and as  $U_{w_i} \subset \mathcal{I}^\perp$  for all  $i$ , it follows that  $U_w \subset \mathcal{I}^\perp$  hence  $w$  is also in the collection. This collection is also closed under binary infimums: If  $w_1$  and  $w_2$  are two such valuations and  $w$  is their infimum, then

$$U_{w_1} \cdot U_{w_2} = U_w,$$

and since  $U_{w_i} \subset \mathcal{I}^\perp$ , we also have  $U_w \subset \mathcal{I}^\perp$ , hence  $w$  is again in the collection. It follows that this collection has a unique minimal element  $v_{\mathcal{I}}$ .

Put  $v = v_{\mathcal{I}}$ . Note that  $v(\mathcal{I}^\perp)$  contains no nontrivial convex subgroups for otherwise the coarsening associated to such a subgroup would contradict the minimality of  $v$ . Conversely, if  $v$  satisfies (1) and (2) and  $w$  is a coarsening of  $v$  satisfying (1), then the convex subgroup of  $vK$  associated to  $w$  must be contained in  $v(\mathcal{I}^\perp)$ . This subgroup must be trivial by condition (2) and thus  $w = v$ . It follows that  $v$  is minimal with respect to condition (1), hence  $v = v_{\mathcal{I}}$ . This concludes the proof.  $\square$

**Lemma 3.4.** *Suppose that  $v$  is a valuation of  $K$  and that  $\mathcal{H}$  is a closed subspace of  $\mathcal{G}_K$ . Then  $v((\mathcal{I}_v \cap \mathcal{H})^\perp)$  is the saturation of  $v(\mathcal{H}^\perp)$  in  $vK$ . In particular, if  $\mathcal{H} \subset \mathcal{I}_v$  then  $v(\mathcal{H}^\perp)$  is saturated.*

*Proof.* Put  $H := (\mathcal{I}_v \cap \mathcal{H})^\perp$ . First, note that  $v(H)$  is indeed saturated in  $vK$ . Indeed, if  $n \cdot v(t) \in v(H)$  for some  $t \in K^\times$  and some positive integer  $n$ , then  $t^n \in U_v \cdot H$  while  $U_v \subset H$  so that  $t^n \in H$ . Since  $H$  is itself saturated, it follows that  $t \in H$  hence  $v(t) \in v(H)$ .

Put  $T := U_v \cdot \mathcal{H}^\perp$ . Since  $\mathcal{I}_v^\perp = U_v$ , it follows that  $\mathcal{G}_{K|T} = \mathcal{I}_v \cap \mathcal{H}$  and thus  $H$  is the saturation of  $T$ . This means that  $H/T$  is torsion, hence  $v(H)/v(T)$  is torsion as well, while  $v(T) = v(\mathcal{H}^\perp)$ . It follows that  $v(H)$  is indeed the saturation of  $v(\mathcal{H}^\perp)$ .  $\square$

**Lemma 3.5.** *Let  $\mathcal{H}$  be a closed subspace of  $\mathcal{G}_K$ , and let  $v$  be a valuation of  $K$ . Then  $v = v_{\mathcal{I}}$  for  $\mathcal{I} = \mathcal{I}_v \cap \mathcal{H}$  if and only if the saturation of  $v(\mathcal{H}^\perp)$  in  $vK$  contains no nontrivial convex subgroups.*

*Proof.* Combine Lemmas 3.3 and 3.4.  $\square$

### 3.2. Detecting valuative subspaces.

**Lemma 3.6.** *Let  $\mathcal{I}$  be a valuative subspace of  $\mathcal{G}_K$  with associated valuation  $v := v_{\mathcal{I}}$ . Then one has  $\mathcal{D}_v = \mathcal{C}_K(\mathcal{I})$ .*

*Proof.* The inclusion  $\mathcal{D}_v \subset \mathcal{C}_K(\mathcal{I})$  follows from Lemma 3.2. Conversely, suppose that  $f \in \mathcal{G}_K$  satisfies  $\mathcal{R}_K(f, g)$  for all  $g \in \mathcal{I}$ . Let  $x \in K^\times$  be an element satisfying  $v(x) > 0$ .

Note that for all  $g \in \mathcal{I}$ , one has  $g(1 - x) = 0$  since  $1 - x \in U_v \subset \mathcal{I}^\perp$ . If there exists some  $g \in \mathcal{I}$  such that  $g(x) \neq 0$ , then one has  $f(1 - x) = 0$  since

$$f(1 - x) \cdot g(x) = f(x) \cdot g(1 - x) = 0.$$

Otherwise, there must exist some  $y \in K^\times$  such that  $0 < v(y) < v(x)$  and some  $g \in \mathcal{I}$  such that  $g(y) \neq 0$ . Indeed, if this does not hold then  $[0, v(x)] \subset vK$  would be contained in  $v(\mathcal{I}^\perp)$ , which cannot happen since  $v = v_{\mathcal{I}}$ . With such a  $y$ , the argument above shows that  $f(1 - y) = 0$  while

$$f(1 - x) = f((1 - x) \cdot (1 - y)) = f(1 - (y + x \cdot (1 - y))).$$

But  $v(y + x \cdot (1 - y)) = v(y)$ , so, again, the argument above shows that  $f(1 - x) = 0$ . In other words,  $U_v^1 \subset f^\perp$ , hence  $f \in \mathcal{D}_v$ .  $\square$

**Proposition 3.7.** *Suppose that  $\mathcal{D}$  is a closed subspace of  $\mathcal{G}_K$  such that  $\mathcal{Z}_K(\mathcal{D}) \neq \mathcal{D}$ . Then  $\mathcal{I} := \mathcal{Z}_K(\mathcal{D})$  is valuative and  $\mathcal{D} \subset \mathcal{D}_v$  for  $v = v_{\mathcal{I}}$ .*

*Proof.* By Lemma 3.6 it suffices to show that  $\mathcal{I}$  is valuative. Since  $\mathcal{I} \neq \mathcal{D}$ , there exists  $f_1, f_2 \in \mathcal{D}$  such that  $\mathcal{R}_K(f_1, f_2)$  does not hold. Put  $\mathcal{D}_i = \mathcal{I} + \mathbb{Q} \cdot f_i$ . Then  $\mathcal{D}_i$  are both closed subspaces of  $\mathcal{G}_K$ , and the following hold:

- (1)  $\mathcal{I}$  has codimension 1 in  $\mathcal{D}_i$ .
- (2)  $\mathcal{I}$  has codimension 2 in  $\mathcal{D}_1 + \mathcal{D}_2$ .
- (3)  $\mathcal{I} = \mathcal{D}_1 \cap \mathcal{D}_2$ .
- (4) For all  $f, g \in \mathcal{D}_i$ , one has  $\mathcal{R}_K(f, g)$ .

By Theorem 3.1, there exist valuations  $v_i$  and closed subspaces  $\mathcal{I}_i$  of  $\mathcal{D}_i$  of codimension  $\leq 1$  such that  $\mathcal{D}_i \subset \mathcal{D}_{v_i}$  and  $\mathcal{I}_i \subset \mathcal{I}_{v_i}$ .

If  $v_1$  and  $v_2$  are not comparable, then, letting  $v$  denote their infimum, one has  $U_{v_1}^1 \cdot U_{v_2}^1 = U_v$  by the approximation theorem for independent valuations. It follows that  $\mathcal{D}_{v_1} \cap \mathcal{D}_{v_2} = \mathcal{I}_v$ , hence  $\mathcal{I} \subset \mathcal{I}_v$ , thereby concluding the proof.

So, assume without loss of generality that  $v_1 \leq v_2$  hence

$$\mathcal{I}_{v_1} \subset \mathcal{I}_{v_2} \subset \mathcal{D}_{v_2} \subset \mathcal{D}_{v_1}.$$

Assume for a contradiction that  $\mathcal{I}$  is *not* valutive. Then  $\mathcal{I}$  is not contained in  $\mathcal{I}_{v_i} \cap \mathcal{D}_i$ , so we may find  $g_i \in \mathcal{I}_{v_i} \cap \mathcal{D}_i$  such that  $\mathcal{D}_i = \mathcal{I} + \mathbb{Q} \cdot g_i$ . In particular, we have

$$\mathcal{D}_1 + \mathcal{D}_2 = \mathcal{I} + \mathbb{Q} \cdot g_1 + \mathbb{Q} \cdot g_2$$

while  $g_1, g_2 \in \mathcal{I}_{v_2}$ . It follows that  $\mathcal{D}_1 + \mathcal{D}_2 \subset \mathcal{D}_{v_2}$  and that the image of  $\mathcal{D}_1 + \mathcal{D}_2$  in  $\mathcal{D}_{v_2}/\mathcal{I}_{v_2}$  agrees with the image of  $\mathcal{D}_2$  in this quotient, which has dimension  $\leq 1$ . This together with Lemma 3.2 would imply that  $\mathcal{R}_K(f, g)$  holds for all  $f, g \in \mathcal{D}_1 + \mathcal{D}_2$ , which is impossible since  $f_1, f_2 \in \mathcal{D}_1 + \mathcal{D}_2$ .  $\square$

**3.3. Visible valuations.** Let  $T$  be a subgroup of  $K^\times$  and  $v$  a valuation of  $K$ . We shall say that  $v$  is *T-visible* provided that the following hold:

- (1) One has  $\mathcal{I}_v \cap \mathcal{G}_{K|T} = \mathcal{Z}_K(\mathcal{D}_v \cap \mathcal{G}_{K|T}) \neq \mathcal{D}_v \cap \mathcal{G}_{K|T}$ .
- (2) One has  $v = v_{\mathcal{I}}$  for  $\mathcal{I} = \mathcal{I}_v \cap \mathcal{G}_{K|T}$ .

These are precisely the valuations  $v$  for which we will be able to characterize  $\mathcal{I}_v \cap \mathcal{G}_{K|T}$  and  $\mathcal{D}_v \cap \mathcal{G}_{K|T}$  using the relation  $\mathcal{R}_K$  restricted to  $\mathcal{G}_{K|T}$ , as we show in the following theorem. If  $H$  is the saturation of  $T$ , then  $\mathcal{G}_{K|T} = \mathcal{G}_{K|H}$  hence a valuation is *T-visible* if and only if it is *H-visible*. When  $T = k^\times$  for a subfield  $k$  of  $K$ , then we shall say “*k-visible*” instead of “*k*<sup>×</sup>-visible.” When  $T = \{1\}$  is trivial, we shall say “visible” instead of “ $\{1\}$ -visible.”

**Theorem 3.8.** *Let  $T$  be a subgroup of  $K^\times$  and  $\mathcal{D} \subset \mathcal{G}_{K|T}$  be a closed subspace. There exists a  $T$ -visible valuation of  $K$  such that  $\mathcal{Z}_K(\mathcal{D}) = \mathcal{I}_v \cap \mathcal{G}_{K|T}$  and  $\mathcal{D} = \mathcal{D}_v \cap \mathcal{G}_{K|T}$  if and only if the following conditions hold:*

- (1) *One has  $\mathcal{Z}_K(\mathcal{D}) \neq \mathcal{D}$ .*
- (2) *One has  $\mathcal{C}_K(\mathcal{Z}_K(\mathcal{D})) \cap \mathcal{G}_{K|T} = \mathcal{D}$ .*

*Proof.* First suppose that  $v$  is indeed *T-visible*. Put  $\mathcal{D} := \mathcal{D}_v \cap \mathcal{G}_{K|T}$  and  $\mathcal{I} := \mathcal{I}_v \cap \mathcal{G}_{K|T}$ . By assumption, we have  $\mathcal{I} = \mathcal{Z}_K(\mathcal{D}) \neq \mathcal{D}$  and by Lemma 3.6 we have  $\mathcal{C}_K(\mathcal{I}) = \mathcal{D}_v$  since  $v = v_{\mathcal{I}}$ , hence both conditions (1) and (2) hold true.

Conversely, suppose that  $\mathcal{D}$  satisfies conditions (1) and (2) and put  $\mathcal{I} := \mathcal{Z}_K(\mathcal{D})$ . By Proposition 3.7 and condition (1),  $\mathcal{I}$  is valutive and, setting  $v = v_{\mathcal{I}}$ , one has  $\mathcal{D} \subset \mathcal{D}_v$ . Lemma 3.6 shows that  $\mathcal{C}_K(\mathcal{I}) = \mathcal{D}_v$  hence condition (2) implies that  $\mathcal{D} = \mathcal{D}_v \cap \mathcal{G}_{K|T}$ . We also know that  $\mathcal{I}_v \cap \mathcal{G}_{K|T} \subset \mathcal{Z}_K(\mathcal{D}_v \cap \mathcal{G}_{K|T}) = \mathcal{I}$  by Lemma 3.2, while  $\mathcal{I} \subset \mathcal{I}_v \cap \mathcal{G}_{K|T}$  because  $v = v_{\mathcal{I}}$ . Thus  $\mathcal{I} = \mathcal{I}_v \cap \mathcal{G}_{K|T}$ . The fact that  $v$  is *T-visible* follows directly from the definition and the observations above.  $\square$

**3.4. Abundant visibility.** This section shows that fields of higher transcendence degree have an abundance of visible valuations.

**Lemma 3.9.** *Suppose that  $k$  is a subfield of  $K$ , and let  $T$  be any subgroup of  $K^\times$  which is contained in  $k^\times$ . Assume that  $\text{trdeg}(K|k) \geq 1$ , and that  $v$  is a valuation of  $K$  such that  $\mathcal{G}_{K|T} \subset \mathcal{D}_v$ . Then  $v$  is trivial.*

*Proof.* If not, then there exists some  $t \in K$  which is transcendental over  $k$  such that  $v(t) > 0$ . Thus  $1 + t \in \mathcal{U}_v^1 \subset \mathcal{D}_v^\perp \subset \mathcal{G}_{K|T}^\perp \subset \mathcal{G}_{K|k}^\perp$ . But  $\mathcal{G}_{K|k}^\perp$  is the saturation of  $k^\times$  in  $K^\times$ . This implies that  $1 + t$  is algebraic over  $k$ , which is impossible.  $\square$

**Proposition 3.10.** *Suppose that  $k$  is a subfield of  $K$ , and let  $T$  be any subgroup of  $K^\times$  which is contained in  $k^\times$ . Let  $v$  be a valuation of  $K$  such that the saturation of  $vk$  in  $vK$  contains no nontrivial convex subgroups and such that  $\text{trdeg}(Kv|kv) \geq 1$ . Then  $v$  is visible over  $T$ .*



*Proof.* Since the saturation of  $T$  in  $K^\times$  is contained in the radical closure of  $k$  in  $K$ , we may assume without loss of generality that  $T$  is saturated and that  $k$  is radically closed in  $K$ . In particular,  $k^\times$  is also saturated in  $K^\times$  and hence  $k^\times = \mathcal{G}_{K|k}^\perp$  while  $T = \mathcal{G}_{K|T}^\perp$ . By Lemma 3.4, we see that the saturation of  $vk$  in  $vK$  is  $v(\mathcal{I}_0^\perp)$  where  $\mathcal{I}_0 = \mathcal{I}_v \cap \mathcal{G}_{K|k}$ . On the other hand, our assumption ensures that  $\mathcal{G}_{K|k} \subset \mathcal{G}_{K|T}$  hence  $\mathcal{I}_0 \subset \mathcal{I}_1 := \mathcal{I}_v \cap \mathcal{G}_{K|T}$ , thus  $\mathcal{I}_1^\perp \subset \mathcal{I}_0^\perp$ . Thus  $v(\mathcal{I}_1^\perp)$  contains no nontrivial convex subgroups, so that  $v$  is indeed the valuation associated to  $\mathcal{I}_1$  due to the characterization from Lemma 3.3.

We must show that

$$\mathcal{I}_v \cap \mathcal{G}_{K|T} = \mathcal{Z}_K(\mathcal{D}_v \cap \mathcal{G}_{K|T}) \neq \mathcal{D}_v \cap \mathcal{G}_{K|T}.$$

Recall that  $\mathcal{D}_v \cap \mathcal{G}_{K|T} / \mathcal{I}_v \cap \mathcal{G}_{K|T} \cong \mathcal{G}_{Kv|Tv}$ . Since  $Tv \subset kv^\times$  we find that  $\mathcal{G}_{Kv|kv} \subset \mathcal{G}_{Kv|Tv}$ , while  $\text{trdeg}(Kv|kv) \geq 1$  ensures that  $\mathcal{G}_{Kv|kv}$  is infinite dimensional. Thus  $\mathcal{I}_v \cap \mathcal{G}_{K|T} \neq \mathcal{D}_v \cap \mathcal{G}_{K|T}$ .

Thus all that remains to show is that  $\mathcal{I}_v \cap \mathcal{G}_{K|T} = \mathcal{Z}_K(\mathcal{D}_v \cap \mathcal{G}_{K|T})$ . Note that  $\mathcal{I}_v \cap \mathcal{G}_{K|T} \subset \mathcal{Z}_K(\mathcal{D}_v \cap \mathcal{G}_{K|T})$  by Lemma 3.2, so we only need to show the other inclusion. As noted above, the image of the composition

$$\mathcal{D}_v \cap \mathcal{G}_{K|T} \hookrightarrow \mathcal{D}_v \twoheadrightarrow \mathcal{D}_v / \mathcal{I}_v \cong \mathcal{G}_{Kv}$$

is precisely  $\mathcal{G}_{Kv|Tv}$ . By Lemma 3.2, it follows that the image of  $\mathcal{Z}_K(\mathcal{D}_v \cap \mathcal{G}_{K|T})$  in  $\mathcal{G}_{Kv|Tv}$  is precisely  $\mathcal{Z}_{Kv}(\mathcal{G}_{Kv|Tv})$ . It suffices to show that this image, or equivalently  $\mathcal{Z}_{Kv}(\mathcal{G}_{Kv|Tv})$ , is trivial.

We now have two cases to consider. If  $\mathcal{Z}_{Kv}(\mathcal{G}_{Kv|Tv}) = \mathcal{G}_{Kv|Tv}$  then by Theorem 3.1, there exists a valuation  $w$  of  $Kv$  such that  $\mathcal{G}_{Kv|Tv} \subset \mathcal{D}_w$  and such that  $\mathcal{I}_w \cap \mathcal{G}_{Kv|Tv}$  has codimension  $\leq 1$  in  $\mathcal{G}_{Kv|Tv}$ . Lemma 3.9 shows that  $w$  is trivial and thus  $\mathcal{I}_w$  is trivial, which is impossible since  $\mathcal{G}_{Kv|kv} \subset \mathcal{G}_{Kv|Tv}$  is infinite dimensional.

Thus we have  $\mathcal{Z}_{Kv}(\mathcal{G}_{Kv|Tv}) \neq \mathcal{G}_{Kv|Tv}$ . In this case, Proposition 3.7 shows that  $\mathcal{Z}_{Kv}(\mathcal{G}_{Kv|Tv})$  is valutive and, letting  $w$  denote the associated valuation, one has  $\mathcal{G}_{Kv|Tv} \subset \mathcal{D}_w$ . Again, Lemma 3.9 shows that  $w$  is trivial, hence  $\mathcal{I}_w$  is trivial, so  $\mathcal{Z}_{Kv}(\mathcal{G}_{Kv|Tv})$  is trivial as well. This concludes the proof of the proposition.  $\square$

#### 4. ALGEBRAIC DEPENDENCE

Let  $k$  be a relatively algebraically closed subfield of  $K$ . Our goal in this section is to provide a characterization of algebraic dependence (over  $k$ ) in  $K$  using the algebra  $\mathcal{K}_*(K|k)$ , or equivalently, using the structure  $\mathcal{A}(K|k)$ .

**4.1. Milnor-closed subspaces.** Let  $T$  be a subgroup of  $K^\times$  and  $\mathcal{H}$  a subspace of  $\mathcal{K}_{K|T}$ . We say that  $\mathcal{H}$  is *Milnor-closed* provided that for all nontrivial  $s \in \mathcal{H}$  and  $t \in \mathcal{K}_{K|T}$  such that  $\{s, t\} = 0$  in  $\mathcal{K}_2(K|T)$ , one has  $t \in \mathcal{H}$  as well. Any subset  $S$  of  $\mathcal{K}_{K|T}$  has a *Milnor-closure* which is the smallest Milnor-closed subspace  $\mathcal{H}$  of  $\mathcal{K}_{K|T}$  that contains  $S$ . Explicitly, the Milnor-closure of  $S$  can be computed as a union

$$\mathcal{H} = \bigcup_{n=0}^{\infty} \mathcal{H}_n$$

where  $\mathcal{H}_0$  is the subspace generated by  $S$ , and  $\mathcal{H}_{n+1}$  is the subspace generated by  $\mathcal{H}_n$  and all  $t \in \mathcal{K}_{K|T}$  such that there exists some nontrivial  $s \in \mathcal{H}_n$  where  $\{s, t\} = 0$  in  $\mathcal{K}_2(K|T)$ .

**Lemma 4.1.** *Let  $\mathcal{H}$  be a Milnor-closed subspace of  $\mathcal{K}_{K|T}$ . Let  $H$  be the preimage of  $\mathcal{H}$  with respect to the canonical map  $K^\times \rightarrow \mathcal{K}_{K|T}$ . Then  $H$  contains the following:*

- (1) *The saturation of  $T$ .*
- (2) *Elements of  $K^\times$  of the form  $a + b \cdot h$  for any  $h \in H$  whose image in  $\mathcal{K}_{K|T}$  is nontrivial and any  $a, b$  in the saturation of  $T$ .*

*Proof.* Let  $R$  denote the saturation of  $T$ . It is clear that  $H$  contains  $R$ . Recall that we have a natural morphism of graded rings

$$K_*^M(K|R) \rightarrow \mathcal{K}_*(K|R) = \mathcal{K}_*(K|T).$$

Consider elements of  $K^\times$  the form  $a + b \cdot h$  as described in (3). We calculate some symbols in  $K_2^M(K|R)$ :

$$\begin{aligned} \{h, a + b \cdot h\} &= \{h, 1 - (-b \cdot a^{-1}) \cdot h\} \\ &= \{(-b \cdot a^{-1}) \cdot h, 1 - (-b \cdot a^{-1}) \cdot h\} \\ &= 0. \end{aligned}$$

The first equality follows from the fact that  $a^{-1} \in R$ , the second from the fact that  $-b \cdot a^{-1} \in R$ , and the last from the Steinberg relations in Milnor K-theory. Since  $\mathcal{H}$  is Milnor-closed and  $h \in H$  it follows that  $a + b \cdot h \in H$  as well.  $\square$

#### 4.2. Detecting algebraic dependence.

**Lemma 4.2.** *Let  $v$  be a valuation on  $K$ . The canonical map*

$$\wedge^*(\mathcal{K}_{K|U_v}) \rightarrow \mathcal{K}_*(K|U_v)$$

*given by the identity in degree one is an isomorphism.*

*Proof.* Note that the map is surjective and that  $U_v = \mathcal{I}_v^\perp$ . We will identify  $\mathcal{K}_{K|U_v}$  with  $\mathbb{Q} \otimes_{\mathbb{Z}} vK$  and for  $t \in K^\times$ , we abuse the notation and write  $v(t)$  for the image of  $t$  in  $\mathbb{Q} \otimes_{\mathbb{Z}} vK = \mathcal{K}_{K|U_v}$ . The kernel of the map in question is generated by  $r_{x,y} := v(x) \wedge v(y)$  where  $x, y \in K^\times$  satisfy  $x + y = 1$ . We claim that all such  $r_{x,y}$  are already trivial. If  $v(x) = 0$  or  $v(y) = 0$  then  $r_{x,y} = 0$ , so there is nothing to show. If  $v(x) > 0$  then  $v(y) = 0$  so  $r_{x,y} = 0$ , and similarly if  $v(y) > 0$ . Otherwise,  $v(x) < 0$  and  $v(y) = v(x)$  so again  $r_{x,y} = 0$ . In any case, the kernel is trivial, so the map in question is an isomorphism.  $\square$

**Lemma 4.3.** *Suppose that  $k$  is a subfield of  $K$  and  $T$  is a subgroup of  $K^\times$  which is contained in  $k^\times$ . Let  $t_1, \dots, t_n \in K^\times$  be given, and let  $\bar{t}_i$  denote the image of  $t_i$  in  $\mathcal{K}_{K|T}$ . If  $\{\bar{t}_1, \dots, \bar{t}_n\} = 0$  in  $\mathcal{K}_n(K|T)$ , then  $t_1, \dots, t_n$  are algebraically dependent over  $k$ .*

*Proof.* If  $t_1, \dots, t_n$  are algebraically independent, then we may find a  $k$ -valuation  $v$  on  $K$  such that  $v(t_1), \dots, v(t_n)$  are linearly independent in  $\mathbb{Q} \otimes_{\mathbb{Z}} vK = \mathcal{K}_{K|U_v}$ . For example, we can take the discrete rank  $n$  valuation on  $k(t_1, \dots, t_n)$  associated to the regular sequence  $(t_1, \dots, t_n)$  and choose  $v$  to be some prolongation of this valuation to  $K$ . Since  $T \subset k^\times \subset U_v$ , the assertion follows from Lemma 4.2.  $\square$

**4.3. Geometric lattices.** Suppose that  $k$  is a relatively algebraically closed subfield of  $K$ . The collection of all relatively algebraically closed subextension of  $K|k$  will be denoted by  $\mathbb{G}(K|k)$ . This is a *complete* lattice, with respect to inclusion of subfields of  $K$ , meaning that any set has a greatest lower bound (the infimum) and a smallest upper bound (the supremum). In this case, the infimum is computed by taking intersections in  $K$ , and the

supremum is computed by taking the relative algebraic closure of the compositum in  $K$ . We call  $\mathbb{G}(K|k)$  the *geometric lattice* associated to  $K|k$ .

For a field  $F$ , write  $F^i$  for the perfect closure of  $F$ . Note that restriction along  $K \hookrightarrow K^i$  induces an isomorphism of geometric lattices

$$\mathbb{G}(K^i|k^i) \cong \mathbb{G}(K|k),$$

where the inverse is given by  $M \mapsto M^i$ . We will make use of the following result from [10], which fundamentally relies on the work of Evans-Hrushovski [8, 9] based on the *group configuration theorem*. This theorem should be thought of as an analogue of the fundamental theorem of projective geometry, but for an incidence geometry associated to  $\mathbb{G}(K|k)$  as opposed to a projective space.

**Theorem 4.4** ([10], Theorem 4.2). *Suppose that  $K|k$  and  $L|l$  are relatively algebraically closed extensions of fields. Assume that  $\text{trdeg}(K|k) \geq 5$ , and that  $\varphi : \mathbb{G}(K|k) \cong \mathbb{G}(L|l)$  is an isomorphism of geometric lattices. Then there exists an isomorphism  $\Phi : K^i \cong L^i$  of fields satisfying  $\Phi(k^i) = l^i$  such that  $\varphi(M) = \Phi(M^i) \cap L$  for all  $M \in \mathbb{G}(K|k)$ . Furthermore,  $\Phi$  is unique with these properties up-to composition with some power of the  $p$ -power Frobenius  $x \mapsto x^p$ , where  $p$  is the characteristic exponent of  $K$ .*

*Proof.* Since  $\mathbb{G}(K|k) \cong \mathbb{G}(L|l)$ , we have  $\text{trdeg}(L|l) = \text{trdeg}(K|k) \geq 5$ , as the transcendence degree of a relatively algebraically closed field extension is the Krull dimension of the associated geometric lattice.

The only part that doesn't follow immediately from [10, Theorem 4.2] is that [8, 9, 10] all write  $\mathbb{G}(K|k)$  for the *combinatorial geometry* associated to  $K|k$  as opposed to the geometric lattice as we have defined above. But the two approaches are easily seen to be equivalent (this is a very well-known fact of matroid theory).

Indeed, let us write  $\mathbb{G}'(K|k)$  for the combinatorial geometry associated to  $K|k$ . This object refers to the set of all relatively algebraically closed subextensions  $M$  of  $K|k$  such that  $\text{trdeg}(M|k) = 1$ , and  $\mathbb{G}'(K|k)$  is endowed with a *closure operator*  $\text{cl}$  which associates to a subset  $S \subset \mathbb{G}'(K|k)$  the set  $\text{cl}(S)$  of all  $M \in \mathbb{G}'(K|k)$  such that  $M \subset \overline{k(S)} \cap K$ . A subset  $S$  of  $\mathbb{G}'(K|k)$  is called *closed* provided that  $\text{cl}(S) = S$ .

One can functorially recover  $\mathbb{G}'(K|k)$  from  $\mathbb{G}(K|k)$ , and vice-versa, as follows. First note that  $\mathbb{G}(K|k)$  has a unique minimal element  $\perp$  corresponding to the field  $k$ . Next, note that  $\mathbb{G}'(K|k)$  is the set of *atoms* of  $\mathbb{G}(K|k)$ , i.e. the set of elements  $M$  of  $\mathbb{G}(K|k)$  which are different from  $\perp$  and minimal with that property. The closure operator  $\text{cl}$  is obtained using the lattice structure of  $\mathbb{G}(K|k)$  as follows:

$$\text{cl}(S) = \{M \in \mathbb{G}'(K|k) \mid M \leq \text{Sup}(S)\},$$

where  $\text{Sup}(S)$  denotes the supremum of  $S$  in  $\mathbb{G}(K|k)$ . Conversely, one may identify  $\mathbb{G}(K|k)$  with the lattice of closed subsets of  $\mathbb{G}'(K|k)$ .

Going back to the context of the theorem, we have an isomorphism  $\varphi : \mathbb{G}(K|k) \cong \mathbb{G}(L|l)$  of lattices, which induces an isomorphism  $\varphi' : \mathbb{G}'(K|k) \cong \mathbb{G}'(L|l)$  of combinatorial geometries. By [10, Theorem 4.2(ii)], there exists an isomorphism  $\Phi : K^i \cong L^i$  of fields with  $\Phi(k^i) = l^i$  such that for all  $M \in \mathbb{G}'(K|k)$ , one has  $\varphi'(M) = \Phi(M^i) \cap L$ .

It remains to show that  $\varphi(M) = \Phi(M^i) \cap L$  for all  $M \in \mathbb{G}(K|k)$ , but this follows easily from the fact that  $\mathbb{G}(K|k)$  and  $\mathbb{G}(L|l)$  are *atomistic* lattices, meaning that every element  $M$  of  $\mathbb{G}(K|k)$  is the supremum of the atoms it bounds from above (and similarly for  $\mathbb{G}(L|l)$ ).  $\square$

**4.4. Geometric subspaces.** Suppose that  $k$  is a relatively algebraically closed subfield of  $K$ . Let  $L$  be any subextension of  $K|k$ . Then the canonical map

$$\mathcal{K}_{L|k} \rightarrow \mathcal{K}_{K|k}$$

is injective, and we will identify  $\mathcal{K}_{L|k}$  with its image in  $\mathcal{K}_{K|k}$ . A subspace of  $\mathcal{K}_{K|k}$  will be called *geometric* if it is of the form  $\mathcal{K}_{L|k}$  for some *relatively algebraically closed* subextension  $L$  of  $K|k$ . The collection of all geometric subspaces of  $\mathcal{K}_{K|k}$  will be denoted by  $\mathbb{G}_{\mathcal{K}}(K|k)$ , considered as a poset with respect to inclusion in  $\mathcal{K}_{K|k}$ .

**Proposition 4.5.** *The canonical map  $\mathbb{G}(K|k) \rightarrow \mathbb{G}_{\mathcal{K}}(K|k)$  sending  $L \in \mathbb{G}(K|k)$  to  $\mathcal{K}_{L|k}$  is an order isomorphism  $\mathbb{G}(K|k) \cong \mathbb{G}_{\mathcal{K}}(K|k)$ . In particular,  $\mathbb{G}_{\mathcal{K}}(K|k)$  is also a complete lattice.*

*Proof.* This map is clearly monotone and surjective. Conversely, suppose that  $L_1$  and  $L_2$  are two elements of  $\mathbb{G}(K|k)$ , that  $\mathcal{K}_{L_1|k} \leq \mathcal{K}_{L_2|k}$  and that  $t \in L_1$  is some element. Letting  $\bar{t}$  denote the image of  $t$  in  $\mathcal{K}_{K|k}$ , we have  $\bar{t} \in \mathcal{K}_{L_1|k}$ , which is contained in  $\mathcal{K}_{L_2|k}$ . Thus there exists some positive integer  $n$  and some constant  $c \in k^\times$  such that  $c \cdot t^n \in L_2^\times$ . This implies that  $t$  is algebraic over  $L_2$  and thus  $t \in L_2$ . In other words, we have  $L_1 \leq L_2$  in  $\mathbb{G}(K|k)$ . In particular, the map in question is then injective, hence bijective, and its inverse is also monotone.  $\square$

**4.5. Milnor closure vs. algebraic closure.** We continue to work with a relatively algebraically closed subfield  $k$  of  $K$ . Recall that we have a canonical pairing

$$\mathcal{K}_{K|k} \times \mathcal{G}_{K|k} \rightarrow \mathbb{Q}.$$

For the rest of this subsection, we will use the notation  $\mathcal{E}^\perp$  to denote the orthogonal of a subspace  $\mathcal{E} \subset \mathcal{G}_{K|k}$  with respect to the above pairing. Although this overloads the notation  $(-)^\perp$  introduced previously, it should be clear from context when  $\mathcal{E}^\perp$  refers to a subspace of  $\mathcal{K}_{K|k}$  as opposed to a subgroup of  $K^\times$ .

The following theorem is the technical core of this paper. It provides a characterization of the elements  $\mathbb{G}_{\mathcal{K}}(K|k)$  as subspaces of  $\mathcal{K}_{K|k}$ .

**Theorem 4.6.** *Assume that  $\text{trdeg}(K|k) \geq 2$ . Let  $k$  be a relatively algebraically closed subfield of  $K$ , and let  $\mathcal{H}$  be a Milnor-closed subspace of  $\mathcal{K}_{K|k}$ . Let  $H$  denote the preimage of  $\mathcal{H}$  with respect to the map  $K^\times \rightarrow \mathcal{K}_{K|k}$ , and let  $L$  denote the relative algebraic closure of  $k(H)$  in  $K$ . Let  $\mathcal{V}_{\mathcal{H}}$  denote the collection of all closed subspaces  $\mathcal{D}$  of  $\mathcal{G}_{K|k}$  satisfying the following conditions:*

- (1) *One has  $\mathcal{Z}_K(\mathcal{D}) \neq \mathcal{D}$ .*
- (2) *One has  $\mathcal{C}_K(\mathcal{Z}_K(\mathcal{D})) \cap \mathcal{G}_{K|k} = \mathcal{D}$ .*
- (3) *One has  $\mathcal{D}^\perp \cap \mathcal{H} = 0$ .*

*Then one has*

$$\mathcal{K}_{L|k} = \bigcap_{\mathcal{D} \in \mathcal{V}_{\mathcal{H}}} \mathcal{Z}_K(\mathcal{D})^\perp.$$

*Conversely, any geometric subspace of  $\mathcal{K}_{K|k}$  arises in this way. More precisely, if  $\mathcal{H} = \mathcal{K}_{L|k}$  for  $L \in \mathbb{G}(K|k)$ , then  $\mathcal{H}$  is Milnor-closed and one has*

$$\mathcal{H} = \bigcap_{\mathcal{D} \in \mathcal{V}_{\mathcal{H}}} \mathcal{Z}_K(\mathcal{D})^\perp.$$

*Proof.* Let  $\mathcal{V}_{\mathcal{H}}$  be as in the statement of the theorem. By Theorem 3.8, the subspaces  $\mathcal{D} \in \mathcal{V}_{\mathcal{H}}$  all have the form  $\mathcal{D} = \mathcal{D}_v \cap \mathcal{G}_{K|k}$  where  $v$  is a  $k$ -visible valuation of  $K$  such that  $\mathcal{U}_{v|k}^1 \cap \mathcal{H} = 0$ . Furthermore, in this case one has  $\mathcal{Z}_K(\mathcal{D}) = \mathcal{I}_v \cap \mathcal{G}_{K|k}$  hence also  $\mathcal{Z}_K(\mathcal{D})^\perp = \mathcal{U}_{v|k}$ . Thus, the intersection in question is precisely

$$\Delta := \bigcap_v \mathcal{U}_{v|k}$$

where  $v$  varies over the  $k$ -visible valuations of  $K$  such that  $\mathcal{U}_{v|k}^1 \cap \mathcal{H} = 0$ .

Let us first handle the case where  $\mathcal{H} = 0$ , so  $L = k$ . By the above discussion, it suffices to show that for every nontrivial  $x \in \mathcal{K}_{K|k}$ , there exists some  $k$ -visible valuation  $v$  of  $K$  such that  $x \notin \mathcal{U}_{v|k}$ . Replace  $x$  by  $n \cdot x$  for some positive integer  $n$  to assume without loss of generality that  $x$  is the image of a transcendental element  $t \in K^\times$ . Extend  $t$  to a transcendence base

$$\mathcal{B} = \{t\} \cup \mathcal{B}_0$$

of  $K|k$ , write  $M = k(\mathcal{B}_0)$ , and let  $v$  be a prolongation of the  $t$ -adic valuation on  $M(t)$  to  $K$ . Note that  $Kv|Mv$  is algebraic and  $Mv = M$  while  $vk = 0$ . By Proposition 3.10 and our assumptions on  $\text{trdeg}(K|k)$ , we see that  $v$  is indeed visible, while  $v(t) > 0$  by construction. Finally, if  $x \in \mathcal{U}_{v|k}$  then  $c \cdot t \in \mathcal{U}_v$  for some  $c \in k^\times$ , but  $v(c \cdot t) = v(t) > 0$ , so this cannot happen. Thus the assertion of the theorem holds true in the case where  $\mathcal{H} = 0$ . Assume for the rest of the proof that  $\mathcal{H} \neq 0$ .

In order to show that  $\Delta = \mathcal{K}_{L|k}$ , it suffices to show that a  $k$ -visible valuation  $v$  of  $K$  is trivial on  $L$  if and only if  $\mathcal{U}_{v|k}^1 \cap \mathcal{H} = 0$ . Indeed, in this case  $\Delta = \bigcap_v \mathcal{U}_{v|k}$  where  $v$  varies over the  $k$ -visible valuations which are trivial on  $L$ . Hence  $\mathcal{K}_{L|k} \subset \Delta$ . If  $x \in \mathcal{K}_{K|k} \setminus \mathcal{K}_{L|k}$ , then choose some  $t \in K \setminus L$  such that the image of  $t$  in  $\mathcal{K}_{K|k}$  is  $n \cdot x$  for some positive integer  $n$ . This  $t$  is then transcendental over  $L$ . Complete  $t$  to a transcendence basis  $\mathcal{B} = \{t\} \cup \mathcal{B}_0$  of  $K|L$ , put  $M := L(\mathcal{B}_0)$ , and let  $v$  be an extension of the  $t$ -adic valuation on  $M(t)$  to  $K$ . Arguing similarly to the above, we see that  $v$  is  $k$ -visible, trivial on  $L$ , and that  $x \notin \mathcal{U}_{v|k}$ . This shows that indeed,  $\Delta = \mathcal{K}_{L|k}$  provided that a  $k$ -visible valuation is trivial on  $L$  if and only if  $\mathcal{U}_{v|k}^1 \cap \mathcal{H} = 0$ .

Suppose that  $v$  is trivial on  $L$ . We have a canonical injective map

$$L^\times / k^\times \rightarrow Kv^\times / kv^\times$$

and after tensoring with  $\mathbb{Q}$  we obtain

$$\mathcal{K}_{L|k} \rightarrow \mathcal{K}_{Kv|kv}$$

which is again injective. This last map is precisely the composition

$$\mathcal{K}_{L|k} \hookrightarrow \mathcal{U}_{v|k} \twoheadrightarrow \mathcal{U}_{v|k} / \mathcal{U}_{v|k}^1 = \mathcal{K}_{Kv|kv}$$

and thus  $\mathcal{U}_{v|k}^1 \cap \mathcal{K}_{L|k} = 0$ . Since  $\mathcal{H} \subset \mathcal{K}_{L|k}$ , we also have  $\mathcal{U}_{v|k}^1 \cap \mathcal{H} = 0$ .

We must now show that a  $k$ -visible valuation  $v$  is trivial on  $L$  provided that  $\mathcal{U}_{v|k}^1 \cap \mathcal{H} = 0$ . *This is the crux of the proof.* So, assume that  $v$  is  $k$ -visible and *nontrivial* on  $L$ . Since  $L|k(H)$  is algebraic, it follows that  $v$  is nontrivial on  $k(H)$ .

We observe that  $k(H)^\times = H$  as subgroups of  $K^\times$ . Indeed, first note that  $k^\times \subset H$  and  $H$  is multiplicatively closed. Thus, it suffices to show that  $H \cup \{0\}$  is additively closed, and since  $H$  is a subgroup, for this it suffices to show that  $1 + t \in H \cup \{0\}$  whenever  $t \in H$ . If  $t \in k^\times$ , then this is obvious, and if not, then its image in  $\mathcal{K}_{K|k}$  is nontrivial, so that  $1 + t \in H$  by Lemma 4.1, using the assumption that  $\mathcal{H}$  is Milnor-closed.

In any case, we have  $k(H)^\times = H$ . Since  $\mathcal{H} \neq 0$ , hence  $k^\times \neq k(H)^\times = H$  there must exist some element  $t \in H \setminus k^\times$  such that  $v(t) > 0$ . Since  $1 + t \in H \setminus k^\times$  as well, the image of  $1 + t$  in  $\mathcal{K}_{K|k}$  is a nontrivial element of  $\mathcal{U}_{v|k}^1 \cap \mathcal{H}$ , showing that  $\mathcal{U}_{v|k}^1 \cap \mathcal{H} \neq 0$ .

The final assertion is easy. If  $\mathcal{H} = \mathcal{K}_{L|k}$  then  $\mathcal{H}$  is Milnor-closed by Lemma 4.3, while the preimage of  $\mathcal{K}_{L|k}$  in  $K^\times$  is  $L^\times$  since  $L$  is relatively algebraically closed in  $K$ . So, the first part of the theorem, which is already proved, gives the desired claim.  $\square$

**4.6. Recovering transcendence degree.** We continue with the context above where  $k$  is a relatively algebraically closed subfield of  $K$ .

**Lemma 4.7.** *Let  $d$  be any positive integer. Assume that there exists a closed subspace  $\mathcal{D}$  of  $\mathcal{G}_{K|k}$  such that the following conditions hold true:*

- (1) *One has  $\mathcal{Z}_K(\mathcal{D}) \neq \mathcal{D} = \mathcal{C}_K(\mathcal{Z}_K(\mathcal{D})) \cap \mathcal{G}_{K|k}$ .*
- (2) *One has  $d \leq \dim_K(\mathcal{Z}_K(\mathcal{D}))$ .*

*Then  $\text{trdeg}(K|k) \geq d$ .*

*Proof.* This is a simple consequence of Theorem 3.8 in conjunction with Abhyankar's inequality. By Theorem 3.8, there exists a  $k$ -visible valuation  $v$  of  $K$  such that  $\mathcal{I} := \mathcal{Z}_K(\mathcal{D}) = \mathcal{I}_v \cap \mathcal{G}_{K|k}$ . Dualizing  $\mathcal{I}$ , we obtain  $\mathcal{K}_{K|k}/\mathcal{U}_{v|k} = \mathbb{Q} \otimes_{\mathbb{Z}} (vK/vk)$ , and our assumption tells us that  $\dim_{\mathbb{Q}}(\mathbb{Q} \otimes_{\mathbb{Z}} (vK/vk)) \geq d$ . Choose  $t_1, \dots, t_d \in K$  whose images in  $\mathbb{Q} \otimes_{\mathbb{Z}} (vK/vk)$  are linearly independent, and consider  $L := k(t_1, \dots, t_d)$ , a finitely-generated subextension of  $K|k$ , which has the property that

$$\dim_{\mathbb{Q}}(\mathbb{Q} \otimes_{\mathbb{Z}} (vL/vk)) \geq d.$$

By Abhyankar's inequality, we find

$$d \leq \dim_{\mathbb{Q}}(\mathbb{Q} \otimes_{\mathbb{Z}} (vL/vk)) \leq \text{trdeg}(L|k) \leq \text{trdeg}(K|k),$$

as claimed in the lemma.  $\square$

**Lemma 4.8.** *Suppose that  $d$  is a positive integer and that  $\text{trdeg}(K|k) > d$ . Then there exists a closed subspace  $\mathcal{D}$  of  $\mathcal{G}_{K|k}$  such that the following conditions hold true:*

- (1) *One has  $\mathcal{Z}_K(\mathcal{D}) \neq \mathcal{D} = \mathcal{C}_K(\mathcal{Z}_K(\mathcal{D})) \cap \mathcal{G}_{K|k}$ .*
- (2) *One has  $d \leq \dim_K(\mathcal{Z}_K(\mathcal{D}))$ .*

*Proof.* Let  $t_1, \dots, t_d \in K$  be algebraically independent over  $k$ , and extend  $\{t_1, \dots, t_d\}$  to a transcendence base of  $K|k$  of the form

$$\mathcal{B} = \{t_1, \dots, t_d\} \cup \mathcal{B}_0.$$

By assumption on  $\text{trdeg}(K|k)$ , the set  $\mathcal{B}_0$  is nonempty. Put  $L := k(\mathcal{B}_0)$  and  $M := k(\mathcal{B}) = L(t_1, \dots, t_d)$ . Consider the valuation associated to the system of regular parameters  $(t_1, \dots, t_d)$  on  $M$ . This is a discrete rank  $d$  valuation which is trivial on  $L$ . Extend it in some way to a valuation  $v$  on  $K$ . Since  $\text{trdeg}(L|k) > 0$ , it follows from Proposition 3.10 that  $v$  is visible, while  $vk = 0$  and the images of  $t_1, \dots, t_d$  in  $\mathbb{Q} \otimes_{\mathbb{Z}} vK$  are rationally independent. Thus  $\dim_{\mathbb{Q}}(\mathcal{I}_v \cap \mathcal{G}_{K|k}) \geq d$  as well, while condition (1) follows from Theorem 3.8.  $\square$

We will use Lemmas 4.7 and 4.8 primarily to provide a *lower bound* on  $\text{trdeg}(K|k)$ , which will be required in order to apply Theorem 4.6. We summarize this observation in the following lemmas.

**Lemma 4.9.** *Suppose that  $K|k$  and  $L|l$  are two relatively algebraically closed field extensions, and that  $\mathcal{K}_*(K|k) \cong \mathcal{K}_*(L|l)$  as algebras. If  $d$  is a positive integer such that  $d < \text{trdeg}(K|k)$ , then  $d \leq \text{trdeg}(L|l)$ .*

*Proof.* The isomorphism  $\mathcal{K}_*(K|k) \cong \mathcal{K}_*(L|l)$  induces an isomorphism of structures  $\mathcal{A}(K|k) \cong \mathcal{A}(L|l)$  by Fact 2.1 and the surrounding discussion. By Lemma 4.8, there exists a closed subspace  $\mathcal{D}$  of  $\mathcal{G}_{K|k}$  such that

- (1) One has  $\mathcal{Z}_K(\mathcal{D}) \neq \mathcal{D} = \mathcal{C}_K(\mathcal{D}) \cap \mathcal{G}_{K|k}$ .
- (2) One has  $d \leq \dim_K(\mathcal{Z}_K(\mathcal{D}))$ .

Transferring this subspace across the isomorphism  $\mathcal{G}_{K|k} \cong \mathcal{G}_{L|l}$ , we obtain a closed subspace of  $\mathcal{G}_{L|l}$  satisfying the conditions of Lemma 4.7, which implies that  $d \leq \text{trdeg}(L|l)$ .  $\square$

**Lemma 4.10.** *Suppose that  $K|k$  and  $L|l$  are two relatively algebraically closed field extensions, and that  $\mathcal{K}_*(K|k) \cong \mathcal{K}_*(L|l)$  as algebras. Assume that  $\text{trdeg}(K|k) \geq 3$ . Then  $\text{trdeg}(K|k) = \text{trdeg}(L|l)$ .*

*Proof.* By Lemma 4.7, we see that  $\text{trdeg}(L|l) \geq 2$ , hence Theorem 4.6 applies to both  $K|k$  and  $L|l$ . This theorem provides us with an isomorphism of lattices

$$\mathbb{G}_{\mathcal{K}}(K|k) \cong \mathbb{G}_{\mathcal{K}}(L|l)$$

hence also  $\mathbb{G}(K|k) \cong \mathbb{G}(L|l)$  by Proposition 4.5. Since the transcendence degree of a relatively algebraically closed field extension is the Krull dimension of the corresponding geometric lattice, the claim follows.  $\square$

## 5. MAIN RESULTS

In this section, we present and prove the main results of this paper. We split up this section into three subsections:

- (1) The first regarding *relative results*, dealing with relatively algebraically closed field extensions  $K|k$  of sufficiently large transcendence degree, while using  $\mathcal{K}_*(K|k)$ .
- (2) The second regarding *absolute results*, dealing with arbitrary fields of sufficiently large *absolute* transcendence degree (i.e. transcendence degree over the prime subfield), while using  $\mathcal{K}_*(K)$ .
- (3) The third dealing with finitely-generated relatively algebraically closed extensions  $K|k$  over perfect fields and finitely-generated fields  $L$ , using  $\mathcal{K}_*^M(K|k)$ ,  $\mathcal{K}_*^M(L)$ .

In cases (1) and (2), our key tool is one of the main results of Evans-Hrushovski and Gismatullin [8, 9, 10], which we have summarized in Theorem 4.4 above. In case (3), our key tools will be the reconstruction result due to Cadoret-Pirutka [4, Theorem 4].

In the first two cases, we will only be able to recover fields *up-to inseparable extensions*, or more precisely, we will be able to recover the perfect closure of the field in question. Recall that we write  $F^i$  for its perfect closure of a field  $F$ . Of course, if  $F$  has characteristic zero, then  $F = F^i$ . If  $F$  has positive characteristic  $p$ , then one has

$$F^i = \bigcup_{n \geq 0} F^{1/p^n}.$$

Note that if  $K|k$  is a relatively algebraically closed extension of fields, then  $K^i|k^i$  is also relatively algebraically closed. The canonical map  $K^\times \rightarrow (K^i)^\times$  induces an *injective* map  $K^\times/k^\times \rightarrow (K^i)^\times/(k^i)^\times$ , and the induced map of  $\mathbb{Q}$ -modules  $\mathcal{K}_{K|k} \rightarrow \mathcal{K}_{K^i|k^i}$  is an *isomorphism*.

We first prove an auxiliary lemma that will be necessary to show certain *uniqueness* properties of the isomorphisms of fields that we obtain. This lemma has appeared before in [9, Theorem 1.1] and in [2, Lemma 13], although the proof of [9] is more complicated as it relies on the more general result [8, Theorem 2.2.2], while [2] has a blanket assumption that the fields in question have characteristic zero and the base field is algebraically closed. The argument we give is an adaptation of the proof from [2] which avoids the restrictions on the base field.

**Lemma 5.1.** *Let  $K|k$  be a relatively algebraically closed extension of fields. Suppose that  $x, y \in K$  are algebraically independent over  $k$ . Let  $a, b \in K \setminus k$  be two elements such that  $\text{trdeg}(k(a, x)|k) = 1$ ,  $\text{trdeg}(k(b, y)|k) = 1$  and  $\text{trdeg}(k(a \cdot b, x \cdot y)|k) = 1$ . Then there exist nonzero integers  $m$  and  $n$  such that, modulo  $k^\times$ , one has  $a^n = x^m$ ,  $b^n = y^m$ .*

*Proof.* Embed  $K^\times/k^\times$  into  $\overline{K}^\times/\overline{k}^\times$  to assume without loss of generality that both  $k$  and  $K$  are algebraically closed. The elements  $a, b$  and  $a \cdot b$  are all contained in the compositum  $\overline{k(x) \cdot k(y)} =: M \subset K$ . Since  $x$  and  $y$  are algebraically independent over  $k$ , we may identify

$$\text{Gal}(M|k(x, y)^i) = \text{Gal}(\overline{k(x)}|k(x)^i) \times \text{Gal}(\overline{k(y)}|k(y)^i),$$

where the two projections are the usual restriction maps. This lets us identify  $\text{Gal}(\overline{k(x)}|k(x)^i)$  with the subgroup of  $\text{Gal}(M|k(x, y)^i)$  which fixes  $\overline{k(y)}$  pointwise, and similarly with  $x$  and  $y$  interchanged.

With this identification, any  $\sigma \in \text{Gal}(\overline{k(x)}|k(x)^i)$  acts on  $a \cdot b$  as  $\sigma(a) \cdot b$ , and thus  $\sigma(a)/a = \sigma(a \cdot b)/(a \cdot b) =: t_\sigma$ . As  $x$  and  $x \cdot y$  are algebraically independent, we have

$$t_\sigma \in \overline{k(x)} \cap \overline{k(x \cdot y)} = k.$$

But  $\sigma(a) = t_\sigma \cdot a$  and  $k(x)^i(a)$  is a finite extension of  $k(x)^i$ , hence  $t_\sigma$  must be a root of unity. By symmetry, for any  $\tau \in \text{Gal}(\overline{k(y)}|k(y)^i)$ , we also have  $\tau(b) = s_\tau \cdot b$  for some root of unity  $s_\tau$ .

It follows that for all  $\gamma \in \text{Gal}(M|k(x, y)^i)$ , the elements  $\gamma(a)/a$  and  $\gamma(b)/b$  are both roots of unity. The action of  $\text{Gal}(M|k(x, y)^i)$  on  $k(x, y)^i(a, b)$  factors through a finite quotient, and thus we deduce that there exists some positive integer  $n_1$  such that for all  $\gamma \in \text{Gal}(M|k(x, y)^i)$ , one has  $\gamma(a^{n_1}) = a^{n_1}$  and  $\gamma(b^{n_1}) = b^{n_1}$ . In other words,  $\text{Gal}(M|k(x, y)^i)$  acts trivially on  $a^{n_1}$  and  $b^{n_1}$ , which implies that  $a^{n_1} \in k(x)^i$ ,  $b^{n_1} \in k(y)^i$ . Arguing similarly with  $x \cdot y$  in place of  $y$  and  $x^{-1}$  in place of  $x$ , we see that there exists an integer  $n_2$  such that  $a^{n_2} \in k(x)^i$  and  $(a \cdot b)^{n_2} \in k(x \cdot y)^i$ . Taking  $n = n_1 \cdot n_2$ , it follows that  $a^n \in k(x)^i$ ,  $b^n \in k(y)^i$  and  $(a \cdot b)^n \in k(x \cdot y)^i$ . Further replacing  $n$  by an integer of the form  $n \cdot p^k$ , where  $p$  is the characteristic exponent of  $k$ , we may assume that  $a^n \in k(x)$ ,  $b^n \in k(y)$  and  $(a \cdot b)^n \in k(x \cdot y)$ .

In particular, we may write

$$a^n = \prod_{c \in k} (x - c)^{m_c}, \quad b^n = \prod_{c \in k} (y - c)^{n_c},$$

modulo constants, where all but finitely many of the  $m_c, n_c \in \mathbb{Z}$  are zero. In particular, we must also have

$$(a \cdot b)^n = \prod_{c \in k} (x - c)^{m_c} \cdot (y - c)^{n_c},$$

modulo constants. But this element is contained in  $k(x \cdot y)$ , so the only irreducible polynomials from  $k[x, y]$  that may appear in the factorization of  $(a \cdot b)^n$  are of the form  $(x \cdot y - c)$  for some  $c \in k$ . Combining these observations, we deduce that  $m_c = n_c = 0$  for all  $c \in k^\times$ , and that



$m_0 = n_0$ . In other words, there exists an integer  $m$  such that  $a^n = x^m$  and  $b^n = y^m$  modulo constants, as required.  $\square$

Recall that  $\mathcal{K}_{K|k}$  is written *additively*. It will be useful to reinterpret the above lemma in the context of  $\mathcal{K}_{K|k}$ . First of all, if  $S$  is *any subset* of  $\mathcal{K}_{K|k}$ , we write  $\text{acl}_{K|k}(S)$  for the *smallest* element of  $\mathbb{G}_{\mathcal{K}}(K|k)$  which contains  $S$ . Since  $\mathbb{G}_{\mathcal{K}}(K|k)$  is a complete lattice, this  $\text{acl}_{K|k}(S)$  is well-defined. We say that two elements  $x, y \in \mathcal{K}_{K|k}$  are *dependent* provided that there exists a geometric subspace of the form  $\mathcal{K}_{L|k}$  where  $L \in \mathbb{G}(K|k)$  with  $\text{trdeg}(L|k) \leq 1$  such that  $x, y \in \mathcal{K}_{L|k}$ . If  $x, y$  are *not dependent* then we shall say that they are *independent* and in this case  $\text{acl}_{K|k}(x, y) = \mathcal{K}_{L|k}$  for some  $L \in \mathbb{G}(K|k)$  with  $\text{trdeg}(L|k) = 2$ . If  $x, y \in K^\times$  with images  $\bar{x}, \bar{y}$  in  $\mathcal{K}_{K|k}$ , then  $\bar{x}, \bar{y}$  are (in)dependent in the above sense if and only if  $x, y$  are algebraically (in)dependent over  $k$ .

**Lemma 5.2.** *Let  $x, y \in \mathcal{K}_{K|k}$  be two independent elements. Let  $a, b \in \mathcal{K}_{K|k}$  be two elements such that  $a \in \text{acl}_{K|k}(x)$ ,  $b \in \text{acl}_{K|k}(y)$  and  $a + b \in \text{acl}_{K|k}(x + y)$ . Then there exists some  $r \in \mathbb{Q}^\times$  such that  $a = r \cdot x$  and  $b = r \cdot y$ .*

*Proof.* The map  $\mathcal{K}_{K|k} \rightarrow \mathcal{K}_{\bar{K}|\bar{k}}$  is injective and compatible with the notion of (in)dependence introduced above. Also, the map

$$K_1^M(\bar{K}|\bar{k}) \rightarrow \mathcal{K}_{\bar{K}|\bar{k}}$$

is an *isomorphism* since  $\bar{K}^\times$  is divisible. By mapping into  $\mathcal{K}_{\bar{K}|\bar{k}}$  and identifying this with  $K_1^M(\bar{K}|\bar{k})$ , the assertion of this lemma reduces to that of Lemma 5.1.  $\square$

### 5.1. The relative case.

**Theorem 5.3.** *Let  $K|k$  be a relatively algebraically closed extension of fields satisfying  $\text{trdeg}(K|k) \geq 5$ . Then  $K^i|k^i$  are determined up-to isomorphism by the algebra  $\mathcal{K}_*(K|k)$ . More precisely, if  $L|l$  is another relatively algebraically closed extension of fields and*

$$\varphi_* : \mathcal{K}_*(K|k) \xrightarrow{\cong} \mathcal{K}_*(L|l)$$

*is an isomorphism of  $\mathbb{Q}$ -algebras, then there exists some  $r \in \mathbb{Q}^\times$  and an isomorphism of fields*

$$\Phi : K^i \xrightarrow{\cong} L^i$$

*such that  $\Phi(k^i) = l^i$  and the following diagram commutes:*

$$\begin{array}{ccc} \mathcal{K}_{K^i|k^i} & \xrightarrow{\Phi} & \mathcal{K}_{L^i|l^i} \\ \uparrow & & \uparrow \\ \mathcal{K}_{K|k} & \xrightarrow{r \cdot \varphi_1} & \mathcal{K}_{L|l}. \end{array}$$

Here, the vertical maps are those induced by the inclusions  $K \subset K^i$  and  $L \subset L^i$ , and the map labeled  $\Phi$  is induced by the isomorphism  $\Phi : K^i \cong L^i$  of fields. Finally,  $\Phi$  is unique with these properties up-to composition with some power of the  $p$ -power Frobenius  $x \mapsto x^p$ , where  $p$  is the characteristic exponent of  $K$ .

*Proof.* Let  $\varphi_*$  be as in the statement of the theorem and put  $\varphi = \varphi_1$ . By Lemma 4.10, we see that  $\text{trdeg}(L|l) = \text{trdeg}(K|k) \geq 5$ . By Theorem 4.6 and the discussion around Fact 2.1, we see

that the map  $\mathcal{H} \mapsto \varphi\mathcal{H}$  is an *isomorphism* of lattices  $\mathbb{G}_{\mathcal{K}}(K|k) \cong \mathbb{G}_{\mathcal{K}}(L|l)$ . By Proposition 4.5, we obtain an isomorphism  $\psi : \mathbb{G}(K|k) \cong \mathbb{G}(L|l)$  of geometric lattices satisfying

$$\varphi\mathcal{K}_{M|k} = \mathcal{K}_{\psi M|l}$$

for all  $M \in \mathbb{G}(K|k)$ . Thus, by [10, theorem 4.2] (see Theorem 4.4), there exists an isomorphism  $\Phi : K^i \cong L^i$  of fields satisfying the following conditions:

- (1)  $\Phi(k^i) = l^i$ .
- (2) One has  $\psi(M) = \Phi(M^i) \cap L$  for all  $M \in \mathbb{G}(K|k)$ .

And furthermore, this  $\Phi$  is unique up-to composition with some power of the  $p$ -power Frobenius  $x \mapsto x^p$ , where  $p$  is the characteristic exponent of  $K$ .

To conclude, we must show that there exists some  $r \in \mathbb{Q}^\times$  making the diagram from the statement of the theorem commute. Let  $\varphi' : \mathcal{K}_{K|k} \cong \mathcal{K}_{L|l}$  be the isomorphism induced by  $\Phi$ , i.e.  $\varphi'$  is the composition

$$\mathcal{K}_{K|k} \xrightarrow{\cong} \mathcal{K}_{K^i|k^i} \xrightarrow{\Phi} \mathcal{K}_{L^i|l^i} \xleftarrow{\cong} \mathcal{K}_{L|l}.$$

We must show that there exists some  $r \in \mathbb{Q}^\times$  such that  $r \cdot \varphi = \varphi'$ .

Let  $x, y \in \mathcal{K}_{K|k}$  be two independent elements. The pair  $\varphi(x), \varphi(y)$  is also independent, while the following pairs are all *dependent*:

$$\varphi(x), \varphi'(x); \varphi(y), \varphi'(y); \varphi(x) + \varphi(y), \varphi'(x) + \varphi'(y).$$

By Lemma 5.2, there exists some  $r(x, y) \in \mathbb{Q}^\times$  such that

$$r(x, y) \cdot \varphi(x) = \varphi'(x), \quad r(x, y) \cdot \varphi(y) = \varphi'(y).$$

Since  $\varphi(x)$  and  $\varphi(y)$  are also *linearly independent* over  $\mathbb{Q}$ , it follows that  $r(x, y) = r(y, x)$ .

We claim that  $r(x, y)$  does not depend on the choice of  $x, y$ . First, if  $z$  is another element which is independent from  $x$ , then  $r(x, y) = r(x, z)$  since  $r(x, y) \cdot \varphi(x) = r(x, z) \cdot \varphi(x)$ . Now, if  $x', y'$  is any other pair of independent elements, we show that  $r(x, y) = r(x', y')$  as follows:

- (1) If  $x, x'$  are *dependent*, then  $x', y$  are independent, hence

$$r(x, y) = r(y, x) = r(y, x') = r(x', y) = r(x', y').$$

- (2) If  $x, x'$  are *independent*, then

$$r(x, y) = r(x, x') = r(x', x) = r(x', y').$$

In any case,  $r(x, y)$  doesn't depend on the choice of  $x, y$ , and we put  $r := r(x, y)$  for some choice of an independent pair  $x, y$ .

Now, if  $x$  is any nontrivial element of  $\mathcal{K}_{K|k}$ , we may find some  $y \in \mathcal{K}_{K|k}$  which is independent from  $x$ , and observe that

$$r \cdot \varphi(x) = r(x, y) \cdot \varphi(x) = \varphi'(x).$$

Of course, this equality holds with  $x = 0$  as well, so we deduce that indeed  $r \cdot \varphi = \varphi'$ . This concludes the proof of the theorem.  $\square$

## 5.2. The absolute case.

**Theorem 5.4.** *Let  $K$  be a field whose absolute transcendence degree is at least five. Then  $K^i$  is determined up-to isomorphism by the algebra  $\mathcal{K}_*(K)$ . More precisely, suppose that  $L$  is another field and*

$$\varphi_* : \mathcal{K}_*(K) \xrightarrow{\cong} \mathcal{K}_*(L)$$

*is an isomorphism of  $\mathbb{Q}$ -algebras. Let  $k$  be the relative algebraic closure of the prime subfield of  $K$  and  $l$  the relative algebraic closure of the prime subfield of  $L$ . Then  $\varphi_*$  induces an isomorphism  $\overline{\varphi}_* : \mathcal{K}_*(K|k) \cong \mathcal{K}_*(L|l)$ . Furthermore, there exists some  $r \in \mathbb{Q}^\times$  and an isomorphism of fields*

$$\Phi : K^i \xrightarrow{\cong} L^i$$

*which necessarily satisfies  $\Phi(k^i) = l^i$ , such that the following diagram commutes:*

$$\begin{array}{ccc} \mathcal{K}_{K^i|k^i} & \xrightarrow{\Phi} & \mathcal{K}_{L^i|l^i} \\ \uparrow & & \uparrow \\ \mathcal{K}_{K|k} & \xrightarrow{r \cdot \overline{\varphi}_1} & \mathcal{K}_{L|l}. \end{array}$$

*Here, the vertical maps are those induced by the inclusions  $K \subset K^i$  and  $L \subset L^i$ , and the map labeled  $\Phi$  is induced by the isomorphism  $\Phi : K^i \cong L^i$  of fields. Finally,  $\Phi$  is unique with these properties up-to composition with some power of the  $p$ -power Frobenius  $x \mapsto x^p$ , where  $p$  is the characteristic exponent of  $K$ .*

By using Theorem 5.3, in order to prove Theorem 5.4 it suffices to give a characterization of the kernel of  $\mathcal{K}_K \rightarrow \mathcal{K}_{K|k}$  using  $\mathcal{K}_*(K)$ , where  $k$  is the relative algebraic closure of the prime subfield. If this kernel is denoted by  $\Delta$ , then one has

$$\mathcal{K}_*(K|k) = \frac{\mathcal{K}_*(K)}{\langle \Delta \rangle}$$

so that we can then apply Theorem 5.3. We describe the characterization of this kernel in the following proposition.

For the rest of this subsection, we will use the notation  $\mathcal{E}^\perp$  to denote the orthogonal of a subspace  $\mathcal{E} \subset \mathcal{G}_K$  with respect to the pairing

$$\mathcal{K}_K \times \mathcal{G}_K \rightarrow \mathbb{Q}.$$

As before, it should be clear from context when  $\mathcal{E}^\perp$  refers to a subspace of  $\mathcal{K}_K$  as opposed to a subgroup of  $K^\times$ .

**Proposition 5.5.** *Let  $K$  be any field, and let  $k$  denote the relative algebraic closure of the prime subfield of  $K$ . Assume that  $\text{trdeg}(K|k) \geq 2$ . Put  $\Delta := \ker(\mathcal{K}_K \rightarrow \mathcal{K}_{K|k})$ . For a nonzero  $t \in \mathcal{K}_K$ , write  $\mathcal{H}_t$  for the Milnor closure of the subset  $\{t\}$ . Let  $\mathcal{V}_t$  denote the collection of closed subspaces  $\mathcal{D} \subset \mathcal{G}_K$  satisfying the following conditions:*

- (1) *One has  $\mathcal{Z}_K(\mathcal{D}) \neq \mathcal{D}$ .*
- (2) *One has  $\mathcal{C}_K(\mathcal{Z}_K(\mathcal{D})) = \mathcal{D}$ .*
- (3) *One has  $\mathcal{H}_t \cap \mathcal{D}^\perp = 0$ .*

Then one has

$$\Delta = \bigcap_{t \in \mathcal{K}_K \setminus \{0\}} \bigcap_{\mathcal{D} \in \mathcal{V}_t} \mathcal{Z}_K(\mathcal{D})^\perp.$$

*Proof.* Note that  $\mathcal{H}_t$  remains unchanged if we replace  $t$  by  $n \cdot t$  for any nonzero integer  $n$ . Thus, we may restrict our attention to those nonzero  $t \in \mathcal{K}_K$  which are in the image of  $K^\times$ . Note also that the kernel of  $K^\times \rightarrow \mathcal{K}_K$  is the torsion subgroup of  $K^\times$ .

If  $K$  has positive characteristic then  $k^\times$  is the torsion of  $K^\times$  hence  $\mathcal{K}_*(K) = \mathcal{K}_*(K|k)$ . Letting  $t \in K \setminus k$  be given, we deduce from Theorem 4.6 that the intersection

$$\bigcap_{\mathcal{D} \in \mathcal{V}_t} \mathcal{Z}_K(\mathcal{D})^\perp$$

is precisely the geometric subspace  $\mathcal{K}_{L|k}$  where  $L$  is the relative algebraic closure of  $k(t)$  in  $K$ . As  $t$  varies, the intersection of all these geometric subspaces is trivial, so the assertion of the proposition follows.

Assume for the rest of the proof that  $K$  has characteristic zero. Let  $\mathcal{D}$  be a subspace satisfying conditions (1) and (2). By Theorem 3.8, we see that  $\mathcal{D} = \mathcal{D}_v$  for some visible (i.e.  $\{1\}$ -visible) valuation  $v$  of  $K$ , and that  $\mathcal{I}_v = \mathcal{Z}_K(\mathcal{D})$ .

Let  $t$  be an element of  $K^\times$  which is not a root of unity, and let  $H_t$  denote the preimage of  $\mathcal{H}_{\bar{t}}$  in  $K^\times$ , where  $\bar{t}$  denotes the image of  $t$  in  $\mathcal{K}_K$ . Assume that such a  $\mathcal{D}$  also satisfies condition (3) for this  $\bar{t}$ . We claim that the  $v$  mentioned above must be trivial on  $\mathbb{Q}$  hence also on  $k$ . For this, it suffices to show that  $2 \in H_t$ . Indeed, in this case we would find that  $\{2, 3, \dots\} \subset H_t$  since  $\mathcal{H}_{\bar{t}}$  is Milnor-closed, using Lemma 4.1. Thus, if  $v$  is nontrivial on  $\mathbb{Q}$ , and  $p$  is the prime for which  $v|_{\mathbb{Q}}$  is the  $p$ -adic valuation, we would have  $1 + p \in H_t \cap U_v^1$ , while  $\mathcal{D}^\perp = \mathcal{U}_v^1$ , hence the image of  $1 + p$  would be a nontrivial element of  $\mathcal{H}_{\bar{t}} \cap \mathcal{D}^\perp$ .

If  $t^n = -2$  for some nonzero integer  $n$ , then we have  $2 \in H_t$ , so we may assume this is not the case. By replacing  $t$  with  $t^n$  for some nonzero integer  $n$  (recall that this doesn't change  $H_t$ ), we may assume that  $1 + t$  and  $(2 + t)/t$  are also not roots of unity (note that our assumptions ensure that  $1 + t^n$  and  $(2 + t^n)/t^n$  are both nonzero for any nonzero integer  $n$ ).

To see this, consider the subfield  $\mathbb{Q}(t)$  generated by  $K$ . Assume first that there exists a complex embedding  $\sigma : \mathbb{Q}(t) \rightarrow \mathbb{C}$  such that  $|\sigma(t)| = 1$ , and write  $\sigma(t) = a + ib$ , with  $a, b$  real numbers. In this case, we show that  $1 + t$  and  $(2 + t)/t$  cannot be roots of unity, so that  $n = 1$  works. Indeed, if  $1 + t$  is a root of unity then

$$|\sigma(1 + t)|^2 = (1 + a)^2 + b^2 = 1 = a^2 + b^2,$$

which implies that  $1 + 2a = 0$  hence  $a = -1/2$  and  $b = \pm\sqrt{3}/2$ , so that  $t$  must be a root of unity. Similarly, if  $(2 + t)/t$  is a root of unity, then

$$|\sigma((2 + t)/t)|^2 = \frac{(2 + a)^2 + b^2}{a^2 + b^2} = (2 + a)^2 + b^2 = 1 = a^2 + b^2,$$

which implies that  $4 + 4a = 0$  hence  $a = -1$  and  $b = 0$ , so that again  $t$  would have to be a root of unity.

We can thus assume that for every complex embedding  $\sigma : \mathbb{Q}(t) \rightarrow \mathbb{C}$ , one has  $|\sigma(t)| \neq 1$ . Let  $\sigma$  be such an embedding. Replacing  $t$  with  $t^{-1}$  if needed, assume that  $|\sigma(t)| > 1$  hence  $|\sigma(t^n)| \rightarrow \infty$  as  $n \rightarrow \infty$  and thus  $1 + t^n$  is not a root of unity for all sufficiently large  $n$ . Replace  $t$  with such a  $t^n$  with  $n$  positive, and note that  $t^{n \cdot k}$  would have also worked for any positive  $k$ .

Now if  $(2+t)/t$  is a root of unity, then  $|2+\sigma(t)| = |\sigma(t)|$  which forces  $\sigma(t) = -1 + b \cdot i$  for some nonzero real  $b$ . If  $(2+t^2)/t^2$  is also a root of unity then  $\sigma(t^2)$  must have real part  $-1$  as well, hence  $-1 = 1 - b^2$  so that  $b = \pm\sqrt{2}$ . But in this case the real part of  $\sigma(t^3)$  is 5, so that  $(2+t^3)/t^3$  cannot be a root of unity.

Having made this replacement, we now use Lemma 4.1 repeatedly to find that  $1+t \in H_t$ ,  $2+t \in H_t$ ,  $(2+t)/t \in H_t$  and  $(2+t)/t-1 \in H_t$  as well. Thus,  $2 = t \cdot ((2+t)/t-1) \in H_t$ . In any case, we have obtained that  $2 \in H_t$ , which implies that  $v$  is trivial on  $k$  as noted above.

We deduce that  $\mathcal{K}_k$  is indeed contained in the intersection in question. On the other hand, if  $t$  is transcendental over  $k$ , then we may extend  $t$  to a transcendence base

$$\mathcal{B} = \{t\} \cup \mathcal{B}_0$$

for  $K|k$ . Put  $L := k(\mathcal{B}_0)$  and  $M = k(\mathcal{B}) = L(t)$ . Let  $v$  be an extension of the  $t$ -adic valuation on  $M$  to  $K$  and note that  $v$  is visible by Proposition 3.10 and our assumption on  $\text{trdeg}(K|k)$ . However, the image of  $t$  is not contained in  $\mathcal{U}_v = \mathcal{Z}_K(\mathcal{D}_v)^\perp$ . On the other hand, letting  $s \in \mathcal{B}_0$  be any element (which is nonempty by our assumption on  $\text{trdeg}(K|k)$ ), we have  $\mathcal{D}_v \in \mathcal{V}_{\bar{s}}$  where  $\bar{s}$  is the image of  $s$ , by arguing as in the proof of Theorem 4.6. Thus, the image of  $t$  is not contained in the intersection in question, and so the assertion of the proposition follows.  $\square$

*Proof of Theorem 5.4.* Let  $k$  denote the relative algebraic closure of the prime subfield of  $K$ , and  $l$  the relative algebraic closure of the prime subfield of  $L$ . The algebras  $\mathcal{K}_*(K)$  and  $\mathcal{K}_*(L)$  are isomorphic, while  $K$  has absolute transcendence degree  $\geq 5$ . Such an isomorphism induces an isomorphism of structures  $\mathcal{A}(K) \cong \mathcal{A}(L)$  by Fact 2.1, hence also an isomorphism  $\mathcal{G}_K \cong \mathcal{G}_L$  which is compatible with alternating pairs.

By Lemma 4.8, there exists a closed subspace  $\mathcal{D} \subset \mathcal{G}_{K|k} \subset \mathcal{G}_K$  such that  $\mathcal{Z}_K(\mathcal{D}) \neq \mathcal{D}$  and  $4 \leq \dim(\mathcal{Z}_K(\mathcal{D}))$ . Transferring  $\mathcal{D}$  across  $\mathcal{G}_K \cong \mathcal{G}_L$ , we obtain a closed subspace  $\mathcal{D}' \subset \mathcal{G}_L$  such that  $\mathcal{Z}_L(\mathcal{D}') \neq \mathcal{D}'$  and  $4 \leq \dim(\mathcal{Z}_L(\mathcal{D}'))$ . By Proposition 3.7,  $\mathcal{I} := \mathcal{Z}_L(\mathcal{D}')$  is valutive, and if  $v$  denotes the valuation associated to  $\mathcal{I}$ , it follows that  $\mathbb{Q} \otimes_{\mathbb{Z}} vL$  has dimension  $\geq 4$ . Since  $l$  the algebraic over a prime field, it follows that  $\mathbb{Q} \otimes_{\mathbb{Z}} vL/vl$  has dimension  $\geq 3$ , and thus  $3 \leq \text{trdeg}(L|l)$  by Abhyankar's inequality.

In any case, we may thus apply Proposition 5.5 to both  $K$  and  $L$ . This shows that the isomorphism  $\varphi_1 : \mathcal{K}_K \cong \mathcal{K}_L$  sends  $\mathcal{K}_k$  to  $\mathcal{K}_l$ , and thus  $\varphi_*$  descends to an isomorphism

$$\overline{\varphi}_* : \mathcal{K}_*(K|k) = \frac{\mathcal{K}_*(K)}{\langle \mathcal{K}_k \rangle} \cong \frac{\mathcal{K}_*(L)}{\langle \mathcal{K}_l \rangle} = \mathcal{K}_*(L|l).$$

By Theorem 5.3, we obtain an isomorphism of fields  $K^i \cong L^i$  and a rational number  $r \in \mathbb{Q}^\times$  satisfying the conditions of our theorem.  $\square$

**5.3. The finitely-generated case.** In the case where  $K|k$  is finitely generated, and  $k$  is perfect, we use the work of [4] to obtain a better result.

**Theorem 5.6.** *Let  $k$  be a perfect field and  $K$  a regular function field over  $k$  of transcendence degree  $\geq 2$ . Then  $K|k$  are determined, up-to isomorphism, from  $\mathcal{K}_*^M(K|k)$ . More precisely, if  $L|l$  is another regular function field over a perfect field of transcendence degree  $\geq 2$  and  $\varphi_* : \mathcal{K}_*^M(K|k) \cong \mathcal{K}_*^M(L|l)$  is an isomorphism, then there exists an isomorphism  $\Phi : K \cong L$  of fields satisfying  $\Phi(k) = l$  and some  $\varepsilon \in \{\pm 1\}$  such that  $\varepsilon \cdot \varphi_1$  is the isomorphism  $K^\times/k^\times \cong L^\times/l^\times$  induced by  $\Phi$ . If  $\text{trdeg}(K|k) \geq 3$ , then the assumption on  $\text{trdeg}(L|l)$  can be dropped.*

*Proof.* The isomorphism  $\varphi_1 : K^\times/k^\times \cong L^\times/l^\times$  is compatible with algebraic dependence by Theorem 4.6. Thus the claim follows from [4, Theorem 4]. In the case where  $\text{trdeg}(K|k) \geq 3$ , the fact that  $\text{trdeg}(L|l) \geq 2$  follows from Lemma 4.10.  $\square$

**Theorem 5.7.** *Let  $K$  be a finitely-generated field whose absolute transcendence degree is at least two. Then the isomorphism type of  $K$  is determined by  $K_*^M(K)$ . More precisely, suppose that  $L$  is any other finitely-generated field of absolute transcendence degree  $\geq 2$ ,  $\varphi_* : K_*^M(K) \cong K_*^M(L)$  is an isomorphism,  $k$  denotes the relative algebraic closure of the prime subfield of  $K$  and  $l$  the relative algebraic closure of the prime subfield of  $L$ . Then  $\varphi_*$  induces an isomorphism*

$$\bar{\varphi}_* : K_*^M(K|k) \cong K_*^M(L|l),$$

*and there exists an isomorphism of fields  $\Phi : K \cong L$  and some  $\varepsilon \in \{\pm 1\}$ , such that  $\varepsilon \cdot \bar{\varphi}_1 : K^\times/k^\times \cong L^\times/l^\times$  is the isomorphism induced by  $\Phi$ . If  $K$  has absolute transcendence degree  $\geq 4$ , then the assumption on the transcendence degree of  $L$  can be dropped.*

*Proof.* Use the characterization of  $\mathcal{K}_k$  in  $\mathcal{K}_K$  from Proposition 5.5 and the fact that the preimage of  $\mathcal{K}_k$  in  $K^\times$  is  $k^\times$  to observe that  $\varphi_1 : K^\times \cong L^\times$  sends  $k^\times$  to  $l^\times$ . Conclude by using Theorem 5.6. If  $\text{trdeg}(K|k) \geq 4$ , we may argue as in the proof of Theorem 5.4 to see that  $L$  has absolute transcendence degree  $\geq 2$ .  $\square$

## REFERENCES

1. Fedor Bogomolov and Yuri Tschinkel, *Commuting elements in Galois groups of function fields*, Motives, polylogarithms and Hodge theory. Part I: Motives and polylogarithms. Papers from the International Press conference, Irvine, CA, USA, June 1998, Somerville, MA: International Press, 2002, pp. 75–120 (English).
2. ———, *Milnor  $K_2$  and field homomorphisms*, Geometry, analysis, and algebraic geometry. Forty years of the Journal of Differential Geometry. Based on the 7th conference on geometry and topology, Cambridge, MA, USA, May 2008. In memory of Prof. C.-C. Hsiung, Somerville, MA: International Press, 2009, pp. 223–244 (English).
3. Fedor A. Bogomolov, *On two conjectures in birational algebraic geometry*, Algebraic geometry and analytic geometry. Proceedings of a conference, held in Tokyo, Japan, August 13–17, 1990, Tokyo etc.: Springer-Verlag, 1991, pp. 26–52 (English).
4. Anna Cadoret and Alena Pirutka, *Reconstructing function fields from Milnor  $K$ -theory*, Algebra Number Theory **15** (2021), no. 9, 2261–2288 (English).
5. Leonardo de Moura, Soonho Kong, Jeremy Avigad, Floris van Doorn, and Jakob von Raumer, *The lean theorem prover (system description)*, Automated Deduction - CADE-25 (Cham) (Amy P. Felty and Aart Middeldorp, eds.), Springer International Publishing, 2015, pp. 378–388.
6. Ido Efrat, *Construction of valuations from  $K$ -theory*, Math. Res. Lett. **6** (1999), no. 3–4, 335–343 (English).
7. Antonio José Engler and Jochen Koenigsmann, *Abelian subgroups of pro- $p$  Galois groups*, Trans. Am. Math. Soc. **350** (1998), no. 6, 2473–2485 (English).
8. David M. Evans and Ehud Hrushovski, *Projective planes in algebraically closed fields*, Proc. Lond. Math. Soc. (3) **62** (1991), no. 1, 1–24 (English).
9. ———, *The automorphism group of the combinatorial geometry of an algebraically closed field*, J. Lond. Math. Soc., II. Ser. **52** (1995), no. 2, 209–225 (English).
10. Jakub Gismatullin, *Combinatorial geometries of field extensions*, Bull. Lond. Math. Soc. **40** (2008), no. 5, 789–800 (English).
11. Jochen Koenigsmann, *From  $p$ -rigid elements to valuations (with a galois-characterization of  $p$ -adic fields)*, J. Reine Angew. Math. **465** (1995), 165–182 (English).
12. The mathlib Community, *The lean mathematical library*, Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (New York, NY, USA), CPP 2020, Association for Computing Machinery, 2020, p. 367–381.

13. Adam Topaz, *Commuting-liftable subgroups of Galois groups. II*, J. Reine Angew. Math. **730** (2017), 65–133 (English).
14. ———, *A Torelli theorem for higher-dimensional function fields*, 2017.
15. Adam Topaz, <https://github.com/adamtopaz/lean-acl-pairs>, 2022.

MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, 632 CENTRAL ACADEMIC BUILDING, EDMONTON AB T6G 2G1, CANADA

*Email address:* `topaz@ualberta.ca`

*URL:* <https://adamtopaz.com/>