1. A client taking risperidone presents with weight gain and hyperglycemia. What condition is likely occurring? **Metabolic Syndrome**
2. This medication carries a black box warning for suicidal ideation. **Montelukast**
3. Many immunosuppressants, corticosteroids, and biologic DMARDs share this major safety concern. **Risk for infection**
4. This antidepressant class is associated with anticholinergic effects. **Tricyclic antidepressant (TCA)**
5. A nurse should monitor this lab value in a client taking phenytoin. **Serum drug level**
6. This adverse effect is most closely associated with venlafaxine. **Hypertension**
7. Swollen and bleeding gums are associated with this adverse effect. **Gingival hyperplasia**
8. Lip smacking and tongue movements are characteristic of this adverse effect. **Tardive dyskinesia**
9. This inhaled corticosteroid reduces airway inflammation and requires the client to rinse the mouth after use. **Fluticasone**
10. This medication is used for acute wheezing and sudden asthma symptoms. **Albuterol**
11. This medication is used to stop an active seizure. **Lorazepam**
12. All antidepressants share this important teaching point regarding time to effectiveness. **"It may take several weeks to work"**
13. This mood stabilizer requires consistent sodium and fluid intake. **Lithium**
14. This is a common anticholinergic side effect of ipratropium. **Dry mouth**
15. These are common side effects of albuterol. **Tachycardia, tremors, nervousness**
16. Which vital sign should be carefully monitored in a client receiving lorazepam. **Respiratory rate**
17. This serious adverse effect may occur in a client taking valproic acid and presents with jaundice. **Hepatotoxicity**
18. This adverse effect is most closely associated with venlafaxine. **Hypertension**
19. A client with a seizure disorder should generally avoid this antidepressant. **Bupropion**
20. A client taking fluoxetine presents with tachycardia, fever, and mental status change. What life threatening condition could be occurring. **Serotonin syndrome**
21. This mood stabilizer requires consistent sodium and fluid intake. **Lithium**
22. Lip smacking and tongue movements are characteristics of this adverse effect. **Tardive dyskinesia**
23. Fever, muscle rigidity, and autonomic instability may indicate this rare antipsychotic emergency. **Neuroleptic malignant syndrome (NMS)**
24. A client taking an MAOI should avoid foods high in tyramine. Name some foods high in tyramine. **Aged cheese, cured meats, fermented foods, and wine**
25. This major adverse effect explains why a client receiving chemotherapy may develop fatigue, infection, and bleeding. **Myelosuppression**
26. This process is used to identify the most effective antibiotic for a bacterial infection. **Culture and sensitivity testing**
27. This class of medications used to treat autoimmune diseases differs from NSAIDs because it slows disease progression rather than simply relieving symptoms. **DMARDs**
28. This type of antineoplastic treatment acts on specific cellular receptors or proteins. **Biologic**