| | Hepatitis A | Hepatitis B | Hepatitis C | Hepatitis D | Hepatitis E |
|---|---|---|---|---|---|
| **Transmission** | Fecal-oral | Blood and sexual (semen, vaginal secretions, saliva, almost every body fluid) | Blood | Blood and sexual | Fecal-oral |
| **Isolation** | Standard | Standard | Standard | Standard | Standard |
| **Acute or Chronic?** | Acute | Both<br>Can also be carrier | Both | Both | Acute; chronic if immunosuppressed |
| **Risk Factors** | Developing countries (contaminated water or shellfish)<br>Day care/preschool<br>MSM | MSM/multiple sex partners<br>IVDU<br>HD<br>Healthcare workers<br>Public safety workers<br>Blood transfusion<br>Mom to baby | IVDU<br>MSM/multiple sex partners<br>HIV<br>Healthcare workers<br>Mom to baby | MSM/multiple sex partners<br>IVDU<br>HD<br>Healthcare workers<br>Public safety workers<br>Blood transfusion<br>Mom to baby | Developing countries<br>Poor hygiene<br>Pregnant women may be affected severely |
| **Dx** | Anti-HAV IgM = acute infection<br>HAV IgG = past infection | HBsAg x 6 months or longer = chronic<br>Anti-HBs = immunity | Anti-HCV in blood + positive viral load = active infection<br>Anti-HCV in blood but no viral load = past infection | Need HBV in past<br>Anti-HDV | |
| **Prevent** | Hand washing<br>Vaccination | Vaccination<br>Don't share personal items<br>Condoms | Reduce risk<br>(No vaccine) | HBV vaccination | Hand washing |
| **Extra Things** | S/S only for >6yo | Increased risk for liver CA, cirrhosis | Jaundice uncommon<br>80% develop chronic hepatitis<br>Increased risk for liver CA, cirrhosis | Can worsen hepatitis B | |
| **S/S (Acute)** | *Prodromal (pre-icterus) period* - abrupt/insidious; malaise, myalgias, arthralgias, anorexia, N/V/D/C, increased AST/ALT (1-2 weeks)<br>*Icterus phase* - increased bilirubin, RUQ tenderness, mild weight loss, spider angiomas (2-12 weeks)<br>*Recovery phase* - increased well-being, return of appetite, no jaundice (1-4 weeks) | | | | |