	Ischemia	Necrosis	Symptoms
Peripheral Vascular Disease (PVD)	Intermittent claudication	Critical limb ischemia and gangrene	Intermittent claudication; increased cap refill, skin cold, weak or absent pulses in extremity; ruddy when dependent and pale when elevated
Brain	TIA	CVA	BEFAST
Heart	Angina	MI	Chest pain, SOB, diaphoresis, nausea, anxiety

Symptoms worsen

Risk Factors

Atherosclerosis & Thrombi	Emboli
Hypertension Hyperlipidemia Smoking Increased age Genetics • Diabetes mellitus • Alcohol intake • Stress • Menopause	<u>Virchow's triad</u>: Venous stasis - immobility, obesity, crossing legs, dehydration Hypercoagulability - increased age, menopause, PO contraceptives, smoking Vessel damage - endothelial dysfunction • (hypertension), trauma

Chest Pain Classification

Angina or MI?	Type	EKG Changes	Troponin	S/S
Angina	Stable/Classic	None	Negative	Chest pain with exertion that is relieved with rest
	Unstable	ST depression T changes	Negative	Prolonged chest pain with rest OR more severe than previous
MI	NSTEMI	ST depression T inversion	Positive	Chest pain that can be treated with nitroglycerin
	STEMI	ST elevation	Positive	Chest pain that may not be relieved by nitroglycerin