Exam Three Practice Quiz

Adam Weber — 2nd Attempt

Due Date	Time Limit	Number of Questions
No due date	**No time limit**	**36**

This quiz does not count toward the final grade



36
Out of 36 points

17:34
Time for this attempt

This assessment has unlimited attempts. Take Now

Attempt History

Results	Points	Score	(Highest score is kept)
Attempt 1	16.517 of 36	45.88%	
Attempt 2	**36 of 36**	**100%**	**(Highest score)**

Your Answers:

1 1 / 1 point Multiple Choice

A client with suspected peripheral vascular disease requires a neck vessel assessment. Which instruction should the nurse provide before palpating the carotid arteries?

- Hold the breath briefly during palpation
- Turn the head fully toward the side being examined
- Perform the palpation immediately after ambulation
- ✓ Avoid pressing on both sides of the neck at the same time

2 1 / 1 point Multiple Choice

A client has neck vessels that are difficult to visualize. Which assessment strategy should the nurse use to obtain the best jugular venous data?

- Compress both jugular veins briefly to make pulsations easier to see
- Palpate the external jugular vein while the client is sitting upright
- Assess jugular venous pressure after the client performs several deep breaths
- ✓ Observe the right internal jugular vein with the client positioned at 30 to 45 degrees

Feedback

Based on your answer

The right internal jugular vein is preferred because it is in direct line with the superior vena cava and best reflects right-sided hemodynamics. Positioning at a semi-Fowler angle improves visualization of venous pulsations. Bilateral compression is unsafe, and deep breathing or relying on the external jugular vein alone may reduce accuracy or alter the reading.

3 1 / 1 point Multiple Choice

A nurse is assessing the feet of a client with a history of diabetes mellitus and peripheral neuropathy. Which assessment finding is most consistent with a neuropathic ulcer?

- An irregularly shaped ulcer near the ankle with brown discoloration of the surrounding skin
- A painful, shallow ulcer near the medial ankle with surrounding edema
- A deep, painful ulcer on the toes with pale skin and diminished pedal pulses

✓ A painless ulcer on the plantar surface of the foot surrounded by thickened callused skin

4 1 / 1 point Multiple Answer

Which of the following are ***non-modifiable*** risk factors of coronary artery disease?

- [ ] High cholesterol
- [ ] Smoking
- [x] Male gender
- [x] Greater than 65 years of age
- [x] Family history of heart disease
- [ ] Hypertension

5 1 / 1 point Multiple Choice

A client with chronic lung disease reports increasing ankle swelling, abdominal fullness, and a rapid weight gain over 3 days. Which finding is most consistent with right-sided heart failure?

- ○ Narrowed pulse pressure
- ○ Crackles heard best at the lung bases
- ○ Bounding peripheral pulses
- ✓ ● Jugular venous distention

Feedback

Based on your answer

Right-sided heart failure causes systemic venous congestion, which commonly produces jugular venous distention, dependent edema, hepatomegaly, ascites, and weight gain. Crackles

are more typical of pulmonary congestion from left-sided failure. Bounding pulses and narrowed pulse pressure are not the classic pattern for right-sided failure.

6 1 / 1 point Multiple Choice

A 15-year-old client reports breast tenderness and a small amount of breast enlargement. Which nursing response is best?

- Tell the client that breast enlargement in males is always abnormal
- Teach that this finding requires immediate surgical evaluation
- Advise stopping all physical activity until the breasts return to baseline
- ✓ Explain that this can occur with pubertal changes and is usually temporary

7 1 / 1 point Multiple Choice

A client has an intact area of skin over the sacrum with localized non-blanchable erythema after prolonged immobility. Which lesion classification should the nurse document?

- Stage 4 Pressure Injury
- Unstageable Pressure injury
- Stage 2 Pressure Injury
- ✓ Stage 1 Pressure Injury

Feedback

Based on your answer

Intact skin with localized non-blanchable erythema is consistent with a stage 1 pressure injury. Stage 2 involves partial-thickness skin loss or a blister, stage 3 has full-thickness skin loss with visible adipose, and unstageable injuries are obscured by slough or eschar. The nurse should use wound staging based on the depth of tissue loss.

8 1 / 1 point Multiple Choice

A client asks for teaching about testicular self-examination after reporting concern about testicular cancer risk. Which instruction should the nurse include?

- ○ Inspect the penis first and then palpate the prostate
- ○ Repeat the exam only if pain develops in the scrotum
- ✓ ● Perform the exam after a warm bath or shower when the scrotum is relaxed
- ○ Use both hands to squeeze the testicle firmly to assess for tenderness.

Feedback

Based on your answer

A warm bath or shower helps relax the scrotal tissue, making palpation easier and more effective for detecting changes. The exam should focus on palpating each testis gently and systematically. Inspecting the penis and palpating the prostate are not part of testicular self-examination. Repeating the exam only when pain occurs can delay recognition of abnormal changes, and squeezing firmly is inappropriate because the exam should be gentle.

9 1 / 1 point True or False

A nurse should position the client supine with the arm on the side being examined raised above the head when palpating breast tissue.

- ✓ ● True
- ○ False

10 1 / 1 point Multiple Choice

A client asks how to lower breast cancer risk during a wellness visit. Which teaching should the nurse include?

- ○ Limit intentional exercise to 20 minutes on most days
- ○ Use breast self-examination as the primary screening test
- ✓ ● Avoid alcohol intake and maintain a healthy adult weight

Stop eating dairy products completely to prevent cancer

Feedback

Based on your answer

Regular intentional exercise, avoiding alcohol, and preventing excessive weight gain are included in risk-reduction teaching for breast cancer. These measures support healthier modifiable risk management. The exercise option is incorrect because the recommended amount is much higher than 20 minutes on most days. Breast self-examination is not the primary screening test, and complete avoidance of dairy is not a standard evidence-based prevention strategy.

11 1 / 1 point Multiple Choice

During a cardiac assessment, the nurse recognizes that the first heart sound (S1) occurs during which event of the cardiac cycle?

Closure of the semilunar valves

✓ Closure of the atrioventricular valves

Isometric contraction

Beginning of diastole

12 1 / 1 point Multiple Answer

A nurse is teaching a client about factors that increase the risk for cervical cancer. Which findings should the nurse include? ***Select all that apply.***

Routine screening with Pap testing every 3 years

✓ Family history of cervical cancer

✓ Previous human papillomavirus infection

✓ Diet low in fruits and vegetables

Feedback

Previous HPV infection is a recognized risk factor for cervical cancer, and diet low in fruits and vegetables is also associated with increased risk. A family history of cervical cancer both increase risk as well. Routine Pap testing is a protective screening measure, not a risk factor.

13 1 / 1 point Multiple Choice

A nurse is performing a breast assessment on a female client. Which area should the nurse palpate to assess the tail of Spence?

- ✓ The upper outer quadrant of the breast extending toward the axilla
- The lower inner quadrant of the breast near the sternum
- The lower outer quadrant of the breast near the inframammary fold
- The upper inner quadrant of the breast near the clavicle

14 1 / 1 point Multiple Choice

A client has a small, fluid-filled lesion on the forearm after poison ivy exposure. Which description best fits the nurse's assessment finding?

- Macule
- Tumor
- Papule
- ✓ Vesicle

Feedback

Based on your answer

A vesicle is a circumscribed elevated lesion containing serous fluid and is less than 0.5 cm, which is consistent with poison ivy-type lesions. A macule is flat, a papule is solid and elevated,

and a tumor is a larger deeper mass. The nurse should match the lesion's fluid content and size to the correct morphology.

15 1 / 1 point Ordering

A nurse is preparing to auscultate a client's precordium during a cardiovascular assessment. Place the cardiac auscultation sites in the correct sequence.

- ✓ Aortic
- ✓ Pulmonic
- ✓ Erb's
- ✓ Tricuspid
- ✓ Mitral

16 1 / 1 point Multiple Choice

A client with a known abdominal aortic aneurysm suddenly reports sharp back pain and dizziness. Which nursing action is the priority?

- ○ Reassess abdominal girth for comparison with prior measurements
- ○ Place the client in a sitting position to ease discomfort
- ✓ ● Activate emergency response and prepare for immediate escalation of care
- ○ Encourage oral fluids to support circulation

17 1 / 1 point Multiple Choice

A client with a sibling diagnosed with breast cancer asks which additional history findings raise concern. Which finding should the nurse identify as a risk factor?

- Menopause before age 40
- ✓ Radiation therapy to the chest during childhood
- Recent increase in water intake
- Frequent use of over-the-counter antacids

Feedback

Based on your answer

Radiation therapy to the chest or breast during childhood or young adulthood is a recognized breast cancer risk factor. The family history already increases concern, making this additional exposure more significant. Early menopause is not the risk pattern described here, and antacids or water intake are not relevant risk factors.

18 1 / 1 point Multiple Choice

A client using oral contraceptives reports new unilateral leg swelling and shortness of breath. Which action should the nurse take first?

- Ask whether the client has recently changed oral contraceptive brands
- Recommend an over-the-counter analgesic and rest
- ✓ Instruct the client to seek urgent evaluation for possible thromboembolism
- Teach the client to continue the medication and monitor symptoms for 1 week

Feedback

Based on your answer

Oral contraceptives are associated with thromboembolic events, and unilateral swelling with dyspnea is concerning for venous thromboembolism or pulmonary embolism. This is a time-sensitive safety issue requiring immediate evaluation. Delaying care for observation or symptom management is inappropriate.

19 1 / 1 point Multiple Choice

An older adult client presents with cramping-type leg pain when walking, which is relieved by rest. The client also has cool, pale feet and capillary refill in the toes of 4 to 6 seconds. What would the nurse suspect?

- Musculoskeletal weakness
- ✓ Arterial insufficiency
- Venous insufficiency
- Diabetic neuropathy

20 1 / 1 point Multiple Choice

A newborn assessment reveals the urethral opening located on the top of the glans. Which interpretation should the nurse document?

- Hypospadias
- ✓ Epispadias
- Paraphimosis
- Phimosis

21 1 / 1 point Matching

A nurse is performing an abdominal assessment on clients with suspected gastrointestinal conditions. Match each assessment technique with the appropriate finding or condition.

Obturator Sign	✓ Right lower quadrant pain with internal rotation of the flexed right hip, suggesting appendicitis
Blumberg Sign	✓ Abdominal pain upon release of pressure, suggesting peritonitis or appendicitis

Psoas Sign	✓ Right lower quadrant pain when the right hip is extended, suggesting appendicitis
Murphy Sign	✓ Palpate under the liver border at the right costal margin, while the client inhales deeply, suggesting cholecystitis

22 1 / 1 point Multiple Answer

A nurse is assessing a client abdominal aortic aneurysm (AAA). Which of the following findings would suggest a ruptured aneurysm? *Select all that apply.*

- ✓ ☑ Unresponsiveness
- ☐ Decreased appetite
- ✓ ☑ Clammy skin
- ✓ ☑ Sudden pain in the abdomen and back
- ✓ ☑ Tachycardia

23 1 / 1 point Multiple Choice

A client with hemorrhoids says, "I want to prevent this from getting worse." Which teaching should the nurse prioritize?

- ✓ ◉ Increase fiber, hydrate well, and avoid straining during bowel movements
- ○ Use stronger soaps to prevent skin breakdown around the anus
- ○ Hold bowel movements until the end of the day to reduce urgency

Spend more time sitting on the toilet to fully empty the bowel

Feedback

Based on your answer

The most effective prevention strategy is to reduce constipation and straining, which lowers pressure in the hemorrhoidal veins. Adequate fluids and fiber support stool softening and regular elimination. Stronger soaps can irritate inflamed tissue. Prolonged toilet sitting increases pressure and worsens symptoms. Delaying bowel movements can increase stool retention and constipation.

24 1 / 1 point Multiple Answer

A nurse is providing education to a group of adults about preventing skin cancer. Which recommendations should the nurse include? ***Select all that apply.***

- [ ] Use tanning oils to prevent sunburn
- [x] Educate the client to increase niacin in their diet
- [ ] Apply sunscreen only when temperatures exceed 80 degrees farenheit
- [x] Avoid tanning beds and sunlamps
- [x] Wear protective clothing when outdoors
- [x] Seek shade during peak UV exposure

25 1 / 1 point Multiple Choice

A client with chest pain says, “My heart must be acting up.” Which nurse response is the best example of therapeutic communication while gathering more data?

- “That is probably just stress from work.”
- ✓ “Tell me more about what changes you have noticed over the last few months.”
- “You should not worry until the test results are back.”

"Chest pain is usually caused by indigestion."

26 1 / 1 point Multiple Choice

The nurse is reviewing symptom data for a client with possible advanced prostate cancer. Which finding should the nurse identify as most concerning for disease progression?

- ✓ Unexplained weight loss
- Blood in the urine or semen
- Trouble urinating at night
- Decreased force in the urine stream

Feedback

Based on your answer

Unexplained weight loss can indicate more advanced disease and requires prompt follow-up. The other findings are important but are more often associated with local urinary tract or prostate involvement and are not as specific for advanced progression. The nurse should still report all abnormal symptoms, but weight loss is the most concerning cue in this group.

27 1 / 1 point Multiple Choice

A nurse is providing discharge teaching to a client diagnosed with gastroesophageal reflux disease (GERD). Which of the following statements by the client indicates a **need for further teaching**?

- "I will wait two hours after eating before lying down."
- ✓ "I can have a glass of wine with dinner to help me relax."
- "I'll try to eat smaller meals throughout the day."
- "I will avoid drinking soda."

28 1 / 1 point Matching

A nurse is performing breast assessments on several clients. Match each assessment finding with the most appropriate condition or interpretation.

Peau D' Orange	✓ Orange peel appearance of the breast
Breast malignancy	✓ Nipple retraction
Fibrocystic Breast Disease	✓ Painful, lumpy rope like cysts improving after menses
Paget's Disease	✓ Redness and flaking of the nipple
Mastitis	✓ Periductal inflammation of breast tissue resulting in redness and pain

29 1 / 1 point Matching

A nurse is assessing bowel sounds in clients with various gastrointestinal conditions. Match each type of bowel sound with the condition most commonly associated with the finding.

Hypoactive	✓ Postoperative ileus, intestinal obstruction
Hyperactive	✓ Gastroenteritis, diarrhea
Absent	✓ Ruptured appendix, paralytic ileus, advanced bowel obstruction

30 1 / 1 point Multiple Choice

The nurse is documenting a normal nail assessment. Which charting entry is most appropriate?

- Fingernails pale, spoon-shaped, and tender to touch
- Fingernails yellow, thick, and painful with delayed refill
- Fingernails chipped, uneven, and separated from the nail bed
- ✓ Fingernails pink, smooth, hard, immobile; capillary refill <2 seconds; cuticles intact

31 1 / 1 point Multiple Choice

A nurse is performing a genitourinary assessment on an older adult male client. Which assessment finding should the nurse recognize as an expected age-related change?

- Increased firmness and enlargement of the testes
- Presence of painless, hard nodules on the testes
- ✓ Decreased scrotal muscle tone with a more pendulous appearance
- Increased pubic hair growth and distribution

32 1 / 1 point Multiple Answer

A nurse is assessing a hospitalized client for risk factors associated with pressure injury development. Which findings increase the client's risk? ***Select all that apply.***

- ✓ Poor nutritional status
- ✓ Limited mobility
- Adequate nutritional intake
- ✓ Urinary incontinence
- ✓ Decreased sensory perception

- [ ] Indepent repositioning

33 1 / 1 point Multiple Choice

A nurse is assessing a client with chronic venous insufficiency. Which finding is most consistent with this condition?

- Cool, pale lower extremities with diminished pulses
- Thick, shiny skin with absent hair growth
- ✓ Brown discoloration around the ankles with bilateral lower leg edema
- Pallor of the foot when elevated with delayed capillary refill

34 1 / 1 point Multiple Choice

A client with a history of peptic ulcer disease wants to reduce recurrence risk and asks which behavior change is the best starting point. Which response should the nurse give?

- Replace all meals with bland foods
- ✓ Stop smoking and avoid chewing tobacco
- Increase intake of caffeine to improve digestion
- Lie down after meals to reduce acid exposure

35 1 / 1 point Multiple Choice

A client with chronic constipation is using laxatives frequently and reports anxiety about bowel movements. Which nursing concern is most appropriate to identify?

- Acute confusion related to stool retention
- ✓ Poor health behavior related to inappropriate laxative use
- Impaired gas exchange related to constipation

Risk for dehydration related to normal bowel elimination

36 1 / 1 point Multiple Choice

A client in an intensive care unit has a central venous line, is receiving broad-spectrum antibiotics, and has a family member with a recent MRSA infection. Which factor should the nurse recognize as the strongest contributor to MRSA risk?

Family member visiting the unit

The client is receiving a liquid diet

Reporting of dry skin on the elbows

✓ Presence of an invasive medical device