Results

Adam Weber — 1st Attempt

100%

31

Out of 31 points

12:29

Time for this attempt

This assessment has unlimited attempts. Take Now

Attempt History

Results	Points	Score	(Highest score is kept)
Attempt 1	31 of 31	100%	(Highest score)

Your Answers:

1 1 / 1 point Multiple Choice

A nurse is assessing a client who reports recurring headaches. Which of the following assessment findings would indicate that the client is experiencing a migraine headache?

- ○ Severe, unilateral pain around one eye accompanied by tearing and nasal congestion.
- ○ Facial pressure and discomfort that worsens when bending forward, accompanied by nasal congestion.
- ○ Bilateral head pain described as a tight band around the forehead and temples.
- ✓ ● Throbbing pain on one side of the head accompanied by nausea and sensitivity to light.

2 1 / 1 point Multiple Choice

A nurse is teaching a client about the appearance of an acute gout flare. Which description should the nurse include?

- A foot with no arch and generalized surface swelling
- Hard nodules over the distal finger joints without warmth
- ✓ A tender, swollen, red, and hot big toe joint
- A painless, fluid-filled cyst on the dorsum of the wrist

Feedback

Based on your answer

Acute gouty arthritis classically affects the first metatarsophalangeal joint and produces tenderness, redness, heat, and swelling.

3 1 / 1 point Multiple Choice

The nurse is reviewing risk factors with a client who smokes and has seasonal allergies. Which client statement shows understanding of teaching to lower sinusitis risk?

- "I can keep using smoke exposure if I drink more water."
- "I only need treatment when facial pain becomes severe."
- ✓ "I should avoid tobacco smoke and manage allergy triggers."
- "Bacterial sinusitis is prevented by skipping vaccines."

4 1 / 1 point Multiple Choice

A nurse is performing a neurological assessment on a client. The nurse observes that both pupils are equal in size, round, and constrict when exposed to light. When the client shifts focus from a distant object to a nearby object, both pupils constrict. How should the nurse interpret these findings?

- The client demonstrates an abnormal pupillary response that requires further neurological assessment.
- The client exhibits impaired accommodation despite having an intact pupillary light reflex.
- The client demonstrates an abnormal response of the optic and oculomotor nerves.
- ✓ The client demonstrates normal pupillary responses to light and accommodation.

5 1 / 1 point Multiple Choice

The nurse is teaching a client with chronic obstructive pulmonary disease how to breathe during episodes of dyspnea. Which instruction should the nurse include?

- ✓ Exhale slowly through pursed lips and prolong the out-breath
- Hold the breath after each inhalation to improve oxygen exchange
- Take rapid shallow breaths while leaning backward in bed
- Inhale quickly through the mouth and exhale slowly through pursed lips

6 1 / 1 point Multiple Choice

The nurse prepares to assess cranial nerve 5 (V). Which technique is most appropriate?

- Ask the client to identify the sound of a vibrating tuning fork on the forehead and cheek
- Ask the client to follow a penlight in all directions
- ✓ Lightly touch the forehead, cheeks, and chin while asking about sensation
- Place sugar on the anterior tongue and ask for identification

7 1 / 1 point Matching

A nurse is performing an otoscopic examination on several clients. Match each tympanic membrane assessment finding with the most likely condition or disorder.

Serous otitis media	✓ Yellowish, bulging membrane with fluid bubbles
Skull trauma	✓ Bluish or dark red color behind the tympanic membrane
Normal tympanic membrane	✓ Pearly gray, translucent tympanic membrane with visible landmarks and a cone of light.
Acute otitis media	✓ Red, bulging eardrum with absent light reflex

8 1 / 1 point Multiple Choice

A nurse is assessing a client who reports a headache, facial pressure, and halitosis (bad breath) . Which action should the nurse implement ***first*** during the sinus portion of the exam?

- ✓ ● Palpate the maxillary and frontal areas gently for tenderness
- ○ Administer antibiotic use for a sinus infection
- ○ Auscultate over the cheekbones for abnormal sounds
- ○ Inspect the tonsils for exudate using a tongue depressor

9 1 / 1 point Multiple Choice

The nurse notes that a client cannot puff out the cheeks and has asymmetry when smiling. Which cranial nerve is most likely impaired?

- ✓ ● Cranial nerve VII (7)
- ○ Cranial nerve XII (12)
- ○ Cranial nerve V (5)

Cranial nerve IX (9)

10 1 / 1 point Multiple Choice

During an eye assessment, a client states that distant signs are hard to read, but close-up reading is still clear. Which interpretation should the nurse make?

Diplopia related to extraocular muscle weakness

Acute glaucoma with increased intraocular pressure

✓ Myopia causing difficulty with far vision

Presbyopia causing loss of peripheral vision

11 1 / 1 point Multiple Choice

During a head-to-toe neurologic examination, the nurse needs to assess cranial nerve II (2). Which action should the nurse select?

Check sound localization with a whispered voice test

Ask the client to move the eyes through the six cardinal fields of gaze

✓ Test peripheral vision while the client focuses straight ahead

Ask the client to protrude the tongue and move it side to side

12 1 / 1 point Multiple Choice

Which otoscopic finding should the nurse identify as expected in a healthy ear assessment?

✓ Pearly gray tympanic membrane with a cone of light

White dome-shaped lesion in the ear canal

Bulging membrane with no visible landmarks

Erythematous canal with drainage

13 1 / 1 point Multiple Choice

The nurse is reviewing findings after an eye assessment. Which result best supports a normal six cardinal fields of gaze examination?

- The client maintains focus on the target but demonstrates jerky eye movements in multiple directions.
- ✓ Smooth, symmetric eye movements with no nystagmus.
- The client reads the Snellen chart with missed letters.
- One pupil constricts more slowly than the other.

14 1 / 1 point Multiple Answer

A nurse is teaching a community group about cataracts. Which findings should the nurse include as risk factors for cataracts? ***Select all that apply.***

- Maintaining blood pressure within the recommended range
- ✓ Excessive exposure to sunlight
- Low-fat diet
- ✓ Smoking tobacco
- ✓ Type 2 diabetes mellitus

15 1 / 1 point Multiple Choice

The nurse is reviewing risk factors with a client who reports frequent marijuana use, heavy alcohol intake, and poor diet with few fruits and vegetables. Which finding should the nurse prioritize as the strongest contributor to oral cancer risk?

- Use of a manual toothbrush instead of an electric toothbrush
- Preference for soft foods over crunchy foods

✓ ● Frequent heavy alcohol intake combined with tobacco-related behaviors

○ Drinking water only between meals

16 1 / 1 point Multiple Choice

A client reports elbow stiffness after an arm injury. During the musculoskeletal assessment, which action should the nurse use to evaluate elbow range of motion?

✓ ● Ask the client to flex and extend the forearm while the nurse compares the movement with the opposite arm

○ Measure grip strength against resistance and infer elbow mobility from the result

○ Palpate the elbow joint for crepitus and document the finding as normal if no pain occurs

○ Observe the client walk across the room and note arm swing symmetry

Feedback

Based on your answer

Elbow range of motion is assessed by observing and guiding the client through flexion and extension, then comparing bilateral findings.

17 1 / 1 point Multiple Answer

A client is admitted for treatment of hypothyroidism. What will the nurse expect to assess in this client? **Select all that apply.**

☐ Decreased ability to taste

✓ ☑ Dry, coarse, and sparse hair

✓ ☑ Edema around the eyes

✓ ☑ Puffy, dull face

☐ Exopthalmus

18 1 / 1 point Multiple Choice

A family member reports several changes in an older adult client's behavior. Which statement should the nurse recognize as ***most*** concerning for Alzheimer's disease?

- ✓ "My mother can no longer manage her monthly bills, even though she handled them independently for years."
- "My mother has started taking afternoon naps more frequently."
- "My mother occasionally forgets the names of people she rarely sees."
- "My mother sometimes needs extra time to remember where she placed her belongings."

19 1 / 1 point Multiple Choice

A community nurse is reviewing stroke-prevention teaching with a client who has multiple risk factors. Which finding suggests the client needs further teaching?

- "I will ask about my blood pressure checks at each visit."
- "I will limit foods that are high in saturated fat."
- "I plan to eat more fruits, vegetables, and whole grains."
- ✓ "I will smoke less and quit later when things settle down."

20 1 / 1 point Multiple Choice

A nurse is assessing hearing in an older adult client. Which assessment finding is most consistent with presbycusis?

- Ear pain and decreased hearing following an upper respiratory infection.
- Sudden hearing loss in one ear accompanied by dizziness.
- Intermittent ringing in the ears accompanied by a feeling of fullness.
- ✓ Gradual difficulty hearing high-pitched sounds

21 1 / 1 point Multiple Choice

A nurse palpates a client's cervical lymph nodes and identifies several small, soft, movable, nontender nodes. Which documentation is **most appropriate**?

- ✓ "Cervical lymph nodes palpable, soft, movable, and nontender."
- "Cervical lymph nodes enlarged, indicating an infectious process."
- "Lymphatic system functioning normally with no abnormalities."
- "Cervical lymph nodes abnormal; further evaluation required."

22 1 / 1 point Matching

A nurse is performing respiratory auscultation. Match each normal breath sound with the location where it is typically heard.

Bronchovesicular	✓ Located in the major bronchi
Bronchial	✓ Over the trachea and larynx
Vesicular	✓ Located over peripheral lung fields

23 1 / 1 point Multiple Answer

A nurse is assessing a client with Kussmaul respirations. Which findings should the nurse expect? ***Select all that apply.***

- ☐ Shallow respirations
- ✓ ☑ Labored breathing
- ☐ Retention of air
- ☐ Periods of apnea

- [x] Rapid respiratory rate
- [x] Deep respirations

24 1 / 1 point Multiple Choice

A nurse is assessing a client who reports experiencing recurrent headaches over the past several months. Which nursing action is **most appropriate** to help identify potential contributing factors?

- ( ) Advise the client to eliminate all caffeinated beverages immediately to reduce headache frequency.
- ( ) Recommend that the client avoid all physical activity until the headaches completely resolve.
- ( ) Instruct the client to take over-the-counter pain medications daily to prevent headaches.
- (x) Encourage the client to maintain a headache diary documenting the frequency, duration, and potential triggers of headaches.

25 1 / 1 point Matching

A nurse is reviewing common eye disorders with a group of nursing students. Match each eye disorder with its characteristic assessment finding

Glaucoma	Loss of peripheral vision
Cataracts	Clouding of the lens that causes blurred vision
Macular degeneration	Gradual loss of central vision
Presbyopia	Age-related difficulty focusing on nearby objects, often re-quiring reading glasses.

26 1 / 1 point Multiple Choice

A community health nurse is educating a group of adults about strategies to reduce the risk of traumatic brain injuries. Which statement by a participant indicates understanding of the teaching?

- ✓ "Wearing a properly fitted helmet when riding a bicycle can reduce my risk of a serious head injury."
- "Removing loose rugs and improving lighting at home is only necessary for older adults."
- "Wearing a seat belt is only necessary when traveling at high speeds."
- "I should focus on preventing falls only if I have a history of balance problems."

27 1 / 1 point Multiple Choice

The nurse is screening cranial nerve VIII (8) during admission. Which finding supports intact function?

- The client turns the head against resistance without difficulty
- ✓ The client repeats whispered words heard from 3 feet away
- The client identifies sweet and salty substances on the tongue
- The client lifts the uvula symmetrically when saying “ah”

Feedback

Based on your answer

Hearing of whispered words is a standard way to screen vestibulocochlear nerve function. Taste recognition on the tongue relates to cranial nerves VII, IX, and X, uvular elevation reflects cranial nerves IX and X, and resisted head turning assesses cranial nerve XI.

28 1 / 1 point Multiple Choice

A client with chronic obstructive pulmonary disease is leaning forward with hands on the knees and is using accessory muscles to breathe. Which finding should the nurse interpret as a compensatory breathing pattern?

✓ Tripod position with accessory muscle use

Symmetric chest expansion with quiet respirations

No nasal flaring during inspiration

Respiratory rate of 16/min with regular depth

29 1 / 1 point Multiple Choice

During an assessment, a client opens the eyes to speech, answers questions appropriately, and remains awake. Which interpretation should the nurse make?

Obtundation

✓ Alertness with intact arousal

Stupor

Lethargy

30 1 / 1 point Multiple Choice

A nurse is assessing an older adult client who reports difficulty maintaining balance while walking. Which age-related change may contribute to the client's increased risk for falls?

✓ Decreased sense of positioning

Improved coordination

Increased joint flexibility

Increased bone density

31 1 / 1 point Multiple Answer

A client asks the nurse about osteoporosis risk after a recent health screening. Which factors increase the client's risk for osteoporosis or fracture? ***Select all that apply.***

✓ Use of prednisone for long-term inflammatory disease management

- [x] Celiac disease with poor nutrient absorption

Feedback

Low body mass index is a modifiable risk factor for osteoporosis and fracture because lower body mass is associated with reduced bone reserve. Long-term glucocorticoid use, such as prednisone, is a fixed risk factor because it accelerates bone loss. Alcohol intake greater than 2 drinks per day and medical conditions that impair nutrient absorption, such as celiac disease, also increase fracture risk. Regular weight-bearing exercise and adequate calcium and vitamin D intake are protective rather than risk increasing.

- [ ] Regular weight-bearing exercise four times each week
- [ ] High calcium intake with adequate vitamin D supplementation
- [x] Low body mass index of 19 kg/m^2
- [x] Daily alcohol intake of three drinks