

1. This major chemotherapy adverse effect results in decreased RBCs, WBC, and low platelets. - **Myelosuppression**
2. This antineoplastic medication is associated with hemorrhagic cystitis. - **Cyclophosphamide**
3. These vaccines may be contraindicated in patients receiving immunosuppressive therapy. - **live vaccines**
4. This corticosteroid may increase infection risk and should not be stopped abruptly. - **Prednisone**
5. This medication is associated with peripheral neuropathy. **Vincristine**
6. These are the required elements of every medication order. **Name, dose, route, frequency**
7. This medication treats some forms of bacteria as well as protozoa. **Metronidazole**
8. Medications administered by this route are most commonly affected by first-pass metabolism. **Oral route**
9. This medication causes cardiotoxicity and may turn body fluids red-orange. **Doxorubicin**
10. This second generation NSAID has lower risk of GI bleeding compared with first-generation NSAIDs. **Celecoxib**
11. This biologic DMARD and TNF blocker increases the risk of serious infections and requires TB screening. **Adalimumab**
12. This medication treats tuberculosis and may cause orange-red body fluid. **Rifampin**
13. These are three major medication classes commonly used to treat cancer. **Chemotherapy, hormonal agents, and biologics**
14. This DMARD is associated with visual changes. **Hydroxychloroquine**
15. This branch of pharmacology describes how the body absorbs, distributes, metabolizes, and excretes a medication. **Pharmacokinetics**
16. This medication is commonly used to treat MRSA and is associated with an infusion reaction. **Vancomycin**
17. This medication should be avoided in children with viral illnesses because of the risk of Reye Syndrome. **Aspirin**
18. This phenomenon occurs when a medication is metabolized by the liver before reaching systemic circulation, reducing the amount of active drug available. **First-pass effect**
19. This medication is commonly used after organ transplantation and is associated with nephrotoxicity. **Cyclosporine**
20. This is the major risk associated with many immunosuppressant medications. **Risk for infection**
21. This first-pass NSAID is commonly associated with GI bleeding. **Ibuprofen**
22. This fluoroquinolone is associated with tendon rupture. **Ciprofloxacin**
23. This pain medication is commonly associated with hepatotoxicity. **Acetaminophen**
24. This branch of pharmacology describes what the drug does to the body, including therapeutic effects and adverse effects. **Pharmacodynamics**
25. This aminoglycoside carries the risk of ototoxicity. **Gentamicin**