Results

Adam Weber — 2nd Attempt

05:38

Time for this attempt

This survey has unlimited submissions. Take Now

Attempt History

Results
Attempt 1
Attempt 2

Your Answers:

1 Multiple Choice

The nurse is alerted to a fire on the hospital unit. The nurse turns off oxygen equipment and places wet towels along the base of the doors. Which step of the RACE acronym is the nurse implementing?

- (●) Contain the fire
- ( ) Evacuate
- ( ) Rescue
- ( ) Activate the fire alarm

Feedback

General Feedback

Answer: contain the fire

2 Multiple Choice

What action by the nursing student indicates a need for further instruction when speaking to a client with hearing loss?

- ○ The student speaks in a louder voice
- ◉ The student encourages the client to increase the volume of their hearing aides to maximum level
- ○ The student speaks in a low tone of voice
- ○ The student provides written information when possible

Feedback

General Feedback

Answer: increase volume of hearing aides to max level

3 Multiple Choice

Which of the following statements is true of acute pain?

- ◉ Acute pain usually develops quickly and can range from mild to severe
- ○ Acute pain typically lasts longer than six months
- ○ Acute pain is not indicative of protection of an injured area
- ○ Acute pain only causes sharp pain

Feedback

General Feedback

Answer: Acute pain usually develops quickly and can range from mild to severe

4 Fill in the Blank

A nurse is caring for a client who has a prescription for diazepam IV. The client weighs 122 pounds. The minimum therapeutic dose for diazepam is 0.17 mg/kg/day. What is the minimum therapeutic dosage the client should be prescribed per day? Round to the nearest tenth. 5.5

5 Multiple Choice

The nurse notices a small, contained fire in a client's trash can. What is the priority action?

- (●) Remove the client and others in danger
- ( ) Attempt to extinguish fire using proper equipment
- ( ) Pull the fire alarm to alert local fire authorities
- ( ) If safe to do so, turn off oxygen and close the room door

Feedback

General Feedback

Answer: remove client and others in danger

6 Multiple Choice

A client with advanced dementia who is oriented to self only experienced a fall on the previous shift. The nurse conducts a pain assessment. What assessment method will provide the nurse with the best information?

- ( ) Show the client a simple visual analog scale and ask the client to rate any pain according to the scale
- ( ) Assess vital signs and observe for evidence of stress response, such as elevated heart rate and respiratory rates
- (●) Ask the client yes-or-no questions related to pain in areas of the body specific to the type of fall experienced
- ( ) Observe the client for evidence of guarding or grimacing or other behavioral evidence of pain during activities

Feedback

General Feedback

Answer: yes or no questions; ask the patient subjective data to see if they are capable before relying on objective findings since objective findings could be indicative of another problem besides pain

7 Fill in the Blank

A nurse is preparing to administer acetaminophen 500 mg by mouth every 6 hours. Available is acetaminophen 1000 mg per tablet. How many tablet(s) should the nurse administer per dose?

0.5

8 Multiple Choice

While performing passive ROM, the patient reports discomfort with an abduction exercise. After stopping the exercise, which action does the nurse take next?

- ○ Report the pain to the healthcare provider
- ○ Offer a dose of PRN analgesia to the client
- ○ Ask the client if heat or ice to the area is preferred
- ● Complete a focused pain assessment

Feedback

General Feedback

Answer: assess pain (ADPIE!)

9 Multiple Choice

What is the best way the nurse should ask the client about current medication use when completing a medication reconciliation?

- ○ "What medications do you take at home?"
- ◉ "What medications, supplements, or vitamins do you currently take?"
- ○ "Do you take anything at home?"
- ○ "What do you use at home to help with your health problems?"

Feedback

General Feedback

Answer: What meds, supplements, vitamins do you currently take?

10 Multiple Choice

The nurse is caring for a patient who reports nausea in the post-anesthesia care unit (PACU). Which nurse statement demonstrates establishing priorities to promote patient safety?

- ◉ "I'm going to increase your head of the bed."
- ○ "Feel free to reposition yourself to get comfortable."
- ○ "Lay on your stomach to decrease nausea."
- ○ "Let's recline the bed flat to support your back."

Feedback

General Feedback

Answer: increase head of bed (to prevent aspiration)

11 Fill in the Blank

A nurse calculates output in the past four hours for a patient undergoing diuresis. The patient drank 200mL water, but then vomited 100mL. Normal saline ran at 100mL/hr. The patient also drank 100mL juice. Urine output is measured at 1,500mL, and there is 50mL of NGT drainage. The nurse calculates the patient's negative fluid volume to be what amount? Round to the nearest whole number. 950

12 Fill in the Blank

A nurse prepares digoxin 150mcg IM to administer to a patient. The nurse retrieves digoxin 0.1mg/mL from the medication dispensing system. How many mL does the nurse prepare? Round to the nearest tenth. 1.5

13 Multiple Answer

Which of the following techniques would assist postoperative clients in prevention of atelectasis and pneumonia? Select all that apply.

- [x] Coughing and deep breathing
- [x] Use of incentive spirometer
- [ ] Chest physiotherapy
- [ ] Pursed-lip breathing

Feedback

General Feedback

Answer: incentive spirometer, cough and deep breathe

14 Fill in the Blank

The nurse receives an order to administer 400mL of normal saline over 30 minutes. The tubing has a drop factor of 15gtt/mL. How many drops per minute should the nurse deliver? Round to the nearest whole number. 200

15 Multiple Choice

You are the nurse on the medical-surgical unit and reviewing the chart of a newly admitted patient to your unit for refractory chronic lower back pain. Which pharmacological intervention would you

expect to be ordered?

- ○ PO acetaminophen
- ○ PO morphine
- ○ PO ibuprofen
- ◉ IV morphine on PCA

Feedback

General Feedback

Answer: IV morphine on PCA

16 Multiple Choice

Your client with a pleural effusion is on 4L NC. She calls you to her room because her nose is starting to bleed. You stop the bleeding and the client's airway is clear. What intervention will help prevent future epistaxis?

- ○ Switching the client from a NC to a simple mask
- ○ Putting Vaseline in the client's nares
- ○ Allowing the client to shower daily
- ◉ Adding humidification to oxygen

Feedback

General Feedback

Answer: adding humidification

17 Fill in the Blank

The nurse prepares to initiate an IV iron dextran infusion at a rate of 50mg/min. The nurse retrieves normal saline 100mL with 1,250mg iron dextran. At what rate does the nurse set the IV infusion

pump in mL/hr? Round to the nearest whole number. 240

18 Multiple Choice

Which of the following steps are included in tracheostomy care? Select all that apply.

- ○ Monitor blood pressure when suctioning the tracheostomy
- ◉ Change the tracheostomy ties daily
- ○ Remove the outer cannula
- ○ Change the gauze under the tracheostomy plate weekly

Feedback

General Feedback

Answer: change ties daily

19 Multiple Answer

Which of the following are possible client risks due to peripheral sensory impairments? Select all that apply.

- ☐ Risk of impaired gas exchange
- ☑ Risk of falls
- ☑ Risk of infection
- ☑ Risk of altered skin integrity
- ☐ Risk of impaired nutrition

Feedback

General Feedback

Infection, falls, altered skin integrity

20 Multiple Choice

You are caring for a patient who has the following vital signs recorded during your assessment:

BP 105/62, HR 105, RR 20, Temp 102.9F oral, O2 sat 98% RA

Skin is hot, dry, and flushed. Patient reports a headache and refuses breakfast.

Based on the vital signs and clinical cues, which of the following actions should you prioritize?

- (●) Administer ordered antipyretics
- ( ) Document the findings and continue monitoring
- ( ) Encourage the patient to drink fluids
- ( ) Notify the healthcare provider about the elevated temperature

Feedback

General Feedback

Answer: give ordered antipyretics

Fever management is the first thing we would do because reducing the fever would prevent dehydration (and the need for increased PO intake). We would want to tell the doctor but after we intervened with an order that was already available to us. If the patient did not have an order for antipyretic then notifying the HCP would be the correct choice. We need to document the vitals but it's not the priority. The patient's BP is also probably lower and HR higher due to fever.

21 Multiple Choice

The student nurse cares for clients in a long-term care facility. The nurse reviews evidence-based fall prevention interventions with the student. Which statement by the student nurse requires follow up?

- ( ) "Clients at risk for falls should be encouraged to change positions slowly."
- ( ) "Clients should have bed alarms on with their beds in the low and locked position."
- ( ) "Clients that are high fall risk should wear socks with grippers on the bottom."

"Client rounding should be conducted every 90 minutes to assess needs."

Feedback

General Feedback

Answer: rounding every 90 minutes

22 Multiple Answer

Which of the following statements are true of PCAs? Select all that apply.

- [ ] PCAs are the best way to control any type of pain
- [x] PCA dose change or initiation must be verified by an additional RN
- [x] Complications of PCA use include respiratory depression and altered level of consciousness
- [x] The PCA button must be pushed only by the patient
- [x] Clients with a PCA should also be on a bowel regimen

Feedback

General Feedback

Answers: all but "PCAs are the best way to control any type of pain."

23 Fill in the Blank

A nurse prepares to administer a dose of methylprednisolone 25mg IV to a patient. The nurse retrieves methylprednisolone 0.05g/5mL from the medication dispensing system. How many mL does the nurse prepare? Round to the nearest tenth. 2.5

24 Multiple Choice

The nurse cares for a 178lb patient prescribed IV dobutamine at 0.7mcg/kg/min. Dobutamine is supplied in a concentration of 250mg/250mL. The nurse finds that the IV pump is infusing at 20mL/hr. What is the priority nursing action?

- ○ Assess the patient's vital signs
- ○ Assess the patient's EKG reading
- ● Stop the dobutamine infusion
- ○ Notify the healthcare provider

Feedback

General Feedback

Answer: stop the infusion!

1. Math: you need to figure out if the pump is set correctly in mL/hr. So first we need to find what the mL/hr should be:
    1. mL/hr = 250mL/250mg x 1mg/1000mcg x 0.7mcg/kg x 1 kg/2.2lb x 178lb/min x 60min/hr = 3.3982 ~ 3.4mL/hr is what the pump should be set at.
    2. Could also convert your lb to kg first (178lb/2.2 = 80.9091kg), then remember 0.7mcg PER kg PER minute = 0.7mcg x 80.9091kg = 56.6364mcg/min. Then you can plug in your numbers from there - mL/hr = 250mL/250mg x 1mg/1000mcg x 56.6364mcg/min x 60min/1 day = 3.4mL/hr
2. Based on your calculation, the pump is set at 20mL/hr and that is no where close to 3.4mL/hr. *We need to stop that infusion!!* Dobutamine is a medication that increases cardiac contractility and thus cardiac output. We use it for people with bad ejection fractions who are in cardiogenic shock because their heart cannot beat strong enough to perfuse their body. It can definitely mess with their VS and EKG, but the priority action is to stop the infusion immediately!

25 Multiple Choice

A nurse is admitting a patient with a history of frequent falls at home. Which intervention does the nurse implement as the priority?

- ○ Teach the patient to wear non-slip socks
- ○ Determine the patient's ability to use assistive devices
- ◉ Place the bed in the lowest position
- ○ Make a bedside commode easily available

Feedback

General Feedback

Answer: bed low; priority would be lowering the bed because getting out of the bed is the first step the patient has to do in order to perform another task. If the bed is high and the patient fell from a height, the patient wouldn't have the opportunity to use a bedside commode, walk with socks, or use assistive devices.