

Results

Adam Weber — 1st Attempt



20

Out of 20 points

06:08

Time for this attempt

1 attempt left

Take Now

▼ Attempt History

Results	Points	Score	(Highest score is kept)
Attempt 1	20 of 20	100%	(Highest score)

Your Answers:

1

1 / 1 point Multiple Choice

What is an adverse effect of gentamycin?

- ☐ cardiac toxicity
- ☐ alcohol toxicity
- ☒ renal toxicity
- ☐ liver toxicity

2

1 / 1 point Multiple Choice

What should be monitored in a client taking vancomycin?

- ☐ respiratory rate
- ☒ hearing test
- ☐ blood pressure
- ☐ blood glucose levels

3

1 / 1 point Multiple Choice

What is the primary mechanism of action of antiviral medications?

- ☐ Prevention of viral entry into host cells
- ☒ Inhibition of viral replication
- ☐ Stimulation of the immune system
- ☐ Neutralization of viral toxins

4

1 / 1 point Multiple Choice

Which of the following antiviral medications is commonly used to treat influenza A?

- ☒ Oseltamivir
- ☐ Acyclovir
- ☐ telbivudine
- ☐ Amantadine

5

1 / 1 point Multiple Choice

Which of the following is a contraindication to amoxicillin administration?

- ☐ poor condition of the teeth

☐ seizure disorder

☒ ✓ ☒ allergy

☐ Parkinson's disease

6 1 / 1 point Multiple Choice

What potential side effect should the nurse monitor for when administering acyclovir?

☐ Neurological impairment

☐ Cardiac dysrhythmias

☒ ✓ ☒ Renal toxicity

☐ Bradycardia

7 1 / 1 point Multiple Choice

Many of the immunosuppressant drugs that are administered following organ transplantation may also be used for treatment of what condition?

☐ AIDS

☒ ✓ ☒ autoimmune disorders

☐ Seizure disorders

☐ increased intracranial pressure

8 1 / 1 point Multiple Choice

Why should clients use non steroidal anti-inflammatory drugs (NSAIDs) cautiously on a long-term basis?

☐ Increased risk of central nervous system impairment

☐ Increased risk of blindness

☐ Increased risk of hearing impairment



☒ Increased risk of gastrointestinal (GI) bleeding

9

1 / 1 point Multiple Choice

What medication may cause reddish-orange discoloration of urine and other body fluids?

☐ tamoxifen

☐ vincristine

☐ methotrexate



☒ doxorubicin

10

1 / 1 point Multiple Choice

Which of these analgesic medications is a non-steroidal anti-inflammatory drug (NSAID)?



☒ ibuprofen

☐ acetaminophen

☐ morphine

☐ gold compound

11

1 / 1 point Multiple Choice

Which laboratory test should be monitored in a client taking acetaminophen?

☐ peak and trough levels

☐ cardiac enzymes

☐ white blood cell count



☒ liver enzymes

12

1 / 1 point Multiple Answer

Which adverse effects would be expected in a client receiving chemotherapy? Select all that apply.

☐ Increase in appetite



☒ Fatigue



☒ Nausea

☐ Increase in hair growth



☒ Anemia

13

1 / 1 point Multiple Choice

What is the therapeutic effect of an immunosuppressant medication?



☒ Decrease immune response

☐ Bradycardia

☐ Myocardial infarction

☐ Wheezing

14

1 / 1 point Multiple Choice

A client taking methotrexate should be instructed to:

☐ Increase alcohol intake

☐ Decrease fluid intake

☐ Stop the medication when symptoms improve



☒ Avoid people with infections

15

1 / 1 point Multiple Choice

A major adverse effect associated with tumor necrosis factor (TNF) blockers such as adalimumab is:



☒ Increased risk of infection

- ☐ Bradycardia
- ☐ Hearing loss
- ☐ Hyperthyroidism

16

1 / 1 point Multiple Choice

Cyclosporine is most commonly used for which purpose?



☒ Preventing organ rejection after transplantation

- ☐ Treating fungal infections
- ☐ Treating influenza infection
- ☐ Managing acute pain

17

1 / 1 point Multiple Choice

Before initiating therapy with a monoclonal antibody, the nurse should be most concerned about:



☒ Active infection

- ☐ Controlled hypertension
- ☐ Seasonal allergies
- ☐ Nearsightedness

18

1 / 1 point Multiple Choice

What important adverse effect is associated with fluconazole?

- ☐ Hypothyroidism
- ☐ Hyperglycemia
- ☐ Tendon rupture



☒ Liver toxicity

19

1 / 1 point Multiple Choice

Tamoxifen is primarily used in the treatment of:

- ☐ Tuberculosis
- ☐ Influenza
- ☐ Rheumatoid arthritis



☒ Breast cancer

20

1 / 1 point Multiple Choice

A client receiving methotrexate, cyclosporine, or a tumor necrosis factor (TNF) blocker should be taught that these medications may:

- ☐ Improve resistance to viral illnesses.
- ☐ Increase white blood cell production.



☒ Increase susceptibility to infection.

- ☐ Prevent all autoimmune reactions.