

1 **Molecular Evolution of the Meiotic Recombination Pathway in Mammals**

2 *Investigations*

3

4 Amy L. Dapper<sup>1,2\*</sup> and Bret A. Payseur<sup>1</sup>

5

6 <sup>1</sup> Laboratory of Genetics, University of Wisconsin, Madison, WI 53706, USA

7 <sup>2</sup> Current Address: Department of Biological Sciences, Mississippi State University, Mississippi State, MS

8 39762, USA

9 Running Title: Evolution of the Recombination Pathway

10 Keywords: (up to 5)

11

12 \* Corresponding Author : Amy L. Dapper

13 Address: 295 E. Lee Blvd., P.O. Box GY, Mississippi State, MS 39762

14 Phone: (662) 325-7575

15 Email: dapper@biology.msstate.edu

## Abstract

Meiotic recombination, the exchange of genetic material between homologous chromosomes during meiosis, is required for successful gametogenesis in most sexually reproducing species. Recombination is also a fundamental evolutionary force, influencing the fate of new mutations and determining the genomic scale over which selection shapes genetic variation. Despite the central importance of recombination, basic questions about its evolution have yet to be addressed. Although many genes that play roles in recombination have been identified, the molecular evolution of most of these genes remains uncharacterized. Using a phylogenetic comparative approach, we measure rates of evolution in 32 recombination pathway genes across 16 mammalian species, spanning primates, murids, and laurasithians. By analyzing a carefully-selected panel of genes involved in key components of recombination – spanning double strand break formation, strand invasion, the crossover/non-crossover decision, and resolution – we generate a comprehensive picture of the evolution of the recombination pathway in mammals. Recombination genes exhibit marked heterogeneity in the rate of protein evolution, both across and within genes. We report signatures of rapid evolution and positive selection that could underlie species differences in recombination rate. [WORK HERE]

Abstract Word Count : ( $< 250$ )

## Introduction

The reciprocal exchange of DNA between homologous chromosomes during meiosis – recombination – is required for successful gametogenesis in most species that reproduce sexually (Hassold and Hunt 2001). The rate of recombination is a major determinant of patterns of genetic diversity in populations, influencing the fate of new mutations (Hill and Robertson 1966), the efficacy of selection (Felsenstein 1974; Charlesworth *et al.* 1993; Comeron *et al.* 1999; Gonen *et al.* 2017), and important features of the genomic landscape (Begun and Aquadro 1992; Charlesworth *et al.* 1994; Duret and Arndt 2008).

Although recombination rate is often treated as a constant, this fundamental parameter evolves over time. Genomic regions ranging in size from short sequences to entire chromosomes vary in recombination rate – both within and between species (Burt and Bell 1987; Broman *et al.* 1998; Jeffreys *et al.* 2005; Coop and Przeworski 2007; Kong *et al.* 2010; Dumont *et al.* 2011; Smukowski and Noor 2011; Comeron *et al.* 2012; Segura *et al.* 2013; Dapper and Payseur 2017; Stapley *et al.* 2017).

Genome-wide association studies are beginning to reveal the genetic basis of differences in recombination rate within species. Individual recombination rates have been associated with variation in specific genes in populations of *Drosophila melanogaster* (Hunter *et al.* 2016), humans (Kong *et al.* 2008, 2014; Chowdhury *et al.* 2009; Fledel-Alon *et al.* 2011), domesticated cattle (Sandor *et al.* 2012; Ma *et al.* 2015; Kadri *et al.* 2016; Shen *et al.* 2018), domesticated sheep (Petit *et al.* 2017), Soay sheep (Johnston *et al.* 2016), and red deer (Johnston *et al.* 2018). Variants in several of these genes correlate with recombination rate in multiple species, including: *Rnf212* (Kong *et al.* 2008; Chowdhury *et al.* 2009; Fledel-Alon *et al.* 2011; Sandor *et al.* 2012; Johnston *et al.* 2016; Kadri *et al.* 2016; Petit *et al.* 2017), *Rnf212B* (Johnston *et al.* 2016, 2018; Kadri *et al.* 2016), *Rec8* (Sandor *et al.* 2012; Johnston *et al.* 2016, 2018), *Hei10/Ccnb1ip1* (Kong *et al.* 2014; Petit *et al.* 2017), *Msh4* (Kong *et al.* 2014; Ma *et al.* 2015; Kadri *et al.* 2016; Shen *et al.* 2018), *Cplx1* (Kong *et al.* 2014; Ma *et al.* 2015; Johnston *et al.* 2016; Shen *et al.* 2018) and *Prdm9* (Fledel-Alon *et al.* 2011; Sandor *et al.* 2012; Kong *et al.* 2014; Ma *et al.* 2015; Shen *et al.* 2018).

In contrast, the genetics of recombination rate variation among species remains poorly understood. Divergence at the di-cistronic gene *mei-217/mei-218* explains much of the disparity in genetic map length between *D. melanogaster* and *D. mauritiana* (Brand *et al.* 2018). *mei-217/mei-218* is the only gene known to confer a recombination rate difference between species, though quantitative trait loci that contribute to shifts in rate among subspecies of house mice have been identified (Dumont and Payseur 2010; Murdoch *et al.* 2010; Balcova *et al.* 2016).

One strategy for understanding how species diverge in recombination rate is to inspect patterns of molecular

evolution at genes involved in the recombination pathway. This approach incorporates knowledge of the molecular and cellular determinants of recombination and is motivated by successful examples. *mei-217/mei-218* was targeted for functional analysis based on its profile of rapid evolution between *D. melanogaster* and *D. mauritiana* (Brand *et al.* 2018). *Prdm9*, a protein that positions recombination hotspots in house mice and humans through histone methylation (Myers *et al.* 2010; Parvanov *et al.* 2010; Grey *et al.* 2011, Paigen2018; 2018), shows accelerated divergence across mammals (Oliver *et al.* 2009). The rapid evolution of *Prdm9* – which localizes to its zinc-finger DNA binding domain (Oliver *et al.* 2009) – appears to be driven by selective pressure to recognize new hotspot motifs as old ones are destroyed via biased gene conversion (Myers *et al.* 2010; Ubeda and Wilkins 2011; Lesecque *et al.* 2014; Latrille *et al.* 2017). Although these examples demonstrate the promise of signatures of molecular evolution for illuminating recombination rate differences between species, patterns of divergence have yet to be reported for most genes involved in meiotic recombination.

Mammals provide a useful system for dissecting the molecular evolution of the recombination pathway for several reasons. First, the evolution of recombination rate has been measured along the mammalian phylogeny (Dumont and Payseur 2008; Segura *et al.* 2013). Second, recombination rate variation has been associated with specific genes in mammalian populations (Kong *et al.* 2008, 2014; Chowdhury *et al.* 2009; Sandor *et al.* 2012; Ma *et al.* 2015; Johnston *et al.* 2016, 2018; Kadri *et al.* 2016; Petit *et al.* 2017; Shen *et al.* 2018). Third, laboratory mice have proven to be instrumental in the identification and functional characterization of recombination genes (Vries *et al.* 1999; Baudat *et al.* 2000; Romanienko and Camerini-Otero 2000; Yang *et al.* 2006; Ward *et al.* 2007; Schramm *et al.* 2011; Bisig *et al.* 2012; Bolcun-Filas and Schimenti 2012; La Salle *et al.* 2012; Kumar *et al.* 2015; Finsterbusch *et al.* 2016; Stanzione *et al.* 2016) .

Work in mice indicates that the mammalian recombination pathway is roughly divided into five major steps, each of which is regulated by a handful of genes. The first step is the formation of hundreds of double strand breaks (DSBs) throughout the genome (Bergerat *et al.* 1997; Keeney *et al.* 1997; Baudat *et al.* 2000; Romanienko and Camerini-Otero 2000; Baudat and Massy 2007; Finsterbusch *et al.* 2016; Lange *et al.* 2016). After formation, DSBs are identified, processed, and paired with their corresponding location on the homologous chromosome through the processes of homology search and strand invasion (Keeney 2007; Cloud *et al.* 2012; Brown and Bishop 2014; Finsterbusch *et al.* 2016; Kobayashi *et al.* 2016; Oh *et al.* 2016; Xu *et al.* 2017). The pairing of homologous chromosomes is then stabilized by a proteinaceous structure referred to as the synaptonemal complex (SC) (Meuwissen *et al.* 1992; Schmekel and Daneholt 1995; Costa *et al.* 2005; Vries *et al.* 2005; Hamer *et al.* 2006; Yang *et al.* 2006; Schramm *et al.* 2011; Fraune *et al.* 2014; Hernández-Hernández *et al.* 2016). The SC also forms a substrate on which the eventual crossover

events will take place [citations]. It is at this point that a small subset of DSBs is designated to mature into crossovers, leaving the majority of DSBs to be resolved as non-crossovers (Snowden *et al.* 2004; Yang *et al.* 2008; Reynolds *et al.* 2013; Finsterbusch *et al.* 2016; Rao *et al.* 2017). Finally, this designation is followed, and each DSB is repaired as a crossover or a non-crossover (Baker *et al.* 1996; Edelman *et al.* 1996; Lipkin *et al.* 2002; Rogacheva *et al.* 2014; Xu *et al.* 2017).

In this article, we examine the molecular evolution of 32 key recombination genes, evenly distributed across each major step in the recombination pathway, in 16 mammalian species spanning Primates, Rodents and Laurasiatherians. In addition to revealing patterns of divergence across diverse mammalian species, we leverage human polymorphism data to make robust evolutionary inferences. Our results provide a comprehensive picture of evolution in the recombination pathway in mammals and identify steps of the pathway most likely to contribute to differences in recombination rate between species.

## Materials and Methods

### Data Acquisition & Processing

We selected a focal panel of 32 recombination genes (See Table1). The genes included in this panel were selected to: (1) cover each major step in the recombination pathway as evenly as possible, (2) choose genes that we know are have integral functions in each step, and (3) include genes that have been associated with variation in recombination rate within mammalian populations. Reference sequences were downloaded for each gene from both NCBI and Ensembl (Release-89) [citations].

Alternative splicing is widespread and presents a challenge for molecular evolution studies [citations]. We used available testes expression datasets to select the transcript expressed in tissues of interest and to validate the computationally imputed annotations for each gene in each species. We downloaded the raw testes expression data for each mammalian species from NCBI GEO (Table S1) [citations]. We converted the SRA files into FASTQ files using SRAToolkit [citation]. The reads were mapped to an indexed reference genome (Table S2,3) [Bowtie2, citation] using tophat [citation]. The resulting bam files were sorted using Samtools and visualized using IGV 2.4.10 [citations]. This allowed us to: (1) identify the transcript that is expressed in the testes tissue, (2) check the reference transcript for errors, (3) revise the reference transcript based upon the transcript data.

We compared expression data to annotations from both the Ensembl & NCBI [citations]. When both transcripts were identical, we selected the NCBI transcript was the default. Ensembl transcript was used

instead when: (1) the NCBI reference sequences was not available for a given gene in a given species, (2) when none of the NCBI matched the expression data, or (3) when there were sequence differences between the two transcripts and the Ensembl transcript was more parsimonious. The use of testes expression data sets was a key data processing step and the inclusion of species in this study was primarily determined by the availability of testes expression data.

## Phylogenetic Comparative Approach in Mammals

For each genes we conducted a phylogenetic analysis by maximum likelihood (PAML 4.8) to measure the rate of evolution across the mammalian phylogeny and to detect molecular signatures indicative of positive selection [citations]. This approach requires a sequences alignment for each gene and a phylogenetic tree. For each gene, sequences were aligned using Translator X, a codon-based alignment tool powered by MUSCLE v3.8.31 (citations). Each alignment was examined by hand and, as necessary, edited. We selected a species tree for our analyses, based upon our current understanding of the phylogenetic relationship of the mammals included in our study (Figure1) [citations].

Due to the ambiguity in the relationship between Laurasithians and the placement of Tree Shrews, we also inferred the gene trees using MrBayes [citations]. This also allowed us to control for effects of incomplete lineage sorting (ILS) [citations]. The results using the gene trees did not differ in any significant manner and can be found in the supplemental info (Table S4).

For the majority of genes, transcripts from all 16 species were used (19 genes). However, for a number of genes, the chimpanzee and bonobo sequences were identical, in which case only the chimpanzee sequence was included in the analyses (11 genes). In one case, the chimpanzee, bonobo and human sequences were all identical, in which case only the human sequence was included in the analyses. In only a small number of instances, a suitable reference sequences could not be identified for a given species.

We estimated rates of synonymous and nonsynonymous substitutions per site using the CODEML program in PAML4.8 (citations). This program takes into account multiple substitutions per site, different rates of transitions and transversions, and effects of codon usage (citations). Rates of substitutions were computed for 6 different models: 0,1,2,7,8,8a (Table 2). The fit of each model was compared using a likelihood ratio test and the rates of substitutions are reported for the model of best fit for each gene.

[TURN INTO TABLE] (1) Comparison 1: Model 0 - one dN/dS ratio for all sites ( $<1$ ); Model 1 - two dN/dS ratios ( $<1$ ,  $=1$ ); Model 2 - three dN/dS ratios ( $<1$ ,  $=1$ ,  $>1$ ). (2) Comparison 2: Model 7 - beta distribution of 10 dN/dS values, all between 0 & 1; Model 8 - beta distribution of 10 dN/dS values, all between 0 & 1,

plus 11th category > 1. (3) Comparison 3: Model 8 - beta distribution of 10 dN/dS values, all between 0 & 1, plus 11th category > 1; Model 8a - beta distribution of 5 dN/dS values, all between 0 & 1, plus 6th category =1.

## **Polymorphism & Divergence in the Primate Lineage**

Human polymorphism data was downloaded from ExAC database.

Not available for 3 genes. Issues with ExAC data for: RNF212, MEI4 (and REC8)?

Pairwise divergence between humans and macaques was calculated using YN00 package in PAML.

Compared polymorphism within humans to divergence between human and macaques using the McDonald-Kreitman test.

## **Identifying Signatures**

Model comparison

## **Multinucleotide Mutations**

Multi-Nucleotide Mutations (MNMs) occur when two mutations happen simultaneously in close proximity (non-independent) [citations]. MNMs violate the assumption of PAMLs maximum likelihood model, which assumes that the probability of two simultaneous mutations in the same codon is zero [citations]. Recent work has shown that MNMs can frequently result in false positive signatures of positive selection in branch-site models in HyPhy [citations]. While we are not using branch-site models, the possibility remains that MNMs could be contributing to the signature of positive selection we are observing in some recombination genes. It is not possible to identify MNMs in our dataset, but we can identify codons with multiple differences (CMDs) that are likely to have arisen on a single branch in the phylogeny. We removed all CMDs that putatively arose on a single branch and then re-analyzed the subset of genes that exhibited a significant signature of positive selection in our original analyses.

To identify CMDs, we used PAML to reconstruct the ancestral sequence at each node in the phylogeny. For the reconstruction, Model 8 was chosen because we were specifically analyzing genes with a significant signature of positive selection when comparing Model 7 & Model 8. From the ancestrally reconstructed sequences, we identified any codons in which PAML inferred more than one substitution on a single branch. All identified CMDs were removed from the sequences in which they occurred. For example, if a CMD was



180 identified in an external branch, that codon would be replaced with ‘—’ only in the sequence of that species.  
181 If a CMD was inferred on an internal branch, the codon would be replaced with ‘—’ in all species connected to  
182 that internal branch. We re-ran our analyses in PAML with the sequences in which all CMDs were removed.

## Results

### Heterogeneous rates of evolution of recombination genes across the mammalian phylogeny

Do we observe elevated rates of evolution in certain steps in the recombination pathway? No - No significant difference in mean omega values among genes that function ( $p = 0.09767$ , Kruskal-Wallis rank sum test).

Do we observe more genes with signatures of positive selection in certain steps in the recombination pathway? Values too small to compare all 5 groups. Compared earlier steps ( $N = 15$ ) to later steps ( $N=17$ ). Significantly more genes with signatures of positive selection in the second half of the recombination pathway ( $p = 0.0457$ , Pearson's Chi-squared test)

### Evidence of positive selection - PAML

Of the 9 genes with significant signatures of positive selection (7vs8), only one (TEX11) retained the significant signature of selection after removing all CMDs. Two additional genes (REC8, RAD21L), also showed that model 8 was a significantly better fit than model 7. However, this is because models that allow a class of sites with a dN/dS of 1 are preferred over models that require all sites to have dN/dS values  $< 1$ . There is limited to no support for a class of sites with dN/dS  $> 1$ .

### Polymorphism & Divergence Data

## Discussion

The rate of meiotic recombination shapes major features of the genomic landscape and influences the efficacy of selection (???)

Recombination rate varies between species (???), between populations (???), within populations (???), and between the sexes (???)

Ample evidence indicates that phenotypic variation in recombination rate has a genetic component.

Recombination rate shows resemblance among relatives in human pedigrees [77,106,107], differs among lines raised in a common environment [74,83,85,96,108,109], and responds to artificial selection in *Drosophila melanogaster* and other insects.

Broad-sense or narrow-sense heritability estimates from humans, mice, insects, and maize range from 0.08 to 0.69 [73,106,107,110–115].

A host of experiments using insects attempted to increase and/or decrease the recombination rate [71,73,110,111,115,145–157], demonstrating the potential for recombination rate to respond to directional selection in nature.

Phylogenetic comparative methods suggest that the genome-wide recombination rate has increased during mammalian evolution [97].

PRDM9, a protein that helps determine the position of crossovers in mice and humans, possesses one of the most rapidly evolving (zinc-finger) domains in mammals [163,164].

Fecundity and recombination rate may be positively correlated in human mothers [77,165,166].

Finally, cellular needs to avoid nondisjunction (by generating at least one crossover per chromosome or chromosome arm) and to minimize costs of double-strand break repair should impose selective bounds on the genome-wide crossover rate in nature [1–4].

There is limited empirical evidence for a role of selection in the evolution of recombination rate.

1. Recombination rate evolves.
2. There is a genetic basis.
3. The same few genes pop up over and over again, suggesting there may be a relatively simple genetic basis to variation in recombination rate.
4. There has recently been an acceleration in our understanding of the genes involved in the recombination pathway.
5. Divergence in a subset of these genes is very likely to underlie trait differences.
6. This should be true whether or not those trait differences are generated by selection or drift.
7. Motivates a phylogenetic comparative study of genes in the recombination pathway among mammals.
8. Analyzing these genes from a pathway perspective may provide insight into which genes, or steps of the pathway, are most likely to be contributing to variation in recombination rate in mammals.
9. Also provides the opportunity to look for molecular signatures of adaptive evolution.

Meiotic recombination begins with the generation of 100's of double strand breaks (DSB) across the genome. **SPO11** directly produces the DSBs, but is recruited and activated by a handful of other proteins. The location of DSB are non-randomly distributed across the genome. **PRDM9** lays down methylation patterns via sequence-specific DNA binding.

## Results

## Discussion

Recombination rate shows resemblance among relatives in human pedigrees [citations], differs among lines raised in a common environment [citations], and responds to artificial selection [citations]. Artificial selection experiments to increase and/or decrease the recombination rate [citations], demonstrate the potential for recombination rate to respond to directional selection in nature. Beyond the lab, comparisons between species and between populations have uncovered pervasive, and in some cases, rapid evolution of recombination rate [citations]. There is limited direct, empirical evidence for a role of selection in the evolution of recombination rate. Indirect evidence includes the observation via phylogenetic comparative methods that the genome-wide recombination rate appears to have increased during mammalian evolution [citation] and the observation that fecundity and recombination rate may be positively correlated in human mothers [citations]. However, due to the importance of recombination rate in shaping the genome and response to evolution, the value of understanding its evolution is not strictly tied to the role of selection in shaping the trait.

While *Prdm9* clearly plays a major role in the positioning of recombination events within the genome, it is less clear that *Prdm9* significantly impacts the total number of recombination events in the genome [citations]. *Prdm9* plays an very early role in the patterning of recombination events.

**Table 1** : Recombination Genes

Gene	Function	Meiosis-Specific?
<b>A)</b>	<b>DSB Formation</b>	
<i>HORMAD1</i>	chromosome axis, promotes DSB formation	Yes
<i>HORMAD2</i>	chromosome axis	Yes
<i>MEI4</i>	promotes DSB formation ( <i>MCD recombinosome</i> )	Yes
<i>REC114</i>	promotes DSB formation ( <i>MCD recombinosome</i> )	Yes
<i>IHO1</i>	promotes DSB formation ( <i>MCD recombinosome</i> )	Yes
<i>SPO11</i>	transesterase, catalyzes the formation of DSBs	Yes
<b>B)</b>	<b>DSB Processing/Strand Invasion</b>	
<i>MRE11</i>	nuclease, required for DSB formation & processing ( <i>MRN Complex</i> )	No
<i>NBS1</i>	phosphopeptide binding, required for DSB formation	No
—	& processing ( <i>MRN Complex</i> )	
<i>RAD50</i>	ATPase/DNA binding protein, required for DSB formation	No

Gene	Function	Meiosis-Specific?
—	& processing ( <i>MRN Complex</i> )	
<i>BRCC3</i>	deubiquitinase, DSB processing	No
<i>DMC1</i>	recombinase, strand invasion & homologous pairing	Yes
<i>RAD51</i>	recombinase, strand invasion & homologous pairing	No
<i>SPATA22</i>	strand invasion & homologous pairing	Yes
<i>MEIOB</i>	oligonucleotide binding, strand invasion & homologous pairing	Yes
<i>MCMDC2</i>	helicase, stabilizes homologous pairing	Yes
<b>C)</b>	<b>Homologous Pairing</b>	
<i>REC8</i>	cohesion core	Yes
<i>RAD21L</i>	cohesion core	Yes
<i>SYCP1</i>	synaptonemal complex - transverse filament	
<i>SYCP2</i>	synaptonemal complex - axial element	
<i>TEX12</i>	synaptonemal complex - central element	
<b>D1)</b>	<b>Crossover vs. Non-Crossover - MutS Recruitment</b>	
<i>TEX11</i>		
<i>SHOC1</i>		
<i>CNTD1</i>		
<i>RNF212</i>		
<i>RNF212B</i>		
<i>MSH4</i>	recombination crossover control	
<i>MSH5</i>	recombination crossover control	
<b>D2)</b>	<b>Crossover vs. Non-Crossover - MutL Recruitment</b>	
<i>MER3</i>		
<i>HEI10</i>		
<i>MLH1</i>	promotion of meiotic crossing over	
<i>MLH3</i>	promotion of meiotic crossing over	
<i>MUS81</i>		



**Table 2:** PAML analysis of 32 recombination genes in mammals (Yang 2007).

<i>Gene</i>	<i>bp</i>	<i>N</i>	$\omega$	<i>M</i>	<i>M1-M2</i>	<i>p-value</i>	<i>M7-M8</i>	<i>p-value</i>	<i>M8a-M8</i>	<i>p-value</i>
<b>A)</b>										
<i>HORMAD1</i>	1212	16	0.3036	7	0	1.000	1.795	0.4076	—	—
<i>HORMAD2</i>	981	15	0.3153	7	0	1.000	3.650	0.1612	—	—
<i>MEI4</i>	1170	16	0.4332	7	0	1.000	0.005	0.9976	—	—
<i>REC114</i>	870	15	0.4003	7	0	1.000	5.384	0.0677	—	—
<i>IHO1</i>	1824	16	0.7095	8	13.061	<b>0.0015</b>	17.571	<b>0.0002</b>	14.527	<b>0.0001</b>
<i>SPO11</i>	1188	15	0.1654	7	0	1.000	4.648	0.0980	—	—
<b>B)</b>										
<i>MRE11</i>	2136	16	0.1688	8	0.363	0.8342	11.931	<b>0.0026</b>	4.706	<b>0.0301</b>
<i>NBS1</i>	2289	15	0.4183	8	0	1.000	12.763	<b>0.0017</b>	4.087	<b>0.0432</b>
<i>RAD50</i>	3936	16	0.1006	7	0	1.000	0.301	0.8605	—	—
<i>BRCC3</i>	954	15	0.0602	7	0	1.000	0.250	0.8826	—	—
<i>DMC1</i>	1020	15	0.0351	1	0.488	0.7835	5.000	0.0821	—	—
<i>RAD51</i>	1017	16	0.0268	7	0	1.000	0	1.000	—	—
<i>SPATA22</i>	1101	16	0.4893	7	0	1.000	0.429	0.8070	—	—
<i>MEIOB</i>	1425	16	0.2341	7	0	1.000	0.665	0.7172	—	—
<i>MCMDC2</i>	2052	16	0.2239	7	0	1.000	0.628	0.7307	—	—
<b>C)</b>										
<i>REC8</i>	1833	16	0.3698	8	0	1.000	14.690	<b>0.0006</b>	5.927	<b>0.0149</b>
<i>RAD21L</i>	1686	15	0.503	8	12.124	<b>0.0023</b>	32.050	<b>&gt;0.0001</b>	12.049	<b>0.0005</b>
<i>SYCP1</i>	3015	16	0.4337	8	8.711	<b>0.0128</b>	26.860	<b>&gt;0.0001</b>	9.243	<b>0.0024</b>
<i>SYCP2</i>	4650	16	0.5572	8	11.584	<b>0.0031</b>	37.200	<b>&gt;0.0001</b>	15.838	<b>0.0001</b>
<i>TEX12</i>	369	14	0.2297	7	0.0565	0.9721	1.549	0.4610	—	—
<b>D1)</b>										
<i>TEX11</i>	2844	15	0.8483	8	60.872	<b>&gt;0.0001</b>	82.665	<b>&gt;0.0001</b>	61.141	<b>&gt;0.0001</b>
<i>SHOC1</i>	4644	16	0.6113	8	12.447	<b>0.0020</b>	30.561	<b>&gt;0.0001</b>	15.645	<b>0.0001</b>
<i>CNTD1</i>	1026	15	0.2496	7	0	1.000	0.936	0.6263	—	—
<i>RNF212</i>	948	16	0.5014	8	0	1.000	16.366	<b>0.0003</b>	5.202	<b>0.0226</b>
<i>RNF212B</i>	906	14	0.4066	7	0	1.000	0.500	0.7788	—	—
<i>MSH4</i>	2814	16	0.2132	8	16.608	<b>0.0002</b>	39.447	<b>&gt;0.0001</b>	23.238	<b>&gt;0.0001</b>

<i>Gene</i>	<i>bp</i>	<i>N</i>	$\omega$	<i>M</i>	<i>M1-M2</i>	<i>p-value</i>	<i>M7-M8</i>	<i>p-value</i>	<i>M8a-M8</i>	<i>p-value</i>
<i>MSH5</i>	2565	15	0.1642	7	0	<i>1.000</i>	4.214	<i>0.1216</i>	—	—
<b>D2)</b>										
<i>MER3</i>	4458	16	0.3633	8a	0	<i>1.000</i>	12.838	<b><i>0.0016</i></b>	3.109	<i>0.0779</i>
<i>HEI10</i>	831	15	0.1226	7	0	<i>1.000</i>	0.250	<i>0.8826</i>	—	—
<i>MLH1</i>	2313	15	0.1652	8a	0	<i>1.000</i>	12.221	<b><i>0.0022</i></b>	0.280	<i>0.5970</i>
<i>MLH3</i>	4419	16	0.4444	7	0	<i>1.000</i>	3.757	<i>0.1528</i>	—	—
<i>MUS81</i>	1665	16	0.2124	7	0	<i>1.000</i>	0.628	<i>0.7304</i>	—	—



**Table 3:** PAML - MNM Analysis

<i>Gene</i>	<i>bp</i>	<i>N</i>	$\omega$	<i>M</i>	<i>M1-M2</i>	<i>p-value</i>	<i>M7-M8</i>	<i>p-value</i>	<i>M8a-M8</i>	<i>p-value</i>
<i>IHO1</i>	1824	16	0.6104	7	0	<i>1.000</i>	0.258	<i>0.8789</i>	—	—
<i>MRE11</i>	2136	16	0.1330	7	0.226	<i>0.8930</i>	3.056	<i>0.2169</i>	—	—
<i>NBS1</i>	2289	15	0.3413	7	0	<i>1.000</i>	1.956	<i>0.3761</i>	—	—
<i>REC8</i>	1833	16	0.2905	7	0	<i>1.000</i>	5.321	<i>0.0699</i>	—	—
<i>RAD21L</i>	1686	15	0.4271	8a	2.329	<i>0.3121</i>	9.497	<b><i>0.0087</i></b>	1.620	<i>0.2031</i>
<i>SYCP1</i>	3015	16	0.3731	8a	3.328	<i>0.1893</i>	13.440	<b><i>0.0012</i></b>	2.122	<i>0.1452</i>
<i>SYCP2</i>	4650	16	0.4752	7	0	<i>1.000</i>	1.758	<i>0.4151</i>	—	—
<i>TEX11</i>	2844	15	0.7287	<b>8</b>	9.989	<b><i>0.0068</i></b>	18.776	<b><i>0.0001</i></b>	10.656	<b><i>0.0011</i></b>
<i>SHOC1</i>	4644	16	0.5519	8a	0	<i>1.000</i>	7.439	<b><i>0.0242</i></b>	0.292	<i>0.5887</i>
<i>RNF212</i>	948	16	0.3685	7	0	<i>1.000</i>	0	<i>1.000</i>	—	—
<i>MSH4</i>	2814	16	0.1509	7	0	<i>1.000</i>	2.079	<i>0.3536</i>	—	—

**Table 4:** Polymorphism & Divergence Data

<i>Gene</i>	<i>bp</i>	<i>Pn</i>	<i>Ps</i>	<i>Pn/Ps</i>	<i>Dn</i>	<i>Ds</i>	<i>Dn/Ds</i>	<i>MK Test</i>	$\alpha$	<i>NI</i>	
<b>A)</b>											
<i>HORMAD1</i>	XXX	84	35	2.4000	5	12	0.4167	<b>0.0018</b>	-4.7600	5.7600	Neg.
<i>HORMAD2</i>	XXX	80	31	2.5806	7	9	0.7778	<b>0.0404</b>	-2.3180	3.3180	Neg.
<i>MEI4</i>	XXX	15	7	2.1429	24	9	2.6667	<i>0.7679</i>	0.1964	0.8036	—
<i>REC114</i>	XXX	76	37	2.0541	11	14	0.7857	<b>0.0392</b>	-1.6143	2.6143	Neg.
<i>IHO1</i>	XXX	130	64	2.0313	36	19	1.8947	<i>0.8718</i>	-0.0720	1.0720	—
<i>SPO11</i>	XXX	118	52	2.2692	11	22	0.5000	<b>0.0001</b>	-3.5385	4.5385	Neg.
<b>B)</b>											
<i>MRE11</i>	XXX	211	86	2.4535	5	35	0.1429	<b>&gt;0.0001</b>	-16.1744	17.1744	Neg.
<i>NBS1</i>	XXX	221	93	2.3763	34	25	1.3600	<i>0.0666</i>	-0.7473	1.7473	—
<i>RAD50</i>	XXX	303	118	2.5678	8	43	0.1860	<b>&gt;0.0001</b>	-12.8019	13.8019	Neg.
<i>BRCC3</i>	XXX	13	21	0.6190	2	6	0.3333	0.6888	-0.8571	1.8571	—
<i>DMC1</i>	XXX	72	42	1.7143	0	11	0.0000	<b>&gt;0.0001</b>	—	—	Neg.
<i>RAD51</i>	XXX	50	48	1.0417	0	13	0.0000	<b>&gt;0.0001</b>	—	—	Neg.
<i>SPATA22</i>	XXX	114	45	2.5333	21	10	2.1000	<i>0.6700</i>	-0.2063	1.2063	—
<i>MEIOB</i>	XXX	91	40	2.2750	20	22	0.9091	<b>0.0200</b>	-1.5025	2.5025	Neg.
<i>MCMDC2</i>	XXX	165	54	3.0556	16	26	0.6154	<b>&gt;0.0001</b>	-3.9653	4.9653	Neg.
<b>C)</b>											
<i>REC8</i>	XXX	147	76	1.9342	38	31	1.2258	<i>0.1164</i>	-0.5779	1.5779	—
<i>RAD21L</i>	XXX	51	17	3.000	27	13	2.0769	<i>0.5051</i>	-0.4444	1.4444	—
<i>SYCP1</i>	XXX	213	100	2.1300	33	37	1.2222	<i>0.0546</i>	-0.7427	1.7427	—
<i>SYCP2</i>	XXX	429	154	2.8506	74	53	1.3962	<b>0.0005</b>	-1.0417	2.0417	Neg.
<i>TEX12</i>	XXX	31	16	1.9375	2	4	0.5000	0.1836	-2.875	3.875	—
<b>D1)</b>											
<i>TEX11</i>	XXX	126	81	1.5556	55	25	2.200	<i>0.2234</i>	0.2929	0.7071	—
<i>SHOC1</i>	XXX	368	124	2.9677	85	37	2.2973	<i>0.2521</i>	-0.2918	1.2918	—
<i>CNTD1</i>	XXX	81	47	1.7234	13	8	1.6250	<i>1.0000</i>	-0.0606	1.0606	—
<i>RNF212</i>	XXX	—	—	—	17	18	0.9444	—	—	—	—
<i>RNF212B</i>	XXX	368	124	2.9677	8	12	0.6667	<b>0.0013</b>	-3.4516	4.4516	Neg.
<i>MSH4</i>	XXX	260	94	2.7660	24	29	0.8276	<b>&gt;0.0001</b>	-2.3422	3.3422	Neg.

<i>Gene</i>	<i>bp</i>	<i>Pn</i>	<i>Ps</i>	<i>Pn/Ps</i>	<i>Dn</i>	<i>Ds</i>	<i>Dn/Ds</i>	<i>MK Test</i>	$\alpha$	<i>NI</i>	
<i>MSH5</i>	XXX	197	104	1.8942	19	33	0.5758	<b>0.0002</b>	-2.2900	3.2900	Neg.
<b>D2)</b>											
<i>MER3</i>	XXX	402	143	2.8112	54	44	1.2273	<b>0.0004</b>	-1.2906	2.2906	Neg.
<i>HEI10</i>	XXX	73	33	2.2121	4	5	0.8000	<i>0.1541</i>	-1.7652	2.7652	—
<i>MLH1</i>	XXX	255	90	2.8333	9	29	0.3103	<b>&gt;0.0001</b>	-8.1296	9.1296	Neg.
<i>MLH3</i>	XXX	437	167	2.6168	77	57	1.3509	<b>0.0012</b>	-0.9370869	1.937087	Neg.
<i>MUS81</i>	XXX	208	81	2.5679	17	40	0.4250	<b>&gt;0.0001</b>	-5.0421	6.0421	Neg.

## Acknowledgements

A.L.D. was supported by NHGRI Training Grant to the Genomic Sciences Training Program 5T32HG002760.

B.A.P. was supported by NIH grant R01 GM100426A and NSF grant DEB 1353737.

**Table R1:** Testes Expression Datasets

<i>Species</i>	<b>GEO Accession</b>	<b>Reference</b>
<i>Bos taurus</i>	GSM1020728 & GSM1020746	Merkin <i>et al.</i> (2012)
<i>Callithrix jacchus</i>	GSM1227961, GSM1227962 & GSM1227963	Cortez <i>et al.</i> (2014)
<i>Canis lupus familiaris</i>	GSM747469 & GSM1359286	Derti <i>et al.</i> (2012), Vandewege <i>et al.</i> (2016)
<i>Eptesicus fuscus</i>	GSM1359287	Vandewege <i>et al.</i> (2016)
<i>Equus caballus</i>	GSM1139276 & GSM1359288	Coleman <i>et al.</i> (2013), Vandewege <i>et al.</i> (2016)
<i>Gorilla gorilla</i>	GSM752663	Brawand <i>et al.</i> (2011)
<i>Homo sapiens</i>	GSM752707 & GSM752708	Brawand <i>et al.</i> (2011)
<i>Macaca mulatta</i>	GSM752642 & GSM752643	Brawand <i>et al.</i> (2011)
<i>Mus musculus</i>	GSM752629 & GSM752630	Brawand <i>et al.</i> (2011)
<i>Ovis aries</i>	GSM1666944 & GSM1666936	Guan <i>et al.</i> (2017)
<i>Pan paniscus</i>	GSM752690	Brawand <i>et al.</i> (2011)
<i>Pan troglodytes</i>	GSM752678	Brawand <i>et al.</i> (2011)
<i>Pongo pygmaeus</i>	GSM1858310 & GSM1858311	Carelli <i>et al.</i> (2016)
<i>Rattus norvegicus</i>	GSM1278058	Cortez <i>et al.</i> (2014)
<i>Sus scrofa</i>	GSM1902350, GSM2033157 & GSM2033163	Li <i>et al.</i> (2016), Yang <i>et al.</i> (2017)
<i>Tupaia chinensis</i>	GSM957062	Fan <i>et al.</i> (2013)

**Table R2:** Reference Genomes (O’Leary *et al.* 2015)

<i>Species</i>	<b>Assembly</b>	<b>RefSeq Accession</b>	<b>WGS Project Reference</b>
<i>Bos taurus</i>	Bos_taurus_UMD_3.1.1	GCF_000003055.6	Zimin <i>et al.</i> (2009)
<i>Callithrix jacchus</i>	Callithrix_jacchus-3.2	GCF_000004665.1	-
<i>Canis lupus familiaris</i>	CanFam3.1	GCF_000002285.3	Lindblad-Toh <i>et al.</i> (2005)
<i>Eptesicus fuscus</i>	EptFus1.0	GCF_000308155.1	-
<i>Equus caballus</i>	EquCab2.0	GCF_000002305.2	Wade <i>et al.</i> (2009)
<i>Gorilla gorilla</i>	gorGor4	GCF_000151905.2	Scally <i>et al.</i> (2012)
<i>Homo sapiens</i>	GRCh38.p10	GCF_000001405.36	-
<i>Macaca mulatta</i>	Mmul_8.0.1	GCF_000772875.2	Zimin <i>et al.</i> (2014)
<i>Mus musculus</i>	GRCm38.p5	GCF_000001635.25	-
<i>Ovis aries</i>	Oar_v4.0	GCF_000298735.2	Consortium <i>et al.</i> (2010)

<i>Species</i>	<i>Assembly</i>	<i>RefSeq Accession</i>	<i>WGS Project Reference</i>
<i>Pan paniscus</i>	panpan1.1	GCF_000258655.2	Prüfer <i>et al.</i> (2012)
<i>Pan troglodytes</i>	Pan_tro_3.0	GCF_000001515.7	Consortium <i>et al.</i> (2005)
<i>Pongo abelii</i>	P_pygmaeus_2.0.2	GCF_000001545.4	Locke <i>et al.</i> (2011)
<i>Rattus norvegicus</i>	Rnor_6.0	GCF_000001895.5	Consortium and others (2004)
<i>Sus scrofa</i>	Sscrofa11.1	GCF_000003025.6	-
<i>Tupaia chinensis</i>	TupChi_1.0	GCF_000334495.1	Fan <i>et al.</i> (2013)

**Table R3:** Ensembl Annotations

<i>Species</i>	<i>Assembly</i>	<i>RefSeq Accession</i>	<i>WGS Project Reference</i>
<i>Bos taurus</i>	Bos_taurus_UMD_3.1	GCF_000003055.3	Zimin <i>et al.</i> (2009)
<i>Callithrix jacchus</i>	Callithrix_jacchus-3.2	GCF_000004665.1	-
<i>Canis lupus familiaris</i>	CanFam3.1	GCF_000002285.3	Lindblad-Toh <i>et al.</i> (2005)
<i>Eptesicus fuscus</i>	-	-	-
<i>Equus caballus</i>	EquCab2.0	GCF_000002305.2	Wade <i>et al.</i> (2009)
<i>Gorilla gorilla</i>	gorGor3.1	GCF_000151905.1	-
<i>Homo sapiens</i>	GRCh38.p10	GCF_000001405.36	-
<i>Macaca mulatta</i>	Mmul_8.0.1	GCF_000772875.2	Zimin <i>et al.</i> (2014)
<i>Mus musculus</i>	GRCm38.p5	GCF_000001635.25	-
<i>Ovis aries</i>	Oar_v3.1	GCF_000298735.1	Consortium <i>et al.</i> (2010)
<i>Pan paniscus</i>	panpan1.1	GCF_000258655.2	Prüfer <i>et al.</i> (2012)
<i>Pan troglodytes</i>	CHIMP2.1.4	GCF_000001515.6	Consortium <i>et al.</i> (2005)
<i>Pongo abelii</i>	PPYG2	GCF_000001545.4	Locke <i>et al.</i> (2011)
<i>Rattus norvegicus</i>	Rnor_6.0	GCF_000001895.5	Consortium and others (2004)
<i>Sus scrofa</i>	Sscrofa11.1	GCF_000003025.6	-
<i>Tupaia chinensis</i>	-	-	-

**Table S1:** Sequence divergence between Human (*Homo sapiens*) and Rhesus Macaque (*Macaca mulatta*) (Yang and Nielsen 2000, Yang (2007)). Steps: A - double strand break (DSB) formation, B - DSB processing & Strand Invasion, C - Homologous Pairing, D1 - crossover (CO) vs. non-crossover (NCO) step1 - MutS, D2 - CO vs. NCO step 2 - MutL.

<i>Gene</i>	<i>bp</i>	$\omega$	<i>S</i>	<i>N</i>	<i>t</i>	$\kappa$	<i>dN</i>	<i>dS</i>
<b>A)</b>								
<i>HORMAD1</i>	XXX	<b>0.0901</b>	273.9	908.1	0.0443	3.8819	0.0044 +- 0.0022	0.0490 +- 0.0137
<i>HORMAD2</i>	XXX	<b>0.295</b>	256.7	664.3	0.0531	4.2164	0.0106 +- 0.0040	0.0360 +- 0.0121
<i>MEI4</i>	XXX	<b>0.7252</b>	331	824	0.0822	4.6295	0.0247 +/- 0.0056	0.0341 +/- 0.0104
<i>REC114</i>	XXX	<b>0.3239</b>	237.2	557.8	0.0974	2.9455	0.0200 +/- 0.0061	0.0618 +/- 0.0168
<i>IHO1</i>	XXX	<b>0.6608</b>	509	1273	0.0951	3.6035	0.0276 +- 0.0047	0.0418 +- 0.0094
<i>SPO11</i>	XXX	<b>0.1434</b>	291.2	896.8	0.0872	2.5317	0.0118 +/- 0.0036	0.0823 +/- 0.0178
<b>B)</b>								
<i>MRE11</i>	XXX	<b>0.0392</b>	479.4	1644.6	0.0597	2.6154	0.0030 +- 0.0014	0.0778 +- 0.0135
<i>NBS1</i>	XXX	<b>0.4155</b>	553.7	1705.3	0.0804	5.0955	0.0199 +- 0.0035	0.0480 +- 0.0097
<i>RAD50</i>	XXX	<b>0.0714</b>	1118.7	2817.3	0.0401	5.0903	0.0028 +- 0.0010	0.0399 +- 0.0062
<i>BRCC3</i>	XXX	<b>0.0979</b>	264	609	0.028	4.6	0.0025 +- 0.0020	0.0252 +- 0.0100
<i>DMC1</i>	XXX	<b>0.0000</b>	273.7	746.3	0.0335	5.1279	0.0000 +- 0.0000	0.0416 +- 0.0127
<i>RAD51</i>	XXX	<b>0.0000</b>	306.5	710.5	0.0398	6.7467	0.0000 +- 0.0000	0.0441 +- 0.0124
<i>SPATA22</i>	XXX	<b>0.4523</b>	247.8	841.2	0.0879	3.6505	0.0230 +- 0.0053	0.0508 +- 0.0150
<i>MEIOB</i>	XXX	<b>0.2462</b>	348.9	1064.1	0.0927	4.3887	0.0176 +- 0.0041	0.0715 +- 0.0151
<i>MCMDC2</i>	XXX	<b>0.2108</b>	534	1509	0.0635	7.8547	0.0107 +- 0.0027	0.0507 +- 0.0101
<b>C)</b>								
<i>REC8</i>	XXX	<b>0.477</b>	497	1138	0.1293	2.8869	0.0323 +- 0.0054	0.0678 +- 0.0122
<i>RAD21L</i>	XXX	<b>0.6334</b>	427.5	1237.5	0.0735	5.6876	0.0213 +- 0.0042	0.0337 +- 0.0091
<i>SYCP1</i>	XXX	<b>0.3676</b>	761.6	2166.4	0.0628	4.8307	0.0145 +- 0.0026	0.0393 +- 0.0074
<i>SYCP2</i>	XXX	<b>0.3873</b>	1070.7	3519.3	0.0854	5.994	0.0208 +- 0.0025	0.0537 +- 0.0074
<i>TEX12</i>	XXX	<b>0.1349</b>	80.2	288.8	0.05	1.9678	0.0070 +- 0.0049	0.0516 +- 0.0260
<b>D1)</b>								
<i>TEX11</i>	XXX	<b>0.9068</b>	805.9	1933.1	0.0897	7.8022	0.0290 +- 0.0040	0.0320 +- 0.0064
<i>SHOC1</i>	XXX	<b>0.7225</b>	1203	3129	0.0865	9.5737	0.0261 +- 0.0029	0.0361 +- 0.0057
<i>CNTD1</i>	XXX	<b>0.6803</b>	335.3	651.7	0.065	8.0721	0.0187 +- 0.0054	0.0274 +- 0.0092
<i>RNF212</i>	XXX	<b>0.387</b>	243.2	572.8	0.1342	4.996	0.0304 +- 0.0074	0.0785 +- 0.0189
<i>RNF212B</i>	XXX	<b>0.2566</b>	255.6	644.4	0.0685	3.4122	0.0125 +- 0.0044	0.0488 +- 0.0143
<i>MSH4</i>	XXX	<b>0.2635</b>	731.3	2073.7	0.058	7.5194	0.0112 +- 0.0023	0.0425 +- 0.0079
<i>MSH5</i>	XXX	<b>0.2106</b>	728.7	1770.3	0.0643	3.9993	0.0102 +- 0.0024	0.0486 +- 0.0085

<i>Gene</i>	<i>bp</i>	$\omega$	<i>S</i>	<i>N</i>	<i>t</i>	$\kappa$	<i>dN</i>	<i>dS</i>
<b>D2)</b>								
<i>MER3</i>	XXX	<b>0.3247</b>	987.6	3317.4	0.0703	7.0099	0.0159 +- 0.0022	0.0488 +- 0.0074
<i>HEI10</i>	XXX	<b>0.3235</b>	241.5	589.5	0.0329	5.9591	0.0068 +- 0.0034	0.0211 +- 0.0095
<i>MLH1</i>	XXX	<b>0.0924</b>	602.3	1665.7	0.0522	2.4752	0.0048 +- 0.0017	0.0521 +- 0.0097
<i>MLH3</i>	XXX	<b>0.4919</b>	1209.8	3149.2	0.0949	6.4296	0.0246 +- 0.0028	0.0500 +- 0.0067
<i>MUS81</i>	XXX	<b>0.1299</b>	465.8	1187.2	0.1106	5.7915	0.0128 +- 0.0033	0.0983 +- 0.0158

**Table S2:** PAML - Gene Trees

<i>Gene</i>	<i>bp</i>	<i>N</i>	$\omega$	<i>M</i>	<i>M1-M2</i>	<i>p-value</i>	<i>M7-M8</i>	<i>p-value</i>	<i>M8a-M8</i>	<i>p-value</i>
<b>A)</b>										
<i>HORMAD1</i>	1212	16	0.3037	7	0	1.000	3.135	0.2086	—	—
<i>HORMAD2</i>	981	15	0.3290	1	0	1.000	3.881	0.1436	—	—
<i>MEI4</i>	1170	16	0.4310	7	0	1.000	0.058	0.9715	—	—
<i>REC114</i>	870	15	0.4237	7	0	1.000	4.1874	XXXX	—	—
<i>IHO1</i>	1824	16	0.7099	8	13.384	0.0012	17.714	0.0001	14.707	0.0001
<i>SPO11</i>	1188	15	0.1701	7	0	1.000	4.697	0.0955	—	—
<b>B)</b>										
<i>MRE11</i>	2136	16	0.1686	8	0.636	0.7277	12.014	XXXX	4.822	XXXX
<i>NBS1</i>	2289	15	0.4185	8	0	1.000	12.899	XXXX	4.298	XXXX
<i>RAD50</i>	3936	16	XXXX	XXXX	XXXX	XXXX	XXXX	XXXX	—	—
<i>BRCC3</i>	954	15	0.0601	7	0	1.000	0.573	XXXX	—	—
<i>DMC1</i>	1020	15	0.0365	7	0	1.000	4.288	0.1172	—	—
<i>RAD51</i>	1017	16	0.0322	1	0	1.000	0.562	XXXX	—	—
<i>SPATA22</i>	1101	16	0.4932	7	0	1.000	0.200	XXXX	—	—
<i>MEIOB</i>	1425	16	0.2340	7	0	1.000	0.221	XXXX	—	—
<i>MCMD2C2</i>	2052	16	0.2242	7	0	1.000	0.610	0.7370	—	—
<b>C)</b>										

270 *REC8* | 1833 | 16 | 0.3698 | **8** | 0 | 1.000 | 14.690 | **0.0006** | 5.927 | **0.0149** |  
 271 *RAD21L* | 1686 | 15 | 0.503 | **8** | 12.124 | **0.0023** | 32.050 | **>0.0001** | 12.049 | **0.0005** | *SYCP1* | 3015 | 16  
 272 | 0.4337 | **8** | 8.711 | **0.0128** | 26.860 | **>0.0001** | 9.243 | **0.0024** | *SYCP2* | 4650 | 16 | 0.5572 | **8** | 11.584 |  
 273 **0.0031** | 37.200 | **>0.0001** | 15.838 | **0.0001** |  
 274 *TEX12* | 369 | 14 | 0.2297 | 7 | 0.0565 | 0.9721 | 1.549 | 0.4610 | — | — **D1**) | | | | | | | | | | *TEX11* | 2844 |  
 275 15 | 0.8483 | **8** | 60.872 | **>0.0001** | 82.665 | **>0.0001** | 61.141 | **>0.0001** | *SHOC1* | 4644 | 16 | 0.6113 | **8**  
 276 | 12.447 | **0.0020** | 30.561 | **>0.0001** | 15.645 | **0.0001** | *CNTD1* | 1026 | 15 | 0.2496 | 7 | 0 | 1.000 | 0.936  
 277 | 0.6263 | — | —  
 278 *RNF212* | 948 | 16 | 0.5014 | **8** | 0 | 1.000 | 16.366 | **0.0003** | 5.202 | **0.0226** | *RNF212B* | 906 | 14 | 0.4066 |  
 279 7 | 0 | 1.000 | 0.500 | 0.7788 | — | — *MSH4* | 2814 | 16 | 0.2132 | **8** | 16.608 | **0.0002** | 39.447 | **>0.0001** |  
 280 23.238 | **>0.0001** | *MSH5* | 2565 | 15 | 0.1642 | 7 | 0 | 1.000 | 4.214 | 0.1216 | — | — **D2**) | | | | | | | | | |



281 *MER3* | 4458 | 16 | 0.3633 | 8a | 0 | 1.000 | 12.838 | **0.0016** | 3.109 | 0.0779 |  
282 *HEI10* | 831 | 15 | 0.1226 | 7 | 0 | 1.000 | 0.250 | 0.8826 | — | — *MLH1* | 2313 | 15 | 0.1652 | 8a | 0 | 1.000 |  
283 12.221 | **0.0022** | 0.280 | 0.5970 | *MLH3* | 4419 | 16 | 0.4444 | 7 | 0 | 1.000 | 3.757 | 0.1528 | — | — *MUS81*  
284 | 1665 | 16 | 0.2124 | 7 | 0 | 1.000 | 0.628 | 0.7304 | — | —

## 285 References

- 286 Baker S. M., A. W. Plug, T. A. Prolla, C. E. Bronner, and A. C. Harris *et al.*, 1996 Involvement of mouse  
287 *mlh1* in dna mismatch repair and meiotic crossing over. *Nature genetics* 13: 336.
- 288 Balcova M., B. Faltusova, V. Gergelits, T. Bhattacharyya, and O. Mihola *et al.*, 2016 Hybrid sterility locus  
289 on chromosome x controls meiotic recombination rate in mouse. *PLoS genetics* 12: e1005906.
- 290 Baudat F., K. Manova, J. P. Yuen, M. Jasin, and S. Keeney, 2000 Chromosome synapsis defects and sexually  
291 dimorphic meiotic progression in mice lacking *spo11*. *Molecular cell* 6: 989–998.
- 292 Baudat F., and B. de Massy, 2007 Regulating double-stranded dna break repair towards crossover or  
293 non-crossover during mammalian meiosis. *Chromosome research* 15: 565–577.
- 294 Begun D. J., and C. F. Aquadro, 1992 Levels of naturally occurring dna polymorphism correlate with  
295 recombination rates in *d. melanogaster*. *Nature* 356: 519.
- 296 Bergerat A., B. de Massy, D. Gadelle, P.-C. Varoutas, and A. Nicolas *et al.*, 1997 An atypical topoisomerase  
297 ii from archaea with implications for meiotic recombination. *Nature* 386: 414.
- 298 Bisig C. G., M. F. Guiraldelli, A. Kouznetsova, H. Scherthan, and C. Höög *et al.*, 2012 Synaptonemal  
299 complex components persist at centromeres and are required for homologous centromere pairing in mouse  
300 spermatocytes. *PLoS genetics* 8: e1002701.
- 301 Bolcun-Filas E., and J. C. Schimenti, 2012 Genetics of meiosis and recombination in mice. *International*  
302 *review of cell and molecular biology* 298: 179–227.
- 303 Brand C. L., M. V. Cattani, S. B. Kingan, E. L. Landeen, and D. C. Presgraves, 2018 Molecular evolution at  
304 a meiosis gene mediates species differences in the rate and patterning of recombination. *Current Biology* 28:  
305 1289–1295.
- 306 Brawand D., M. Soumillon, A. Necsulea, P. Julien, and G. Csárdi *et al.*, 2011 The evolution of gene expression  
307 levels in mammalian organs. *Nature* 478: 343.
- 308 Broman K. W., J. C. Murray, V. C. Sheffield, R. L. White, and J. L. Weber, 1998 Comprehensive human

309 genetic maps: Individual and sex-specific variation in recombination. *The American Journal of Human*  
310 *Genetics* 63: 861–869.

311 Brown M. S., and D. K. Bishop, 2014 DNA strand exchange and recombination homologs in meiosis. *Cold Spring*  
312 *Harbor perspectives in biology* a016659.

313 Burt A., and G. Bell, 1987 Red queen versus tangled bank models. *Nature* 330: 118.

314 Carelli F. N., T. Hayakawa, Y. Go, H. Imai, and M. Warnefors *et al.*, 2016 The life history of retrocopies  
315 illuminates the evolution of new mammalian genes. *Genome research* gr-198473.

316 Charlesworth B., M. Morgan, and D. Charlesworth, 1993 The effect of deleterious mutations on neutral  
317 molecular variation. *Genetics* 134: 1289–1303.

318 Charlesworth B., P. Jarne, and S. Assimakopoulos, 1994 The distribution of transposable elements within and  
319 between chromosomes in a population of *drosophila melanogaster*. iii. element abundances in heterochromatin.  
320 *Genetics Research* 64: 183–197.

321 Chowdhury R., P. R. Bois, E. Feingold, S. L. Sherman, and V. G. Cheung, 2009 Genetic analysis of variation  
322 in human meiotic recombination. *PLoS genetics* 5: e1000648.

323 Cloud V., Y.-L. Chan, J. Grubb, B. Budke, and D. K. Bishop, 2012 Rad51 is an accessory factor for  
324 dmc1-mediated joint molecule formation during meiosis. *Science* 337: 1222–1225.

325 Coleman S. J., Z. Zeng, M. S. Hestand, J. Liu, and J. N. Macleod, 2013 Analysis of unannotated equine  
326 transcripts identified by mRNA sequencing. *PLoS One* 8: e70125.

327 Comeron J. M., M. Kreitman, and M. Aguadé, 1999 Natural selection on synonymous sites is correlated with  
328 gene length and recombination in *drosophila*. *Genetics* 151: 239–249.

329 Comeron J. M., R. Ratnappan, and S. Bailin, 2012 The many landscapes of recombination in *drosophila*  
330 *melanogaster*. *PLoS genetics* 8: e1002905.

331 Consortium R. G. S. P., and others, 2004 Genome sequence of the brown norway rat yields insights into  
332 mammalian evolution. *Nature* 428: 493.

333 Consortium T. C. S. A., R. H. Waterson, E. S. Lander, and R. K. Wilson, 2005 Initial sequence of the  
334 chimpanzee genome and comparison with the human genome. *Nature* 437: 69.

335 Consortium I. S. G., A. Archibald, N. Cockett, B. Dalrymple, and T. Faraut *et al.*, 2010 The sheep genome

reference sequence: A work in progress. *Animal genetics* 41: 449–453.

Coop G., and M. Przeworski, 2007 An evolutionary view of human recombination. *Nature Reviews Genetics* 8: 23.

Cortez D., R. Marin, D. Toledo-Flores, L. Froidevaux, and A. Liechti *et al.*, 2014 Origins and functional evolution of y chromosomes across mammals. *Nature* 508: 488.

Costa Y., R. Speed, R. Öllinger, M. Alsheimer, and C. A. Semple *et al.*, 2005 Two novel proteins recruited by synaptonemal complex protein 1 (sycp1) are at the centre of meiosis. *Journal of cell science* 118: 2755–2762.

Dapper A. L., and B. A. Payseur, 2017 Connecting theory and data to understand recombination rate evolution. *Phil. Trans. R. Soc. B* 372: 20160469.

Derti A., P. Garrett-Engele, K. D. MacIsaac, R. C. Stevens, and S. Sriram *et al.*, 2012 A quantitative atlas of polyadenylation in five mammals. *Genome research* gr-132563.

Dumont B. L., and B. A. Payseur, 2010 Evolution of the genomic recombination rate in murid rodents. *Genetics*.

Dumont B. L., M. A. White, B. Steffy, T. Wiltshire, and B. A. Payseur, 2011 Extensive recombination rate variation in the house mouse species complex inferred from genetic linkage maps. *Genome research* 21: 114–125.

Duret L., and P. F. Arndt, 2008 The impact of recombination on nucleotide substitutions in the human genome. *PLoS genetics* 4: e1000071.

Edelmann W., P. E. Cohen, M. Kane, K. Lau, and B. Morrow *et al.*, 1996 Meiotic pachytene arrest in *mlh1*-deficient mice. *Cell* 85: 1125–1134.

Fan Y., Z.-Y. Huang, C.-C. Cao, C.-S. Chen, and Y.-X. Chen *et al.*, 2013 Genome of the chinese tree shrew. *Nature communications* 4: 1426.

Felsenstein J., 1974 The evolutionary advantage of recombination. *Genetics* 78: 737–756.

Finsterbusch F., R. Ravindranathan, I. Dereli, M. Stanzione, and D. Tränkner *et al.*, 2016 Alignment of homologous chromosomes and effective repair of programmed dna double-strand breaks during mouse meiosis require the minichromosome maintenance domain containing 2 (*mcmcdc2*) protein. *PLoS genetics* 12: e1006393.

Fledel-Alon A., E. M. Leffler, Y. Guan, M. Stephens, and G. Coop *et al.*, 2011 Variation in human

364 recombination rates and its genetic determinants. PLoS one 6: e20321.

365 Fraune J., M. Alsheimer, J. Redolfi, C. Brochier-Armanet, and R. Benavente, 2014 Protein sycp2 is an ancient  
366 component of the metazoan synaptonemal complex. Cytogenetic and genome research 144: 299–305.

367 Gonen S., M. Battagin, S. E. Johnston, G. Gorjanc, and J. M. Hickey, 2017 The potential of shifting  
368 recombination hotspots to increase genetic gain in livestock breeding. Genetics Selection Evolution 49: 55.

369 Grey C., P. Barthès, G. Chauveau-Le Friec, F. Langa, and F. Baudat *et al.*, 2011 Mouse prdm9 dna-binding  
370 specificity determines sites of histone h3 lysine 4 trimethylation for initiation of meiotic recombination. PLoS  
371 biology 9: e1001176.

372 Grey C., F. Baudat, and B. de Massy, 2018 PRDM9, a driver of the genetic map. PLoS genetics 14: e1007479.

373 Guan Y., G. Liang, G. B. Martin, and others, 2017 Functional changes in mRNA expression and alternative  
374 pre-mRNA splicing associated with the effects of nutrition on apoptosis and spermatogenesis in the adult  
375 testis. BMC genomics 18: 64.

376 Hamer G., K. Gell, A. Kouznetsova, I. Novak, and R. Benavente *et al.*, 2006 Characterization of a novel  
377 meiosis-specific protein within the central element of the synaptonemal complex. Journal of cell science 119:  
378 4025–4032.

379 Hassold T., and P. Hunt, 2001 To err (meiotically) is human: The genesis of human aneuploidy. Nature  
380 Reviews Genetics 2: 280.

381 Hernández-Hernández A., S. Masich, T. Fukuda, A. Kouznetsova, and S. Sandin *et al.*, 2016 The central  
382 element of the synaptonemal complex in mice is organized as a bilayered junction structure. J Cell Sci 129:  
383 2239–2249.

384 Hill W. G., and A. Robertson, 1966 The effect of linkage on limits to artificial selection. Genetics Research 8:  
385 269–294.

386 Hunter C. M., W. Huang, T. F. Mackay, and N. D. Singh, 2016 The genetic architecture of natural variation  
387 in recombination rate in drosophila melanogaster. PLoS genetics 12: e1005951.

388 Jeffreys A. J., R. Neumann, M. Panayi, S. Myers, and P. Donnelly, 2005 Human recombination hot spots  
389 hidden in regions of strong marker association. Nature genetics 37: 601.

390 Johnston S. E., C. Bérénos, J. Slate, and J. M. Pemberton, 2016 Conserved genetic architecture underlying  
391 individual recombination rate variation in a wild population of soay sheep (ovis aries). Genetics genetics–115.

392 Johnston S. E., J. Huisman, and J. M. Pemberton, 2018 A genomic region containing rec8 and rnf212b is

393 associated with individual recombination rate variation in a wild population of red deer (*cervus elaphus*). G3:  
394 Genes, Genomes, Genetics g3–200063.

395 Kadri N. K., C. Harland, P. Faux, N. Cambisano, and L. Karim *et al.*, 2016 Coding and noncoding variants  
396 in *hfm1*, *mlh3*, *msh4*, *msh5*, *rnf212*, and *rnf212b* affect recombination rate in cattle. *Genome research*.

397 Keeney S., C. N. Giroux, and N. Kleckner, 1997 Meiosis-specific dna double-strand breaks are catalyzed by  
398 spo11, a member of a widely conserved protein family. *Cell* 88: 375–384.

399 Keeney S., 2007 Spo11 and the formation of dna double-strand breaks in meiosis, pp. 81–123 in *Recombination*  
400 *and meiosis*, Springer.

401 Kobayashi W., M. Takaku, S. Machida, H. Tachiwana, and K. Maehara *et al.*, 2016 Chromatin architecture  
402 may dictate the target site for *dmc1*, but not for *rad51*, during homologous pairing. *Scientific reports* 6:  
403 24228.

404 Kong A., G. Thorleifsson, H. Stefansson, G. Masson, and A. Helgason *et al.*, 2008 Sequence variants in the  
405 *rnf212* gene associate with genome-wide recombination rate. *Science* 319: 1398–1401.

406 Kong A., G. Thorleifsson, D. F. Gudbjartsson, G. Masson, and A. Sigurdsson *et al.*, 2010 Fine-scale  
407 recombination rate differences between sexes, populations and individuals. *Nature* 467: 1099.

408 Kong A., G. Thorleifsson, M. L. Frigge, G. Masson, and D. F. Gudbjartsson *et al.*, 2014 Common and  
409 low-frequency variants associated with genome-wide recombination rate. *Nature genetics* 46: 11.

410 Kumar R., N. Ghyselinck, K.-i. Ishiguro, Y. Watanabe, and A. Kouznetsova *et al.*, 2015 MEI4: A central  
411 player in the regulation of meiotic dna double strand break formation in the mouse. *J Cell Sci jcs*–165464.

412 La Salle S., K. Palmer, M. O’Brien, J. C. Schimenti, and J. Eppig *et al.*, 2012 Spata22, a novel vertebrate-  
413 specific gene, is required for meiotic progress in mouse germ cells. *Biology of reproduction* 86: 45–1.

414 Lange J., S. Yamada, S. E. Tischfield, J. Pan, and S. Kim *et al.*, 2016 The landscape of mouse meiotic  
415 double-strand break formation, processing, and repair. *Cell* 167: 695–708.

416 Latrille T., L. Duret, and N. Lartillot, 2017 The red queen model of recombination hot-spot evolution: A  
417 theoretical investigation. *Phil. Trans. R. Soc. B* 372: 20160463.

418 Lesecque Y., S. Glémin, N. Lartillot, D. Mouchiroud, and L. Duret, 2014 The red queen model of recombination  
419 hotspots evolution in the light of archaic and modern human genomes. *PLoS genetics* 10: e1004790.

420 Li Y., J. Li, C. Fang, L. Shi, and J. Tan *et al.*, 2016 Genome-wide differential expression of genes and small

rnas in testis of two different porcine breeds and at two different ages. Scientific reports 6: 26852.

Lindblad-Toh K., C. M. Wade, T. S. Mikkelsen, E. K. Karlsson, and D. B. Jaffe *et al.*, 2005 Genome sequence, comparative analysis and haplotype structure of the domestic dog. Nature 438: 803.

Lipkin S. M., P. B. Moens, V. Wang, M. Lenzi, and D. Shanmugarajah *et al.*, 2002 Meiotic arrest and aneuploidy in mlh3-deficient mice. Nature genetics 31: 385.

Locke D. P., L. W. Hillier, W. C. Warren, K. C. Worley, and L. V. Nazareth *et al.*, 2011 Comparative and demographic analysis of orang-utan genomes. Nature 469: 529.

Ma L., J. R. O’Connell, P. M. VanRaden, B. Shen, and A. Padhi *et al.*, 2015 Cattle sex-specific recombination and genetic control from a large pedigree analysis. PLoS genetics 11: e1005387.

Merkin J., C. Russell, P. Chen, and C. B. Burge, 2012 Evolutionary dynamics of gene and isoform regulation in mammalian tissues. Science 338: 1593–1599.

Meuwissen R., H. H. Offenberg, A. Dietrich, A. Riesewijk, and M. van Iersel *et al.*, 1992 A coiled-coil related protein specific for synapsed regions of meiotic prophase chromosomes. The EMBO Journal 11: 5091.

Murdoch B., N. Owen, S. Shirley, S. Crumb, and K. W. Broman *et al.*, 2010 Multiple loci contribute to genome-wide recombination levels in male mice. Mammalian genome 21: 550–555.

Myers S., R. Bowden, A. Tumian, R. E. Bontrop, and C. Freeman *et al.*, 2010 Drive against hotspot motifs in primates implicates the prdm9 gene in meiotic recombination. Science 327: 876–879.

Oh J., A. Al-Zain, E. Cannavo, P. Cejka, and L. S. Symington, 2016 Xrs2 dependent and independent functions of the mre11-rad50 complex. Molecular cell 64: 405–415.

Oliver P. L., L. Goodstadt, J. J. Bayes, Z. Birtle, and K. C. Roach *et al.*, 2009 Accelerated evolution of the prdm9 speciation gene across diverse metazoan taxa. PLoS genetics 5: e1000753.

O’Leary N. A., M. W. Wright, J. R. Brister, S. Ciufo, and D. Haddad *et al.*, 2015 Reference sequence (refseq) database at ncbi: Current status, taxonomic expansion, and functional annotation. Nucleic acids research 44: D733–D745.

Parvanov E. D., P. M. Petkov, and K. Paigen, 2010 Prdm9 controls activation of mammalian recombination hotspots. Science 327: 835–835.

Petit M., J.-M. Astruc, J. Sarry, L. Drouilhet, and S. Fabre *et al.*, 2017 Variation in recombination rate and

its genetic determinism in sheep populations. *Genetics* 163: 300123.

Prüfer K., K. Munch, I. Hellmann, K. Akagi, and J. R. Miller *et al.*, 2012 The bonobo genome compared with the chimpanzee and human genomes. *Nature* 486: 527.

Rao H. P., H. Qiao, S. K. Bhatt, L. R. Bailey, and H. D. Tran *et al.*, 2017 A sumo-ubiquitin relay recruits proteasomes to chromosome axes to regulate meiotic recombination. *Science* 355: 403–407.

Reynolds A., H. Qiao, Y. Yang, J. K. Chen, and N. Jackson *et al.*, 2013 RNF212 is a dosage-sensitive regulator of crossing-over during mammalian meiosis. *Nature genetics* 45: 269.

Rogacheva M. V., C. M. Manhart, C. Chen, A. Guarne, and J. Surtees *et al.*, 2014 Mlh1-mlh3, a meiotic crossover and dna mismatch repair factor, is a msh2-msh3-stimulated endonuclease. *Journal of Biological Chemistry* jbc-M113.

Romanienko P. J., and R. D. Camerini-Otero, 2000 The mouse spo11 gene is required for meiotic chromosome synapsis. *Molecular cell* 6: 975–987.

Sandor C., W. Li, W. Coppieters, T. Druet, and C. Charlier *et al.*, 2012 Genetic variants in rec8, rnf212, and prdm9 influence male recombination in cattle. *PLoS genetics* 8: e1002854.

Scally A., J. Y. Dutheil, L. W. Hillier, G. E. Jordan, and I. Goodhead *et al.*, 2012 Insights into hominid evolution from the gorilla genome sequence. *Nature* 483: 169.

Schmekel K., and B. Daneholt, 1995 The central region of the synaptonemal complex revealed in three dimensions. *Trends in cell biology* 5: 239–242.

Schramm S., J. Fraune, R. Naumann, A. Hernandez-Hernandez, and C. Höög *et al.*, 2011 A novel mouse synaptonemal complex protein is essential for loading of central element proteins, recombination, and fertility. *PLoS genetics* 7: e1002088.

Segura J., L. Ferretti, S. Ramos-Onsins, L. Capilla, and M. Farré *et al.*, 2013 Evolution of recombination in eutherian mammals: Insights into mechanisms that affect recombination rates and crossover interference. *Proceedings of the Royal Society of London B: Biological Sciences* 280: 20131945.

Shen B., J. Jiang, E. Seroussi, G. E. Liu, and L. Ma, 2018 Characterization of recombination features and the genetic basis in multiple cattle breeds. *BMC genomics* 19: 304.

Smukowski C., and M. Noor, 2011 Recombination rate variation in closely related species. *Heredity* 107: 496.

Snowden T., S. Acharya, C. Butz, M. Berardini, and R. Fishel, 2004 HSMH4-hSMH5 recognizes holliday junctions and forms a meiosis-specific sliding clamp that embraces homologous chromosomes. *Molecular cell*

15: 437–451.

Stanzione M., M. Baumann, F. Papanikos, I. Dereli, and J. Lange *et al.*, 2016 Meiotic dna break formation requires the unsynapsed chromosome axis-binding protein iho1 (*ccdc36*) in mice. *Nature cell biology* 18: 1208.

Stapley J., P. G. Feulner, S. E. Johnston, A. W. Santure, and C. M. Smadja, 2017 Variation in recombination frequency and distribution across eukaryotes: Patterns and processes. *Phil. Trans. R. Soc. B* 372: 20160455.

Ubeda F., and J. Wilkins, 2011 The red queen theory of recombination hotspots. *Journal of evolutionary biology* 24: 541–553.

Vandewege M. W., R. N. Platt, D. A. Ray, and F. G. Hoffmann, 2016 Transposable element targeting by piRNAs in laurasiatherians with distinct transposable element histories. *Genome biology and evolution* 8: 1327–1337.

Vries S. S. de, E. B. Baart, M. Dekker, A. Siezen, and D. G. de Rooij *et al.*, 1999 Mouse msh5-like protein msh5 is required for proper chromosome synapsis in male and female meiosis. *Genes & Development* 13: 523–531.

Vries F. A. de, E. de Boer, M. van den Bosch, W. M. Baarends, and M. Ooms *et al.*, 2005 Mouse sycp1 functions in synaptonemal complex assembly, meiotic recombination, and xy body formation. *Genes & development* 19: 1376–1389.

Wade C., E. Giolotto, S. Sigurdsson, M. Zoli, and S. Gnerre *et al.*, 2009 Genome sequence, comparative analysis, and population genetics of the domestic horse. *Science* 326: 865–867.

Ward J. O., L. G. Reinholdt, W. W. Motley, L. M. Niswander, and D. C. Deacon *et al.*, 2007 Mutation in mouse hei10, an e3 ubiquitin ligase, disrupts meiotic crossing over. *PLoS genetics* 3: e139.

Xu Y., R. A. Greenberg, E. Schonbrunn, and P. J. Wang, 2017 Meiosis-specific proteins meiob and spata22 cooperatively associate with the single-stranded dna-binding replication protein a complex and dna double-strand breaks. *Biology of reproduction* 96: 1096–1104.

Yang Z., and R. Nielsen, 2000 Estimating synonymous and nonsynonymous substitution rates under realistic evolutionary models. *Molecular Biology and Evolution* 17: 32–43.

Yang F., R. De La Fuente, N. A. Leu, C. Baumann, and K. J. McLaughlin *et al.*, 2006 Mouse sycp2 is required for synaptonemal complex assembly and chromosomal synapsis during male meiosis. *The Journal of Cell Biology* 173: 497–507.

Yang Z., 2007 PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution* 24:



506 1586–1591. <https://doi.org/10.1093/molbev/msm088>

507 Yang F., K. Gell, G. W. Van Der Heijden, S. Eckardt, and N. A. Leu *et al.*, 2008 Meiotic failure in male mice  
508 lacking an x-linked factor. *Genes & development* 22: 682–691.

509 Yang Y., G. Liang, G. Niu, Y. Zhang, and R. Zhou *et al.*, 2017 Comparative analysis of dna methylome and  
510 transcriptome of skeletal muscle in lean-, obese-, and mini-type pigs. *Scientific reports* 7: 39883.

511 Zimin A. V., A. L. Delcher, L. Florea, D. R. Kelley, and M. C. Schatz *et al.*, 2009 A whole-genome assembly  
512 of the domestic cow, *bos taurus*. *Genome biology* 10: R42.

513 Zimin A. V., A. S. Cornish, M. D. Maudhoo, R. M. Gibbs, and X. Zhang *et al.*, 2014 A new rhesus macaque  
514 assembly and annotation for next-generation sequencing analyses. *Biology direct* 9: 20.