

**ELEC 221 Lecture 04**  
**Properties of the CT Fourier series and the**  
**Gibbs phenomenon**

Tuesday 20 September 2022

# Announcements

- Quiz 2 today
- Assignment 1 due tomorrow at 23:59
  - no late submissions
  - include statement of contributions or you will get 0, no exceptions
- Assignment 2 (computational assignment) released tomorrow

## Last time

We explored complex exponential signals, and saw how they are eigenfunctions of LTI systems.

Given

then

## Last time

We expressed CT signals as Fourier series, and computed their Fourier coefficients.

**Fourier synthesis equation:**



**Fourier analysis equation:**

## Last time

We computed the Fourier coefficients of

$$x(t) = e^{-t}, -1 < t < 1$$

but we got a funny result.



What is going on here?

## Learning outcomes:

- State the Dirichlet conditions for convergence of a Fourier series
- Describe the Gibbs phenomenon
- Compute the power and energy of a signal, and state Parseval's theorem for CT periodic signals

# Dirichlet conditions

Can we *always* express a signal as a Fourier series?




Image credit: *Sine, square, triangle, and sawtooth waveforms* (author: Omegatron)

[https://en.wikipedia.org/wiki/Triangle\\_wave#/media/File:Waveforms.svg](https://en.wikipedia.org/wiki/Triangle_wave#/media/File:Waveforms.svg) (CC BY-SA 3.0)

## Dirichlet conditions

The **Dirichlet conditions** can help us here.

If over one period, a signal  $x(t)$

1. is single-valued
2. is absolutely integrable
3. has a finite number of maxima and minima
4. has a finite number of discontinuities

then the Fourier series converges to

- $x(t)$  where it is continuous
- half the value of the jump where it is discontinuous

These conditions are **sufficient** but not necessary.



Recall the synthesis equation:

Consider what happens if we truncate the series:

How good of an approximation is this?

## Approximation error

Can look at the approximation error


How much error is there over a given period?

This should go to 0 as  $N \rightarrow \infty$ .

## Example: the square wave

Let's consider a square wave signal with period  $2\pi$ :

$$x(t) = \begin{cases} 1, & 0 < t < \pi, \\ -1, & \pi < t < 2\pi \end{cases}$$



## Example: the square wave

Let's compute its Fourier coefficients.

First, note that since  $x(t)$  is an odd function, we can express it using only sin terms:

You can derive that for a  $2\pi$ -periodic function, the coefficients have the following form:

## Example: the square wave

$$b_k = \frac{1}{\pi} \int_0^{2\pi} x(\tau) \sin(k\tau) d\tau$$

Since the function is symmetric, we can rewrite this:



## Example: the square wave

Putting everything together,

Does this really converge? Let's try some increasingly good approximations by truncating the series:

# The Gibbs phenomenon

This is called the Gibbs phenomenon.



Truncation to *any*  $N$  results in “ringing” at the discontinuities.

## The Gibbs phenomenon

The amount of “overshoot” is about 9% of the jump of the discontinuity, no matter the size of  $N$ . With a bit of work, we can derive this!



## The Gibbs phenomenon

Let's evaluate what is in the sum...

Let's make use of an identity for complex numbers:

# The Gibbs phenomenon

Let's evaluate what is in the sum...

## The Gibbs phenomenon

$$\sum_{k=1}^N \cos(kx) = \Re \left[ e^{jNx} \frac{\sin(Nx)}{\sin(x)} \right]$$

Apply Euler's relation and apply some trig identities...

# The Gibbs phenomenon

Let's put this back in:

Now, we want to find  $t$  where the most amount of ring occurs;  
take the derivative!

# The Gibbs phenomenon

This is true when

# The Gibbs phenomenon

Evaluate the sum at this point:

Let's make a change of variables:

We care about the case where  $N \rightarrow \infty$ .

## The Gibbs phenomenon

Recall that for small  $x$ ,  $\sin x \approx x$ :

We can get a numerical value for the integral...

The highest peak of the ring is 0.17898 higher than it should be.  
This is  $\approx 9\%$  of the jump!

Break time

Go to `menti.com`



As long as you truncate, the ringing doesn't go away. However, we saw earlier that

This quantity here is actually the **energy** of the error.

Energy of a signal  $x(t)$  over a single period:

Power of a signal over a period:

We can relate this back to the Fourier coefficients...

# Signal power and energy

This is called **Parseval's relation**

The average power of a signal is the sum of the average powers contributed from each harmonic component.

Fourier coefficients have some really useful properties that help us evaluate them.

What happens when we apply the following transformations to the Fourier series representations of our signals?

- Superposition
- Time shift / scale / reversal
- Multiplication

# Properties of Fourier series

Fourier coefficients combine linearly.

Suppose we have two signals  $x(t), y(t)$  with period  $T$ ,

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{jk\omega t},$$

$$y(t) = \sum_{k=-\infty}^{\infty} b_k e^{jk\omega t},$$

Then the signal  $z(t) = Ax(t) + By(t)$  has the form

Time shift  $x(t) \rightarrow x(t - t_0)$ :

## Properties of Fourier series

Time scale  $x(t) \rightarrow x(\alpha t)$ .

If original period was  $T$ , new period is  $T/\alpha$ , new fundamental frequency is  $\omega\alpha$

where



## Properties of Fourier series

Multiplication leads to convolution of the coefficients:

Given

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{jk\omega t},$$

$$y(t) = \sum_{k=-\infty}^{\infty} b_k e^{jk\omega t},$$

Then the signal  $z(t) = x(t)y(t)$  has the form

Today's learning outcomes were:

- State the Dirichlet conditions for convergence of a Fourier series
- Describe the Gibbs phenomenon
- Compute the power and energy of a signal, and state Parseval's theorem for CT periodic signals

What topics did you find unclear today?

# For next time

## Content:

- The discrete-time Fourier series representation

## Action items:

1. Assignment 1 is due **tomorrow**
2. Assignment 2 (computational assignment) will be released tomorrow

## Recommended reading:

- From today's class:
  - Oppenheim 3.4-3.5,
  - Derivation of Gibbs phenomenon:  
<https://www.youtube.com/watch?v=MvXY8MXSit4>
- For next class: Oppenheim 1.3, 3.6-3.8