

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	5
Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	6
Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population.....	7
Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population.....	8
Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population	9
Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population	10
Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population	11
Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population	12
Vaccine Efficacy – First COVID-19 Occurrence After Dose 1 – Dose 1 All-Available Efficacy Population	13
Vaccine Efficacy – First COVID-19 Occurrence Based on CDC-Defined Symptoms From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population	14
Vaccine Efficacy – First COVID-19 Occurrence Based on CDC-Defined Symptoms From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population	15
Vaccine Efficacy – First COVID-19 Occurrence Based on CDC-Defined Symptoms From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population	16
Vaccine Efficacy – First COVID-19 Occurrence Based on CDC-Defined Symptoms From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population.....	17
COVID-19 Occurrence After Dose 1 Based on CDC-Defined Symptoms – Dose 1 All-Available Efficacy Population.....	18
Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	19
Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	21
Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population.....	23
Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population.....	25

Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms After Dose 1 – Dose 1 All-Available Efficacy Population	27
Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	29
Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	31
Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population.....	34
Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population	36
Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population	39
Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population	41
Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population	44
Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population	46
Vaccine Efficacy – First COVID-19 Occurrence After Dose 1, by Subgroup – Dose 1 All-Available Efficacy Population.....	49
Vaccine Efficacy – First Severe COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	52
Vaccine Efficacy – First Severe COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	53
Vaccine Efficacy – First Severe COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population.....	54
Vaccine Efficacy – First Severe COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population.....	55
Vaccine Efficacy – First Severe COVID-19 Occurrence After Dose 1 – Dose 1 All-Available Efficacy Population	56
Demographic Characteristics – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population	57
Demographic Characteristics – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population	59
Demographic Characteristics – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population	61
Demographic Characteristics – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population.....	63
Demographic Characteristics – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population.....	65
Demographic Characteristics – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population.....	67
Demographic Characteristics – Dose 1 All-Available Efficacy Population	69

Efficacy Populations.....	71
Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	73
Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	75
Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population.....	77
Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population	79
Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population	81
Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population	83
Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population	85
Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population	87
Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 1 All-Available Efficacy Population	89
Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population	91
Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population	97
Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population	103
Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population.....	109
Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population.....	115
Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population.....	121
Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population.....	127

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population	133
Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 1 All-Available Efficacy Population.....	139
Summary of Signs and Symptoms for Severe COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population.....	147
Summary of Signs and Symptoms for Severe COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population	149
Summary of Signs and Symptoms for Severe COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population	150
Summary of Signs and Symptoms for Severe COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population	152
Summary of Signs and Symptoms for Severe COVID-19 Occurrence After Dose 1 – Dose 1 All-Available Efficacy Population	153

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95 % CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =18198)		Placebo (N ^a =18325)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First COVID-19 occurrence from 7 days after Dose 2	8	2.214 (17411)	162	2.222 (17511)	95.0	(90.3, 97.6)	>0.9999

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:46)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_7pd2_wo_eval

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =19965)		Placebo (N ^a =20172)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First COVID-19 occurrence from 7 days after Dose 2	9	2.332 (18559)	169	2.345 (18708)	94.6	(89.9, 97.3)	>0.9999

Abbreviations: VE = vaccine efficacy.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:46)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_7pd2_eval

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =18175)		Placebo (N ^a =18261)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First COVID-19 occurrence from 14 days after Dose 2	8	1.887 (16612)	139	1.893 (16663)	94.2	(88.7, 97.2)	>0.9999

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:46)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_14pd2_wo_eval

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =19965)		Placebo (N ^a =20171)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First COVID-19 occurrence from 14 days after Dose 2	8	1.984 (17645)	144	1.995 (17746)	94.4	(89.1, 97.3)	>0.9999

Abbreviations: VE = vaccine efficacy.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:46)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_14pd2_eval

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =18650)		Placebo (N ^a =18570)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First COVID-19 occurrence from 7 days after Dose 2	8	2.266 (17852)	165	2.244 (17746)	95.2	(90.6, 97.7)	>0.9999

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

- a. N = number of subjects in the specified group.
- b. n1 = Number of subjects meeting the endpoint definition.
- c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.
- d. n2 = Number of subjects at risk for the endpoint.
- e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.
- f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102,1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:46)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_7pd2_wo_aai

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =20488)		Placebo (N ^a =20459)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First COVID-19 occurrence from 7 days after Dose 2	9	2.389 (19049)	172	2.370 (18971)	94.8	(90.2, 97.4)	>0.9999

Abbreviations: VE = vaccine efficacy.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:46)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_7pd2_aai

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =18627)		Placebo (N ^a =18506)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First COVID-19 occurrence from 14 days after Dose 2	8	1.931 (17032)	141	1.911 (16876)	94.4	(89.0, 97.3)	>0.9999

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:46)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_14pd2_wo_aai

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =20488)		Placebo (N ^a =20459)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First COVID-19 occurrence from 14 days after Dose 2	8	2.032 (18111)	146	2.015 (17985)	94.6	(89.4, 97.4)	>0.9999

Abbreviations: VE = vaccine efficacy.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:46)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_14pd2_aai

Vaccine Efficacy – First COVID-19 Occurrence After Dose 1 – Dose 1 All-Available Efficacy Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =21669)		Placebo (N ^a =21686)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First COVID-19 occurrence after Dose 1	50	4.015 (21314)	275	3.982 (21258)	82.0	(75.6, 86.9)
After Dose 1 to before Dose 2	39		82		52.4	(29.5, 68.4)
Dose 2 to 7 days after Dose 2	2		21		90.5	(61.0, 98.9)
≥7 Days after Dose 2	9		172		94.8	(89.8, 97.6)
Abbreviations: VE = vaccine efficacy.						
a. N = number of subjects in the specified group.						
b. n1 = Number of subjects meeting the endpoint definition.						
c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from Dose 1 to the end of the surveillance period.						
d. n2 = Number of subjects at risk for the endpoint.						
e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method (adjusted for surveillance time for overall row).						
PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 18NOV2020 (17:06)						
(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001 Efficacy FA 164/adc19ef ve cov pd1 aai						

Vaccine Efficacy – First COVID-19 Occurrence Based on CDC-Defined Symptoms From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)					
	BNT162b2 (30 µg) (N ^a =18198)			Placebo (N ^a =18325)		
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)	VE (%)	(95% CI ^e)
First COVID-19 occurrence based on CDC-defined symptoms from 7 days after Dose 2	8	2.213 (17399)	165	2.220 (17495)	95.1	(90.2, 97.9)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 18NOV2020 (07:39)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_7pd2_wo_cdc_eval

Vaccine Efficacy – First COVID-19 Occurrence Based on CDC-Defined Symptoms From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population						
Efficacy Endpoint	Vaccine Group (as Randomized)					
	BNT162b2 (30 µg) (N ^a =19965)			Placebo (N ^a =20172)		
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)	VE (%)	(95% CI ^e)
First COVID-19 occurrence based on CDC-defined symptoms from 7 days after Dose 2	9	2.330 (18544)	172	2.343 (18690)	94.7	(89.8, 97.6)
Abbreviations: VE = vaccine efficacy. a. N = number of subjects in the specified group. b. n1 = Number of subjects meeting the endpoint definition. c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period. d. n2 = Number of subjects at risk for the endpoint. e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 18NOV2020 (07:39) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_7pd2_cdc_eval						

Vaccine Efficacy – First COVID-19 Occurrence Based on CDC-Defined Symptoms From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)					
	BNT162b2 (30 µg) (N ^a =18175)		Placebo (N ^a =18261)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)	VE (%)	(95% CI ^e)
First COVID-19 occurrence based on CDC-defined symptoms from 14 days after Dose 2	8	1.886 (16600)	141	1.891 (16647)	94.3	(88.5, 97.6)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 18NOV2020 (07:39)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_14pd2_wo_cdc_eval

Vaccine Efficacy – First COVID-19 Occurrence Based on CDC-Defined Symptoms From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)					
	BNT162b2 (30 µg) (N ^a =19965)			Placebo (N ^a =20171)		
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)	VE (%)	(95% CI ^e)
First COVID-19 occurrence based on CDC-defined symptoms from 14 days after Dose 2	8	1.983 (17630)	146	1.993 (17727)	94.5	(88.9, 97.7)

Abbreviations: VE = vaccine efficacy.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 18NOV2020 (07:39)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_14pd2_cdc_eval

COVID-19 Occurrence After Dose 1 Based on CDC-Defined Symptoms – Dose 1 All-Available Efficacy Population		
Efficacy Endpoint	Vaccine Group (as Randomized)	
	BNT162b2 (30 µg) (N^a=21669) n^b	Placebo (N^a=21686) n^b
COVID-19 occurrence after Dose 1 based on CDC-defined symptoms	54	282
After Dose 1 and before Dose 2	42	85
Dose 2 to 7 days after Dose 2	3	22
≥7 days after Dose 2	9	175
a. N = number of subjects in the specified group. b. n = Number of subjects meeting the endpoint definition. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 18NOV2020 (18:59) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_pd1_cdc_aai		

**Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 7 Days After Dose 2 –
Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population**

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =165)	Total (N ^a =173)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (25.0)	76 (46.1)	78 (45.1)
New or increased cough	3 (37.5)	114 (69.1)	117 (67.6)
New or increased shortness of breath	0 (0.0)	25 (15.2)	25 (14.5)
Chills	2 (25.0)	57 (34.5)	59 (34.1)
New or increased muscle pain	1 (12.5)	81 (49.1)	82 (47.4)
New loss of taste or smell	5 (62.5)	43 (26.1)	48 (27.7)
Sore throat	3 (37.5)	68 (41.2)	71 (41.0)
Diarrhea	1 (12.5)	18 (10.9)	19 (11.0)
Vomiting	2 (25.0)	6 (3.6)	8 (4.6)
Additional CDC-defined symptoms			
Fatigue	0 (0.0)	44 (26.7)	44 (25.4)
Headache	2 (25.0)	63 (38.2)	65 (37.6)
Nasal congestion	1 (12.5)	33 (20.0)	34 (19.7)
Runny nose	0 (0.0)	21 (12.7)	21 (12.1)
Nausea	1 (12.5)	3 (1.8)	4 (2.3)

Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =165)	Total (N ^a =173)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence based on CDC-defined symptoms from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:50)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_cdc_7d2_wo_eval

Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =9)	Placebo (N ^a =172)	Total (N ^a =181)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (22.2)	77 (44.8)	79 (43.6)
New or increased cough	3 (33.3)	120 (69.8)	123 (68.0)
New or increased shortness of breath	0 (0.0)	26 (15.1)	26 (14.4)
Chills	2 (22.2)	58 (33.7)	60 (33.1)
New or increased muscle pain	1 (11.1)	82 (47.7)	83 (45.9)
New loss of taste or smell	5 (55.6)	46 (26.7)	51 (28.2)
Sore throat	4 (44.4)	71 (41.3)	75 (41.4)
Diarrhea	1 (11.1)	19 (11.0)	20 (11.0)
Vomiting	2 (22.2)	6 (3.5)	8 (4.4)
Additional CDC-defined symptoms			
Fatigue	0 (0.0)	45 (26.2)	45 (24.9)
Headache	2 (22.2)	65 (37.8)	67 (37.0)
Nasal congestion	1 (11.1)	36 (20.9)	37 (20.4)
Runny nose	0 (0.0)	22 (12.8)	22 (12.2)
Nausea	1 (11.1)	3 (1.7)	4 (2.2)

Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =9)	Placebo (N ^a =172)	Total (N ^a =181)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

a. N = number of subjects with COVID-19 occurrence based on CDC-defined symptoms from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:50)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_cdc_7d2_eval

Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =141)	Total (N ^a =149)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (25.0)	65 (46.1)	67 (45.0)
New or increased cough	3 (37.5)	101 (71.6)	104 (69.8)
New or increased shortness of breath	0 (0.0)	20 (14.2)	20 (13.4)
Chills	2 (25.0)	48 (34.0)	50 (33.6)
New or increased muscle pain	1 (12.5)	67 (47.5)	68 (45.6)
New loss of taste or smell	5 (62.5)	36 (25.5)	41 (27.5)
Sore throat	3 (37.5)	57 (40.4)	60 (40.3)
Diarrhea	1 (12.5)	17 (12.1)	18 (12.1)
Vomiting	2 (25.0)	5 (3.5)	7 (4.7)
Additional CDC-defined symptoms			
Fatigue	0 (0.0)	36 (25.5)	36 (24.2)
Headache	2 (25.0)	54 (38.3)	56 (37.6)
Nasal congestion	1 (12.5)	29 (20.6)	30 (20.1)
Runny nose	0 (0.0)	18 (12.8)	18 (12.1)
Nausea	1 (12.5)	3 (2.1)	4 (2.7)

Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =141)	Total (N ^a =149)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence based on CDC-defined symptoms from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:50)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_cdc_14d2_wo_eva

Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =146)	Total (N ^a =154)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (25.0)	66 (45.2)	68 (44.2)
New or increased cough	3 (37.5)	105 (71.9)	108 (70.1)
New or increased shortness of breath	0 (0.0)	21 (14.4)	21 (13.6)
Chills	2 (25.0)	48 (32.9)	50 (32.5)
New or increased muscle pain	1 (12.5)	68 (46.6)	69 (44.8)
New loss of taste or smell	5 (62.5)	38 (26.0)	43 (27.9)
Sore throat	3 (37.5)	60 (41.1)	63 (40.9)
Diarrhea	1 (12.5)	17 (11.6)	18 (11.7)
Vomiting	2 (25.0)	5 (3.4)	7 (4.5)
Additional CDC-defined symptoms			
Fatigue	0 (0.0)	37 (25.3)	37 (24.0)
Headache	2 (25.0)	56 (38.4)	58 (37.7)
Nasal congestion	1 (12.5)	32 (21.9)	33 (21.4)
Runny nose	0 (0.0)	19 (13.0)	19 (12.3)
Nausea	1 (12.5)	3 (2.1)	4 (2.6)

Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =146)	Total (N ^a =154)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

a. N = number of subjects with COVID-19 occurrence based on CDC-defined symptoms from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:50)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_cdc_14d2_eval

**Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms After Dose 1 –
Dose 1 All-Available Efficacy Population**

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =54)	Placebo (N ^a =282)	Total (N ^a =336)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	20 (37.0)	122 (43.3)	142 (42.3)
New or increased cough	22 (40.7)	186 (66.0)	208 (61.9)
New or increased shortness of breath	4 (7.4)	44 (15.6)	48 (14.3)
Chills	10 (18.5)	86 (30.5)	96 (28.6)
New or increased muscle pain	12 (22.2)	121 (42.9)	133 (39.6)
New loss of taste or smell	24 (44.4)	91 (32.3)	115 (34.2)
Sore throat	18 (33.3)	111 (39.4)	129 (38.4)
Diarrhea	4 (7.4)	35 (12.4)	39 (11.6)
Vomiting	5 (9.3)	11 (3.9)	16 (4.8)
Additional CDC-defined symptoms			
Fatigue	6 (11.1)	65 (23.0)	71 (21.1)
Headache	14 (25.9)	96 (34.0)	110 (32.7)
Nasal congestion	7 (13.0)	56 (19.9)	63 (18.8)
Runny nose	7 (13.0)	31 (11.0)	38 (11.3)
Nausea	2 (3.7)	8 (2.8)	10 (3.0)

a. N = number of subjects with COVID-19 occurrence based on CDC-defined symptoms after dose 1 in the specified group. This value is used as the

Summary of Signs and Symptoms for COVID-19 Occurrence Based on CDC-Defined Symptoms After Dose 1 – Dose 1 All-Available Efficacy Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =54)	Placebo (N ^a =282)	Total (N ^a =336)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:50)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_cdc_d1_aai

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =18198)		Placebo (N ^a =18325)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First COVID-19 occurrence from 7 days after Dose 2						
Overall	8	2.214 (17411)	162	2.222 (17511)	95.0	(90.0, 97.9)
Age group (years)						
16 to 55	5	1.234 (9897)	114	1.239 (9955)	95.6	(89.4, 98.6)
>55	3	0.980 (7500)	48	0.983 (7543)	93.7	(80.6, 98.8)
≥65	1	0.508 (3848)	19	0.511 (3880)	94.7	(66.7, 99.9)
Sex						
Male	3	1.124 (8875)	81	1.108 (8762)	96.4	(88.9, 99.3)
Female	5	1.090 (8536)	81	1.114 (8749)	93.7	(84.7, 98.0)
Race						
White	7	1.889 (14504)	146	1.903 (14670)	95.2	(89.8, 98.1)
Black or African American	0	0.165 (1502)	7	0.164 (1486)	100.0	(31.2, 100.0)
All others ^f	1	0.160 (1405)	9	0.155 (1355)	89.3	(22.6, 99.8)
Ethnicity						

090177e195901438\Final\Final On: 19-Nov-2020 06:23 (GMT)

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =18198)		Placebo (N ^a =18325)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
Hispanic/Latino	3	0.605 (4764)	53	0.600 (4746)	94.4	(82.7, 98.9)
Non-Hispanic/non-Latino	5	1.596 (12548)	109	1.608 (12661)	95.4	(88.9, 98.5)
Country						
Argentina	1	0.351 (2545)	35	0.346 (2521)	97.2	(83.3, 99.9)
Brazil	1	0.119 (1129)	8	0.117 (1121)	87.7	(8.1, 99.7)
USA	6	1.732 (13359)	119	1.747 (13506)	94.9	(88.6, 98.2)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.

f. All others = American Indian or Alaska native, Asian, Native Hawaiian or other Pacific Islander, multiracial, and not reported race categories.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 18NOV2020 (15:55)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_7pd2_wo_sg_eval

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =19965)		Placebo (N ^a =20172)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First COVID-19 occurrence from 7 days after Dose 2						
Overall	9	2.332 (18559)	169	2.345 (18708)	94.6	(89.6, 97.6)
Age group (years)						
16 to 55	6	1.309 (10653)	120	1.317 (10738)	95.0	(88.7, 98.2)
>55	3	1.022 (7892)	49	1.028 (7956)	93.8	(80.9, 98.8)
≥65	1	0.530 (4044)	19	0.532 (4067)	94.7	(66.8, 99.9)
Sex						
Male	4	1.183 (9457)	85	1.170 (9342)	95.3	(87.6, 98.8)
Female	5	1.149 (9102)	84	1.176 (9366)	93.9	(85.2, 98.1)
Race						
White	7	1.975 (15294)	153	1.990 (15473)	95.4	(90.3, 98.2)
Black or African American	0	0.187 (1758)	7	0.188 (1758)	100.0	(30.4, 100.0)
All others ^f	2	0.170 (1507)	9	0.167 (1477)	78.2	(-5.4, 97.7)
Ethnicity						

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)					
	BNT162b2 (30 µg) (N ^a =19965)		Placebo (N ^a =20172)		VE (%)	(95% CI ^e)
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
Hispanic/Latino	3	0.637 (5074)	55	0.638 (5090)	94.5	(83.2, 98.9)
Non-Hispanic/non-Latino	6	1.681 (13380)	114	1.693 (13509)	94.7	(88.1, 98.1)
Country						
Argentina	1	0.366 (2664)	36	0.367 (2684)	97.2	(83.5, 99.9)
Brazil	2	0.134 (1274)	8	0.132 (1257)	75.4	(-23.5, 97.5)
USA	6	1.816 (14141)	124	1.830 (14287)	95.1	(89.1, 98.2)
South Africa	0	0.015 (362)	1	0.015 (363)	100.0	(-3818.9, 100.0)
Prior SARS-CoV-2 Status						
Positive at baseline ^g	1	0.056 (526)	1	0.060 (567)	-7.1	(-8309.9, 98.6)
Negative at baseline but positive prior to 7 days after Dose 2 ^h	0	0.003 (27)	1	0.004 (34)	100.0	(-6004.9, 100.0)
Negative prior to 7 days after Dose 2 ⁱ	8	2.214 (17411)	162	2.222 (17511)	95.0	(90.0, 97.9)
Unknown	0	0.059 (595)	5	0.060 (596)	100.0	(-9.6, 100.0)
Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.						
a. N = number of subjects in the specified group.						
b. n1 = Number of subjects meeting the endpoint definition.						
c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.						
d. n2 = Number of subjects at risk for the endpoint.						

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =19965)		Placebo (N ^a =20172)			
	Surveillance		Surveillance			
	n1 ^b	Time ^c (n2 ^d)	n1 ^b	Time ^c (n2 ^d)		
e.	Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.					
f.	All others = American Indian or Alaska native, Asian, Native Hawaiian or other Pacific Islander, multiracial, and not reported race categories.					
g.	Positive N-binding antibody result at Visit 1, positive NAAT result at Visit 1, or medical history of COVID-19.					
h.	Negative N-binding antibody result and negative NAAT result at Visit 1, positive NAAT result at Visit 2 or at unscheduled visit, if any, prior to 7 days after Dose 2.					
i.	Negative N-binding antibody result at Visit 1, negative NAAT result at Visit 1 and Visit 2, and negative NAAT result at unscheduled visit, if any, prior to 7 days after Dose 2.					
PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 18NOV2020 (15:55) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_7pd2_sg_eval						

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =18175)		Placebo (N ^a =18261)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First COVID-19 occurrence from 14 days after Dose 2						
Overall	8	1.887 (16612)	139	1.893 (16663)	94.2	(88.3, 97.6)
Age group (years)						
16 to 55	5	1.048 (9440)	96	1.052 (9453)	94.8	(87.4, 98.3)
>55	3	0.839 (7172)	43	0.841 (7210)	93.0	(78.2, 98.6)
≥65	1	0.436 (3663)	17	0.438 (3709)	94.1	(62.3, 99.9)
Sex						
Male	3	0.958 (8467)	68	0.944 (8334)	95.7	(86.7, 99.1)
Female	5	0.930 (8145)	71	0.950 (8329)	92.8	(82.4, 97.7)
Race						
White	7	1.616 (13908)	124	1.628 (14025)	94.3	(87.9, 97.8)
Black or African American	0	0.138 (1378)	7	0.136 (1370)	100.0	(31.2, 100.0)
All others ^f	1	0.134 (1326)	8	0.129 (1268)	87.9	(10.1, 99.7)
Ethnicity						

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =18175)		Placebo (N ^a =18261)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
Hispanic/Latino	3	0.515 (4613)	39	0.511 (4575)	92.4	(76.0, 98.5)
Non-Hispanic/non-Latino	5	1.361 (11901)	100	1.371 (11988)	95.0	(87.9, 98.4)
Country						
Argentina	1	0.303 (2545)	25	0.298 (2510)	96.1	(76.0, 99.9)
Brazil	1	0.098 (994)	4	0.097 (974)	75.3	(-149.2, 99.5)
USA	6	1.480 (12815)	110	1.492 (12931)	94.5	(87.6, 98.0)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

- N = number of subjects in the specified group.
- n1 = Number of subjects meeting the endpoint definition.
- Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.
- n2 = Number of subjects at risk for the endpoint.
- Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.
- All others = American Indian or Alaska native, Asian, Native Hawaiian or other Pacific Islander, multiracial, and not reported race categories.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 18NOV2020 (15:58)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_14pd2_wo_sg_eval

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =19965)		Placebo (N ^a =20171)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First COVID-19 occurrence from 14 days after Dose 2						
Overall	8	1.984 (17645)	144	1.995 (17746)	94.4	(88.7, 97.6)
Age group (years)						
16 to 55	5	1.110 (10114)	100	1.117 (10157)	95.0	(87.9, 98.4)
>55	3	0.874 (7531)	44	0.879 (7589)	93.1	(78.6, 98.6)
≥65	1	0.455 (3842)	17	0.455 (3881)	94.1	(62.4, 99.9)
Sex						
Male	3	1.006 (8991)	71	0.995 (8860)	95.8	(87.3, 99.2)
Female	5	0.979 (8654)	73	1.000 (8886)	93.0	(82.9, 97.8)
Race						
White	7	1.688 (14621)	129	1.700 (14750)	94.5	(88.4, 97.8)
Black or African American	0	0.155 (1605)	7	0.155 (1616)	100.0	(30.4, 100.0)
All others ^f	1	0.142 (1419)	8	0.140 (1380)	87.7	(8.4, 99.7)
Ethnicity						

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =19965)		Placebo (N ^a =20171)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
Hispanic/Latino	3	0.542 (4896)	41	0.542 (4900)	92.7	(77.0, 98.5)
Non-Hispanic/non-Latino	5	1.431 (12645)	103	1.440 (12741)	95.1	(88.2, 98.4)
Country						
Argentina	1	0.315 (2664)	26	0.315 (2673)	96.2	(76.6, 99.9)
Brazil	1	0.111 (1122)	4	0.110 (1101)	75.3	(-149.2, 99.5)
USA	6	1.549 (13518)	114	1.562 (13631)	94.7	(88.1, 98.1)
Prior SARS-CoV-2 Status						
Positive at baseline ^g	0	0.046 (486)	1	0.050 (527)	100.0	(-4066.9, 100.0)
Negative prior to 14 days after Dose 2 ⁱ	8	1.887 (16612)	139	1.893 (16663)	94.2	(88.3, 97.6)
Unknown	0	0.049 (522)	4	0.049 (523)	100.0	(-53.1, 100.0)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

- N = number of subjects in the specified group.
- n1 = Number of subjects meeting the endpoint definition.
- Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.
- n2 = Number of subjects at risk for the endpoint.
- Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.
- All others = American Indian or Alaska native, Asian, Native Hawaiian or other Pacific Islander, multiracial, and not reported race categories.
- Positive N-binding antibody result at Visit 1, positive NAAT result at Visit 1, or medical history of COVID-19.

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)					
	BNT162b2 (30 µg) (N ^a =19965)		Placebo (N ^a =20171)		VE (%)	(95% CI ^e)
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
h. Negative N-binding antibody result and negative NAAT result at Visit 1, positive NAAT result at Visit 2 or at unscheduled visit, if any, prior to 7 days after Dose 2.						
i. Negative N-binding antibody result at Visit 1, negative NAAT result at Visit 1 and Visit 2, and negative NAAT result at unscheduled visit, if any, prior to 7 days after Dose 2.						

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 18NOV2020 (15:58)
(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_14pd2_sg_eval

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =18650)		Placebo (N ^a =18570)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First COVID-19 occurrence from 7 days after Dose 2						
Overall	8	2.266 (17852)	165	2.244 (17746)	95.2	(90.3, 98.0)
Age group (years)						
16 to 55	5	1.262 (10145)	117	1.251 (10087)	95.8	(89.8, 98.6)
>55	3	1.004 (7693)	48	0.993 (7646)	93.8	(80.8, 98.8)
≥65	1	0.523 (3965)	19	0.516 (3936)	94.8	(67.3, 99.9)
Sex						
Male	3	1.150 (9100)	84	1.120 (8886)	96.5	(89.5, 99.3)
Female	5	1.116 (8752)	81	1.124 (8860)	93.8	(84.9, 98.0)
Race						
White	7	1.937 (14912)	149	1.922 (14879)	95.3	(90.2, 98.2)
Black or African American	0	0.168 (1525)	7	0.165 (1495)	100.0	(31.8, 100.0)
All others ^f	1	0.162 (1415)	9	0.157 (1372)	89.2	(22.3, 99.8)
Ethnicity						

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =18650)		Placebo (N ^a =18570)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
Hispanic/Latino	3	0.615 (4882)	54	0.608 (4849)	94.5	(83.1, 98.9)
Non-Hispanic/non-Latino	5	1.637 (12866)	111	1.623 (12793)	95.5	(89.3, 98.6)
Country						
Argentina	1	0.358 (2628)	36	0.351 (2599)	97.3	(83.8, 99.9)
Brazil	1	0.119 (1130)	8	0.117 (1123)	87.7	(8.0, 99.7)
USA	6	1.777 (13709)	121	1.764 (13656)	95.1	(89.0, 98.2)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

- N = number of subjects in the specified group.
- n1 = Number of subjects meeting the endpoint definition.
- Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.
- n2 = Number of subjects at risk for the endpoint.
- Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.
- All others = American Indian or Alaska native, Asian, Native Hawaiian or other Pacific Islander, multiracial, and not reported race categories.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 18NOV2020 (15:58)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_7pd2_wo_sg_aai

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =20488)		Placebo (N ^a =20459)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First COVID-19 occurrence from 7 days after Dose 2						
Overall	9	2.389 (19049)	172	2.370 (18971)	94.8	(89.9, 97.7)
Age group (years)						
16 to 55	6	1.341 (10931)	123	1.331 (10890)	95.2	(89.2, 98.3)
>55	3	1.048 (8104)	49	1.039 (8067)	93.9	(81.2, 98.8)
≥65	1	0.545 (4168)	19	0.537 (4126)	94.8	(67.4, 99.9)
Sex						
Male	4	1.212 (9717)	88	1.183 (9477)	95.6	(88.2, 98.8)
Female	5	1.176 (9332)	84	1.187 (9494)	94.0	(85.4, 98.1)
Race						
White	7	2.027 (15749)	156	2.012 (15709)	95.5	(90.6, 98.2)
Black or African American	0	0.189 (1781)	7	0.189 (1768)	100.0	(30.8, 100.0)
All others ^f	2	0.172 (1519)	9	0.169 (1494)	78.1	(-5.6, 97.7)
Ethnicity						

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =20488)		Placebo (N ^a =20459)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
Hispanic/Latino	3	0.649 (5213)	56	0.647 (5214)	94.7	(83.6, 98.9)
Non-Hispanic/non-Latino	6	1.725 (13726)	116	1.708 (13648)	94.9	(88.5, 98.2)
Country						
Argentina	1	0.375 (2765)	37	0.373 (2776)	97.3	(84.1, 99.9)
Brazil	2	0.134 (1277)	8	0.132 (1259)	75.4	(-23.5, 97.5)
USA	6	1.863 (14520)	126	1.849 (14451)	95.3	(89.4, 98.3)
South Africa	0	0.015 (363)	1	0.015 (363)	100.0	(-3802.8, 100.0)
Prior SARS-CoV-2 Status						
Positive at baseline ^g	1	0.058 (541)	1	0.061 (579)	-6.4	(-8250.5, 98.6)
Negative at baseline but positive prior to 7 days after Dose 2 ^h	0	0.003 (29)	1	0.004 (35)	100.0	(-5553.6, 100.0)
Negative prior to 7 days after Dose 2 ⁱ	8	2.266 (17852)	165	2.244 (17746)	95.2	(90.3, 98.0)
Unknown	0	0.063 (627)	5	0.061 (611)	100.0	(-5.9, 100.0)
Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.						
a. N = number of subjects in the specified group.						
b. n1 = Number of subjects meeting the endpoint definition.						
c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.						
d. n2 = Number of subjects at risk for the endpoint.						

Vaccine Efficacy – First COVID-19 Occurrence From 7 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =20488)		Placebo (N ^a =20459)			
	Surveillance		Surveillance			
	n1 ^b	Time ^c (n2 ^d)	n1 ^b	Time ^c (n2 ^d)		
e.	Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.					
f.	All others = American Indian or Alaska native, Asian, Native Hawaiian or other Pacific Islander, multiracial, and not reported race categories.					
g.	Positive N-binding antibody result at Visit 1, positive NAAT result at Visit 1, or medical history of COVID-19.					
h.	Negative N-binding antibody result and negative NAAT result at Visit 1, positive NAAT result at Visit 2 or at unscheduled visit, if any, prior to 7 days after Dose 2.					
i.	Negative N-binding antibody result at Visit 1, negative NAAT result at Visit 1 and Visit 2, and negative NAAT result at unscheduled visit, if any, prior to 7 days after Dose 2.					
PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 18NOV2020 (15:58) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_7pd2_sg_aai						

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =18627)		Placebo (N ^a =18506)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First COVID-19 occurrence from 14 days after Dose 2						
Overall	8	1.931 (17032)	141	1.911 (16876)	94.4	(88.6, 97.6)
Age group (years)						
16 to 55	5	1.072 (9674)	98	1.062 (9572)	94.9	(87.8, 98.4)
>55	3	0.859 (7358)	43	0.849 (7304)	93.1	(78.5, 98.6)
≥65	1	0.448 (3776)	17	0.442 (3759)	94.2	(63.0, 99.9)
Sex						
Male	3	0.980 (8680)	70	0.954 (8447)	95.8	(87.3, 99.2)
Female	5	0.951 (8352)	71	0.957 (8429)	92.9	(82.7, 97.8)
Race						
White	7	1.656 (14298)	126	1.643 (14213)	94.5	(88.3, 97.8)
Black or African American	0	0.139 (1398)	7	0.137 (1378)	100.0	(31.7, 100.0)
All others ^f	1	0.135 (1336)	8	0.131 (1285)	87.9	(9.7, 99.7)
Ethnicity						

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =18627)		Placebo (N ^a =18506)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
Hispanic/Latino	3	0.523 (4726)	39	0.516 (4672)	92.4	(76.1, 98.5)
Non-Hispanic/non-Latino	5	1.397 (12203)	102	1.383 (12104)	95.1	(88.3, 98.5)
Country						
Argentina	1	0.308 (2627)	25	0.301 (2585)	96.1	(76.1, 99.9)
Brazil	1	0.098 (994)	4	0.097 (976)	75.3	(-149.4, 99.5)
USA	6	1.518 (13152)	112	1.507 (13067)	94.7	(88.1, 98.1)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

- N = number of subjects in the specified group.
- n1 = Number of subjects meeting the endpoint definition.
- Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.
- n2 = Number of subjects at risk for the endpoint.
- Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.
- All others = American Indian or Alaska native, Asian, Native Hawaiian or other Pacific Islander, multiracial, and not reported race categories.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 18NOV2020 (15:58)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_14pd2_wo_sg_aai

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =20488)		Placebo (N ^a =20459)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First COVID-19 occurrence from 14 days after Dose 2						
Overall	8	2.032 (18111)	146	2.015 (17985)	94.6	(89.0, 97.7)
Age group (years)						
16 to 55	5	1.136 (10376)	102	1.127 (10294)	95.1	(88.3, 98.5)
>55	3	0.896 (7735)	44	0.888 (7691)	93.2	(78.9, 98.7)
≥65	1	0.467 (3962)	17	0.459 (3934)	94.2	(63.1, 99.9)
Sex						
Male	3	1.030 (9238)	73	1.005 (8983)	96.0	(87.8, 99.2)
Female	5	1.001 (8873)	73	1.010 (9002)	93.1	(83.1, 97.8)
Race						
White	7	1.732 (15055)	131	1.717 (14963)	94.7	(88.8, 97.9)
Black or African American	0	0.157 (1625)	7	0.156 (1625)	100.0	(30.8, 100.0)
All others ^f	1	0.143 (1431)	8	0.141 (1397)	87.7	(8.2, 99.7)
Ethnicity						

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =20488)		Placebo (N ^a =20459)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
Hispanic/Latino	3	0.551 (5030)	41	0.549 (5016)	92.7	(77.2, 98.6)
Non-Hispanic/non-Latino	5	1.469 (12972)	105	1.454 (12864)	95.3	(88.6, 98.5)
Country						
Argentina	1	0.322 (2764)	26	0.320 (2762)	96.2	(76.8, 99.9)
Brazil	1	0.111 (1123)	4	0.110 (1103)	75.3	(-149.2, 99.5)
USA	6	1.590 (13882)	116	1.577 (13779)	94.9	(88.5, 98.2)
Prior SARS-CoV-2 Status						
Positive at baseline ^g	0	0.047 (500)	1	0.050 (537)	100.0	(-4038.2, 100.0)
Negative prior to 14 days after Dose 2 ⁱ	8	1.931 (17032)	141	1.911 (16876)	94.4	(88.6, 97.6)
Unknown	0	0.051 (552)	4	0.050 (538)	100.0	(-47.7, 100.0)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

- N = number of subjects in the specified group.
- n1 = Number of subjects meeting the endpoint definition.
- Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.
- n2 = Number of subjects at risk for the endpoint.
- Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time.
- All others = American Indian or Alaska native, Asian, Native Hawaiian or other Pacific Islander, multiracial, and not reported race categories.
- Positive N-binding antibody result at Visit 1, positive NAAT result at Visit 1, or medical history of COVID-19.

Vaccine Efficacy – First COVID-19 Occurrence From 14 Days After Dose 2, by Subgroup – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =20488)		Placebo (N ^a =20459)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
h. Negative N-binding antibody result and negative NAAT result at Visit 1, positive NAAT result at Visit 2 or at unscheduled visit, if any, prior to 7 days after Dose 2.						
i. Negative N-binding antibody result at Visit 1, negative NAAT result at Visit 1 and Visit 2, and negative NAAT result at unscheduled visit, if any, prior to 7 days after Dose 2.						
PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 18NOV2020 (16:00) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_14pd2_sg_aai						

Vaccine Efficacy – First COVID-19 Occurrence After Dose 1, by Subgroup – Dose 1 All-Available Efficacy Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =21669)		Placebo (N ^a =21686)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First COVID-19 occurrence after Dose 1	50	4.015 (21314)	275	3.982 (21258)	82.0	(75.6, 86.9)
Age group (years)						
12 to 15	1	0.004 (49)	2	0.004 (51)	48.3	(-893.5, 99.1)
16 to 55	38	2.294 (12488)	200	2.274 (12469)	81.2	(73.2, 87.0)
>55	11	1.718 (8777)	73	1.704 (8738)	85.1	(71.7, 92.9)
≥65	6	0.887 (4467)	24	0.877 (4461)	75.3	(37.9, 91.7)
Sex						
Male	25	2.047 (10968)	149	1.992 (10702)	83.7	(74.9, 89.8)
Female	25	1.969 (10346)	126	1.990 (10556)	79.9	(69.0, 87.5)
Race						
White	45	3.366 (17544)	250	3.338 (17508)	82.1	(75.4, 87.3)
Black or African American	2	0.345 (2023)	10	0.344 (2024)	80.0	(6.4, 97.9)
All others ^c	3	0.304 (1747)	15	0.299 (1726)	80.3	(30.3, 96.3)
Ethnicity						

Vaccine Efficacy – First COVID-19 Occurrence After Dose 1, by Subgroup – Dose 1 All-Available Efficacy Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =21669)		Placebo (N ^a =21686)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
Hispanic/Latino	21	1.085 (5590)	101	1.076 (5582)	79.4	(66.8, 87.8)
Non-Hispanic/non-Latino	29	2.907 (15606)	172	2.883 (15558)	83.3	(75.1, 89.1)
Not reported	0	0.023 (118)	2	0.023 (118)	100.0	(-441.9, 100.0)
Country						
Argentina	16	0.600 (2856)	70	0.592 (2855)	77.5	(60.8, 87.8)
Brazil	2	0.244 (1439)	18	0.241 (1425)	89.0	(54.2, 98.8)
Germany	2	0.018 (246)	0	0.018 (249)	-∞	(NA, NA)
South Africa	0	0.044 (372)	2	0.044 (368)	100.0	(-427.5, 100.0)
Turkey	2	0.009 (245)	0	0.009 (240)	-∞	(NA, NA)
USA	28	3.099 (16156)	185	3.079 (16121)	85.0	(77.5, 90.3)
Baseline SARS-CoV-2 status						
Positive ^d	10	0.106 (633)	9	0.113 (670)	-17.9	(-227.9, 56.9)
Negative ^e	36	3.814 (19913)	259	3.777 (19818)	86.2	(80.4, 90.6)
Unknown	4	0.095 (768)	7	0.093 (770)	44.2	(-119.6, 88.0)
Abbreviations: N-binding = SARS-CoV-2 nucleoprotein–binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.						

Vaccine Efficacy – First COVID-19 Occurrence After Dose 1, by Subgroup – Dose 1 All-Available Efficacy Population

Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =21669)		Placebo (N ^a =21686)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
a. N = number of subjects in the specified group. b. n1 = Number of subjects meeting the endpoint definition. c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from Dose 1 to the end of the surveillance period. d. n2 = Number of subjects at risk for the endpoint. e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method adjusted for surveillance time. f. All others = American Indian or Alaska native, Asian, Native Hawaiian or other Pacific Islander, multiracial, and not reported race categories. g. Positive N-binding antibody result at Visit 1, positive NAAT result at Visit 1, or medical history of COVID-19. h. Negative N-binding antibody result at Visit 1, negative NAAT result at Visit 1, and no medical history of COVID-19. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 18NOV2020 (17:11) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_cov_pd1_sg_aai						

Vaccine Efficacy – First Severe COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =18198)		Placebo (N ^a =18325)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First severe COVID-19 occurrence from 7 days after Dose 2	1	2.215 (17411)	3	2.232 (17511)	66.4	(-124.8, 96.3)	0.7429

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

- a. N = number of subjects in the specified group.
- b. n1 = Number of subjects meeting the endpoint definition.
- c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.
- d. n2 = Number of subjects at risk for the endpoint.
- e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.
- f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:47)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_sev_cov_7pd2_wo_eval

Vaccine Efficacy – First Severe COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI) ^e	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =19965)		Placebo (N ^a =20172)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First severe COVID-19 occurrence from 7 days after Dose 2	1	2.333 (18566)	3	2.358 (18733)	66.3	(-125.5, 96.3)	0.7419

Abbreviations: VE = vaccine efficacy.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 7 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:46)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_sev_cov_7pd2_eval

Vaccine Efficacy – First Severe COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =18175)		Placebo (N ^a =18261)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First severe COVID-19 occurrence from 14 days after Dose 2	1	1.888 (16612)	3	1.901 (16663)	66.4	(-124.7, 96.3)	0.7432

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; VE = vaccine efficacy.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:46)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_sev_cov_14pd2_wo_eval

Vaccine Efficacy – First Severe COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Efficacy Endpoint	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)	Pr (VE >30% data) ^f
	BNT162b2 (30 µg) (N ^a =19965)		Placebo (N ^a =20171)				
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)			
First severe COVID-19 occurrence from 14 days after Dose 2	1	1.985 (17652)	3	2.007 (17792)	66.3	(-125.6, 96.3)	0.7418

Abbreviations: VE = vaccine efficacy.

a. N = number of subjects in the specified group.

b. n1 = Number of subjects meeting the endpoint definition.

c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from 14 days after Dose 2 to the end of the surveillance period.

d. n2 = Number of subjects at risk for the endpoint.

e. Credible interval for VE was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

f. Posterior probability (Pr) was calculated using a beta-binomial model with prior beta (0.700102, 1) adjusted for surveillance time. Refer to the statistical analysis plan, Appendix 2, for more details.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 17NOV2020 (16:47)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adc19ef_ve_sev_cov_14pd2_eval

Vaccine Efficacy – First Severe COVID-19 Occurrence After Dose 1 – Dose 1 All-Available Efficacy Population						
Efficacy Endpoint Subgroup	Vaccine Group (as Randomized)				VE (%)	(95% CI ^e)
	BNT162b2 (30 µg) (N ^a =21669)		Placebo (N ^a =21686)			
	n1 ^b	Surveillance Time ^c (n2 ^d)	n1 ^b	Surveillance Time ^c (n2 ^d)		
First severe COVID-19 occurrence after Dose 1	1	4.021 (21314)	9	4.006 (21259)	88.9	(20.1, 99.7)
After Dose 1 to before Dose 2	0		4		100.0	(-51.5, 100.0)
Dose 2 to 7 days after Dose 2	0		1		100.0	(-3800.0, 100.0)
≥7 Days after Dose 2	1		4		75.0	(-152.6, 99.5)
Abbreviations: VE = vaccine efficacy.						
a. N = number of subjects in the specified group.						
b. n1 = Number of subjects meeting the endpoint definition.						
c. Total surveillance time in 1000 person-years for the given endpoint across all subjects within each group at risk for the endpoint. Time period for COVID-19 case accrual is from Dose 1 to the end of the surveillance period.						
d. n2 = Number of subjects at risk for the endpoint.						
e. Confidence interval (CI) for VE is derived based on the Clopper and Pearson method (adjusted for surveillance time for overall row).						
PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:48) Source Data: adc19ef Table Generation: 18NOV2020 (17:43)						
(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001 Efficacy FA 164/adc19ef ve sev cov pd1 aai						

Demographic Characteristics – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population			
	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N^a=18242) n^b (%)	Placebo (N^a=18379) n^b (%)	Total (N^a=36621) n^b (%)
Sex			
Male	9318 (51.1)	9225 (50.2)	18543 (50.6)
Female	8924 (48.9)	9154 (49.8)	18078 (49.4)
Race			
White	15110 (82.8)	15301 (83.3)	30411 (83.0)
Black or African American	1617 (8.9)	1617 (8.8)	3234 (8.8)
American Indian or Alaska native	118 (0.6)	106 (0.6)	224 (0.6)
Asian	815 (4.5)	810 (4.4)	1625 (4.4)
Native Hawaiian or other Pacific Islander	48 (0.3)	29 (0.2)	77 (0.2)
Multiracial	448 (2.5)	402 (2.2)	850 (2.3)
Not reported	86 (0.5)	114 (0.6)	200 (0.5)
Ethnicity			
Hispanic/Latino	4886 (26.8)	4857 (26.4)	9743 (26.6)
Non-Hispanic/non-Latino	13253 (72.7)	13412 (73.0)	26665 (72.8)
Not reported	103 (0.6)	110 (0.6)	213 (0.6)
Country			
Argentina	2561 (14.0)	2539 (13.8)	5100 (13.9)
Brazil	1232 (6.8)	1223 (6.7)	2455 (6.7)

Demographic Characteristics – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population			
	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N^a=18242) n^b (%)	Placebo (N^a=18379) n^b (%)	Total (N^a=36621) n^b (%)
Germany	121 (0.7)	126 (0.7)	247 (0.7)
South Africa	287 (1.6)	279 (1.5)	566 (1.5)
USA	14041 (77.0)	14212 (77.3)	28253 (77.1)
Age group			
12-15 Years	46 (0.3)	42 (0.2)	88 (0.2)
16-55 Years	10428 (57.2)	10507 (57.2)	20935 (57.2)
>55 Years	7768 (42.6)	7830 (42.6)	15598 (42.6)
≥65 Years	3980 (21.8)	4038 (22.0)	8018 (21.9)
Age at vaccination (years)			
Mean (SD)	50.6 (15.70)	50.4 (15.81)	50.5 (15.76)
Median	52.0	52.0	52.0
Min, max	(12, 89)	(12, 91)	(12, 91)
Note: HIV-positive subjects are included in this summary but not included in the analyses of the overall study objectives. a. N = number of subjects in the specified group, or the total sample. This value is the denominator for the percentage calculations. b. n = Number of subjects with the specified characteristic. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (10:49) Source Data: adsl Table Generation: 17NOV2020 (18:29) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsl_demo_7d_eval_eff			

**Demographic Characteristics – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 –
Evaluable Efficacy (7 Days) Population**

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N^a=20033) n^b (%)	Placebo (N^a=20244) n^b (%)	Total (N^a=40277) n^b (%)
Sex			
Male	10239 (51.1)	10137 (50.1)	20376 (50.6)
Female	9794 (48.9)	10107 (49.9)	19901 (49.4)
Race			
White	16387 (81.8)	16619 (82.1)	33006 (81.9)
Black or African American	1957 (9.8)	1972 (9.7)	3929 (9.8)
American Indian or Alaska native	131 (0.7)	122 (0.6)	253 (0.6)
Asian	880 (4.4)	883 (4.4)	1763 (4.4)
Native Hawaiian or other Pacific Islander	54 (0.3)	29 (0.1)	83 (0.2)
Multiracial	523 (2.6)	493 (2.4)	1016 (2.5)
Not reported	101 (0.5)	126 (0.6)	227 (0.6)
Ethnicity			
Hispanic/Latino	5272 (26.3)	5281 (26.1)	10553 (26.2)
Non-Hispanic/non-Latino	14652 (73.1)	14847 (73.3)	29499 (73.2)
Not reported	109 (0.5)	116 (0.6)	225 (0.6)
Country			
Argentina	2686 (13.4)	2711 (13.4)	5397 (13.4)
Brazil	1430 (7.1)	1425 (7.0)	2855 (7.1)

**Demographic Characteristics – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 –
Evaluable Efficacy (7 Days) Population**

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =20033) n ^b (%)	Placebo (N ^a =20244) n ^b (%)	Total (N ^a =40277) n ^b (%)
Germany	198 (1.0)	201 (1.0)	399 (1.0)
South Africa	390 (1.9)	393 (1.9)	783 (1.9)
Turkey	31 (0.2)	26 (0.1)	57 (0.1)
USA	15298 (76.4)	15488 (76.5)	30786 (76.4)
Age group			
12-15 Years	48 (0.2)	47 (0.2)	95 (0.2)
16-55 Years	11589 (57.8)	11743 (58.0)	23332 (57.9)
>55 Years	8396 (41.9)	8454 (41.8)	16850 (41.8)
≥65 Years	4294 (21.4)	4319 (21.3)	8613 (21.4)
Age at vaccination (years)			
Mean (SD)	50.3 (15.73)	50.1 (15.78)	50.2 (15.76)
Median	51.0	51.0	51.0
Min, max	(12, 89)	(12, 91)	(12, 91)

Note: HIV-positive subjects are included in this summary but not included in the analyses of the overall study objectives.

a. N = number of subjects in the specified group, or the total sample. This value is the denominator for the percentage calculations.

b. n = Number of subjects with the specified characteristic.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (10:49) Source Data: adsl Table Generation: 17NOV2020 (18:29)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsl_demo_7d_wwo_eval_eff

Demographic Characteristics – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population			
	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N^a=18219) n^b (%)	Placebo (N^a=18315) n^b (%)	Total (N^a=36534) n^b (%)
Sex			
Male	9306 (51.1)	9194 (50.2)	18500 (50.6)
Female	8913 (48.9)	9121 (49.8)	18034 (49.4)
Race			
White	15092 (82.8)	15245 (83.2)	30337 (83.0)
Black or African American	1614 (8.9)	1613 (8.8)	3227 (8.8)
American Indian or Alaska native	118 (0.6)	105 (0.6)	223 (0.6)
Asian	813 (4.5)	809 (4.4)	1622 (4.4)
Native Hawaiian or other Pacific Islander	48 (0.3)	29 (0.2)	77 (0.2)
Multiracial	448 (2.5)	402 (2.2)	850 (2.3)
Not reported	86 (0.5)	112 (0.6)	198 (0.5)
Ethnicity			
Hispanic/Latino	4883 (26.8)	4838 (26.4)	9721 (26.6)
Non-Hispanic/non-Latino	13234 (72.6)	13367 (73.0)	26601 (72.8)
Not reported	102 (0.6)	110 (0.6)	212 (0.6)
Country			
Argentina	2560 (14.1)	2526 (13.8)	5086 (13.9)
Brazil	1231 (6.8)	1219 (6.7)	2450 (6.7)

**Demographic Characteristics – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 –
Evaluable Efficacy (14 Days) Population**

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =18219) n ^b (%)	Placebo (N ^a =18315) n ^b (%)	Total (N ^a =36534) n ^b (%)
Germany	121 (0.7)	125 (0.7)	246 (0.7)
South Africa	287 (1.6)	279 (1.5)	566 (1.5)
USA	14020 (77.0)	14166 (77.3)	28186 (77.2)
Age group			
12-15 Years	46 (0.3)	42 (0.2)	88 (0.2)
16-55 Years	10416 (57.2)	10464 (57.1)	20880 (57.2)
>55 Years	7757 (42.6)	7809 (42.6)	15566 (42.6)
≥65 Years	3971 (21.8)	4027 (22.0)	7998 (21.9)
Age at vaccination (years)			
Mean (SD)	50.6 (15.70)	50.4 (15.81)	50.5 (15.76)
Median	52.0	52.0	52.0
Min, max	(12, 89)	(12, 91)	(12, 91)

Note: HIV-positive subjects are included in this summary but not included in the analyses of the overall study objectives.

a. N = number of subjects in the specified group, or the total sample. This value is the denominator for the percentage calculations.

b. n = Number of subjects with the specified characteristic.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (10:49) Source Data: adsl Table Generation: 17NOV2020 (18:29)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsl_demo_14d_eval_eff

**Demographic Characteristics – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 –
Evaluable Efficacy (14 Days) Population**

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =20033) n ^b (%)	Placebo (N ^a =20243) n ^b (%)	Total (N ^a =40276) n ^b (%)
Sex			
Male	10239 (51.1)	10136 (50.1)	20375 (50.6)
Female	9794 (48.9)	10107 (49.9)	19901 (49.4)
Race			
White	16387 (81.8)	16618 (82.1)	33005 (81.9)
Black or African American	1957 (9.8)	1972 (9.7)	3929 (9.8)
American Indian or Alaska native	131 (0.7)	122 (0.6)	253 (0.6)
Asian	880 (4.4)	883 (4.4)	1763 (4.4)
Native Hawaiian or other Pacific Islander	54 (0.3)	29 (0.1)	83 (0.2)
Multiracial	523 (2.6)	493 (2.4)	1016 (2.5)
Not reported	101 (0.5)	126 (0.6)	227 (0.6)
Ethnicity			
Hispanic/Latino	5272 (26.3)	5281 (26.1)	10553 (26.2)
Non-Hispanic/non-Latino	14652 (73.1)	14846 (73.3)	29498 (73.2)
Not reported	109 (0.5)	116 (0.6)	225 (0.6)
Country			
Argentina	2686 (13.4)	2711 (13.4)	5397 (13.4)
Brazil	1430 (7.1)	1425 (7.0)	2855 (7.1)

**Demographic Characteristics – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 –
Evaluable Efficacy (14 Days) Population**

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =20033) n ^b (%)	Placebo (N ^a =20243) n ^b (%)	Total (N ^a =40276) n ^b (%)
Germany	198 (1.0)	201 (1.0)	399 (1.0)
South Africa	390 (1.9)	393 (1.9)	783 (1.9)
Turkey	31 (0.2)	26 (0.1)	57 (0.1)
USA	15298 (76.4)	15487 (76.5)	30785 (76.4)
Age group			
12-15 Years	48 (0.2)	47 (0.2)	95 (0.2)
16-55 Years	11589 (57.8)	11742 (58.0)	23331 (57.9)
>55 Years	8396 (41.9)	8454 (41.8)	16850 (41.8)
≥65 Years	4294 (21.4)	4319 (21.3)	8613 (21.4)
Age at vaccination (years)			
Mean (SD)	50.3 (15.73)	50.1 (15.78)	50.2 (15.76)
Median	51.0	51.0	51.0
Min, max	(12, 89)	(12, 91)	(12, 91)

Note: HIV-positive subjects are included in this summary but not included in the analyses of the overall study objectives.

a. N = number of subjects in the specified group, or the total sample. This value is the denominator for the percentage calculations.

b. n = Number of subjects with the specified characteristic.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (10:49) Source Data: adsl Table Generation: 17NOV2020 (18:29)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsl_demo_14d_wwo_eval_eff

Demographic Characteristics – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population				
	Vaccine Group (as Randomized)			
	BNT162b2 (30 µg) (N^a=18701) n^b (%)	Placebo (N^a=18627) n^b (%)	Total (N^a=37328) n^b (%)	
Sex				
Male	9557 (51.1)	9359 (50.2)	18916 (50.7)	
Female	9144 (48.9)	9268 (49.8)	18412 (49.3)	
Race				
White	15531 (83.0)	15521 (83.3)	31052 (83.2)	
Black or African American	1645 (8.8)	1627 (8.7)	3272 (8.8)	
American Indian or Alaska native	118 (0.6)	108 (0.6)	226 (0.6)	
Asian	823 (4.4)	820 (4.4)	1643 (4.4)	
Native Hawaiian or other Pacific Islander	49 (0.3)	30 (0.2)	79 (0.2)	
Multiracial	449 (2.4)	407 (2.2)	856 (2.3)	
Not reported	86 (0.5)	114 (0.6)	200 (0.5)	
Ethnicity				
Hispanic/Latino	5005 (26.8)	4963 (26.6)	9968 (26.7)	
Non-Hispanic/non-Latino	13588 (72.7)	13554 (72.8)	27142 (72.7)	
Not reported	108 (0.6)	110 (0.6)	218 (0.6)	
Country				
Argentina	2644 (14.1)	2618 (14.1)	5262 (14.1)	
Brazil	1234 (6.6)	1226 (6.6)	2460 (6.6)	

**Demographic Characteristics – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 –
Dose 2 All-Available Efficacy Population**

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =18701) n ^b (%)	Placebo (N ^a =18627) n ^b (%)	Total (N ^a =37328) n ^b (%)
Germany	127 (0.7)	131 (0.7)	258 (0.7)
South Africa	288 (1.5)	279 (1.5)	567 (1.5)
USA	14408 (77.0)	14373 (77.2)	28781 (77.1)
Age group			
12-15 Years	46 (0.2)	42 (0.2)	88 (0.2)
16-55 Years	10689 (57.2)	10647 (57.2)	21336 (57.2)
>55 Years	7966 (42.6)	7938 (42.6)	15904 (42.6)
≥65 Years	4101 (21.9)	4096 (22.0)	8197 (22.0)
Age at vaccination (years)			
Mean (SD)	50.6 (15.72)	50.4 (15.81)	50.5 (15.76)
Median	52.0	52.0	52.0
Min, max	(12, 89)	(12, 91)	(12, 91)

Note: HIV-positive subjects are included in this summary but not included in the analyses of the overall study objectives.

a. N = number of subjects in the specified group, or the total sample. This value is the denominator for the percentage calculations.

b. n = Number of subjects with the specified characteristic.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (10:49) Source Data: adsl Table Generation: 17NOV2020 (18:29)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsl_demo_7d_d2_aai

**Demographic Characteristics – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 –
Dose 2 All-Available Efficacy Population**

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =20566) n ^b (%)	Placebo (N ^a =20536) n ^b (%)	Total (N ^a =41102) n ^b (%)
Sex			
Male	10527 (51.2)	10293 (50.1)	20820 (50.7)
Female	10039 (48.8)	10243 (49.9)	20282 (49.3)
Race			
White	16874 (82.0)	16880 (82.2)	33754 (82.1)
Black or African American	1987 (9.7)	1985 (9.7)	3972 (9.7)
American Indian or Alaska native	131 (0.6)	124 (0.6)	255 (0.6)
Asian	894 (4.3)	893 (4.3)	1787 (4.3)
Native Hawaiian or other Pacific Islander	55 (0.3)	30 (0.1)	85 (0.2)
Multiracial	524 (2.5)	498 (2.4)	1022 (2.5)
Not reported	101 (0.5)	126 (0.6)	227 (0.6)
Ethnicity			
Hispanic/Latino	5420 (26.4)	5411 (26.3)	10831 (26.4)
Non-Hispanic/non-Latino	15032 (73.1)	15009 (73.1)	30041 (73.1)
Not reported	114 (0.6)	116 (0.6)	230 (0.6)
Country			
Argentina	2793 (13.6)	2806 (13.7)	5599 (13.6)
Brazil	1434 (7.0)	1428 (7.0)	2862 (7.0)

**Demographic Characteristics – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 –
Dose 2 All-Available Efficacy Population**

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =20566) n ^b (%)	Placebo (N ^a =20536) n ^b (%)	Total (N ^a =41102) n ^b (%)
Germany	204 (1.0)	206 (1.0)	410 (1.0)
South Africa	392 (1.9)	394 (1.9)	786 (1.9)
Turkey	33 (0.2)	26 (0.1)	59 (0.1)
USA	15710 (76.4)	15676 (76.3)	31386 (76.4)
Age group			
12-15 Years	48 (0.2)	47 (0.2)	95 (0.2)
16-55 Years	11899 (57.9)	11914 (58.0)	23813 (57.9)
>55 Years	8619 (41.9)	8575 (41.8)	17194 (41.8)
≥65 Years	4425 (21.5)	4383 (21.3)	8808 (21.4)
Age at vaccination (years)			
Mean (SD)	50.3 (15.74)	50.1 (15.77)	50.2 (15.76)
Median	51.0	51.0	51.0
Min, max	(12, 89)	(12, 91)	(12, 91)

Note: HIV-positive subjects are included in this summary but not included in the analyses of the overall study objectives.

a. N = number of subjects in the specified group, or the total sample. This value is the denominator for the percentage calculations.

b. n = Number of subjects with the specified characteristic.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (10:49) Source Data: adsl Table Generation: 17NOV2020 (18:29)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsl_demo_7d_wwd_d2_aai

Demographic Characteristics – Dose 1 All-Available Efficacy Population			
	Vaccine Group (as Randomized)		Total (N^a=43551) n^b (%)
	BNT162b2 (30 µg) (N^a=21768) n^b (%)	Placebo (N^a=21783) n^b (%)	
Sex			
Male	11215 (51.5)	10969 (50.4)	22184 (50.9)
Female	10553 (48.5)	10814 (49.6)	21367 (49.1)
Race			
White	17876 (82.1)	17898 (82.2)	35774 (82.1)
Black or African American	2100 (9.6)	2116 (9.7)	4216 (9.7)
American Indian or Alaska native	160 (0.7)	159 (0.7)	319 (0.7)
Asian	935 (4.3)	933 (4.3)	1868 (4.3)
Native Hawaiian or other Pacific Islander	56 (0.3)	32 (0.1)	88 (0.2)
Multiracial	538 (2.5)	514 (2.4)	1052 (2.4)
Not reported	103 (0.5)	131 (0.6)	234 (0.5)
Ethnicity			
Hispanic/Latino	5680 (26.1)	5673 (26.0)	11353 (26.1)
Non-Hispanic/non-Latino	15967 (73.4)	15990 (73.4)	31957 (73.4)
Not reported	121 (0.6)	120 (0.6)	241 (0.6)
Country			
Argentina	2882 (13.2)	2882 (13.2)	5764 (13.2)
Brazil	1452 (6.7)	1448 (6.6)	2900 (6.7)
Germany	249 (1.1)	250 (1.1)	499 (1.1)

Demographic Characteristics – Dose 1 All-Available Efficacy Population			
	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N^a=21768) n^b (%)	Placebo (N^a=21783) n^b (%)	Total (N^a=43551) n^b (%)
South Africa	401 (1.8)	399 (1.8)	800 (1.8)
Turkey	249 (1.1)	249 (1.1)	498 (1.1)
USA	16535 (76.0)	16555 (76.0)	33090 (76.0)
Age group			
12-15 Years	49 (0.2)	51 (0.2)	100 (0.2)
16-55 Years	12781 (58.7)	12822 (58.9)	25603 (58.8)
>55 Years	8938 (41.1)	8910 (40.9)	17848 (41.0)
≥65 Years	4545 (20.9)	4539 (20.8)	9084 (20.9)
Age at vaccination (years)			
Mean (SD)	50.0 (15.76)	49.8 (15.86)	49.9 (15.81)
Median	51.0	51.0	51.0
Min, max	(12, 89)	(12, 91)	(12, 91)
Note: HIV-positive subjects are included in this summary but not included in the analyses of the overall study objectives. a. N = number of subjects in the specified group, or the total sample. This value is the denominator for the percentage calculations. b. n = Number of subjects with the specified characteristic. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (10:49) Source Data: adsl Table Generation: 17NOV2020 (18:29) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsl_demo_d1_aai			

Efficacy Populations			
	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) n^a (%)	Placebo n^a (%)	Total n^a (%)
Randomized ^b	21823 (100.0)	21828 (100.0)	43651 (100.0)
Dose 1 all-available efficacy population	21768 (99.7)	21783 (99.8)	43551 (99.8)
Subjects without evidence of infection before Dose 1	20314 (93.1)	20296 (93.0)	40610 (93.0)
Subjects excluded from Dose 1 all-available efficacy population	55 (0.3)	45 (0.2)	100 (0.2)
Reason for exclusion ^c			
Did not receive at least 1 vaccination	54 (0.2)	45 (0.2)	99 (0.2)
Did not provide informed consent	1 (0.0)	0	1 (0.0)
Dose 2 all-available efficacy population	20566 (94.2)	20536 (94.1)	41102 (94.2)
Subjects without evidence of infection prior to 7 days after Dose 2	18701 (85.7)	18627 (85.3)	37328 (85.5)
Subjects without evidence of infection prior to 14 days after Dose 2	18678 (85.6)	18563 (85.0)	37241 (85.3)
Subjects excluded from Dose 2 all-available efficacy population	1257 (5.8)	1292 (5.9)	2549 (5.8)
Reason for exclusion ^c			
Did not receive 2 vaccinations	1256 (5.8)	1292 (5.9)	2548 (5.8)
Did not provide informed consent	1 (0.0)	0	1 (0.0)
Evaluable efficacy (7 days) population	20033 (91.8)	20244 (92.7)	40277 (92.3)
Subjects without evidence of infection prior to 7 days after Dose 2	18242 (83.6)	18379 (84.2)	36621 (83.9)
Evaluable efficacy (14 days) population	20033 (91.8)	20243 (92.7)	40276 (92.3)
Subjects without evidence of infection prior to 14 days after Dose 2	18219 (83.5)	18315 (83.9)	36534 (83.7)
Subjects excluded from evaluable efficacy (7 days) population	1790 (8.2)	1584 (7.3)	3374 (7.7)

Efficacy Populations			
	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) n^a (%)	Placebo n^a (%)	Total n^a (%)
Subjects excluded from evaluable efficacy (14 days) population	1790 (8.2)	1585 (7.3)	3375 (7.7)
Reason for exclusion ^c			
Randomized but did not meet all eligibility criteria	36 (0.2)	26 (0.1)	62 (0.1)
Did not provide informed consent	1 (0.0)	0	1 (0.0)
Did not receive all vaccinations as randomized or did not receive Dose 2 within the predefined window (19-42 days after Dose 1)	1550 (7.1)	1561 (7.2)	3111 (7.1)
Had other important protocol deviations on or prior to 7 days after Dose 2	311 (1.4)	60 (0.3)	371 (0.8)
Had other important protocol deviations on or prior to 14 days after Dose 2	311 (1.4)	61 (0.3)	372 (0.9)
Note: HIV-positive subjects are included in this summary but not included in the analyses of the overall study objectives. a. n = Number of subjects with the specified characteristic. b. These values are the denominators for the percentage calculations. c. Subjects may have been excluded for more than 1 reason. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (10:49) Source Data: adsl Table Generation: 17NOV2020 (18:29) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001 Efficacy FA 164/adsl_eff_pop			

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =162)	Total (N ^a =170)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (25.0)	76 (46.9)	78 (45.9)
New or increased cough	3 (37.5)	114 (70.4)	117 (68.8)
New or increased shortness of breath	0 (0.0)	25 (15.4)	25 (14.7)
Chills	2 (25.0)	57 (35.2)	59 (34.7)
New or increased muscle pain	1 (12.5)	81 (50.0)	82 (48.2)
New loss of taste or smell	5 (62.5)	43 (26.5)	48 (28.2)
Sore throat	3 (37.5)	68 (42.0)	71 (41.8)
Diarrhea	1 (12.5)	18 (11.1)	19 (11.2)
Vomiting	2 (25.0)	6 (3.7)	8 (4.7)
Subjects with specific number of signs and symptoms			
1	1 (12.5)	24 (14.8)	25 (14.7)
2	3 (37.5)	46 (28.4)	49 (28.8)
3	4 (50.0)	34 (21.0)	38 (22.4)
4	0 (0.0)	33 (20.4)	33 (19.4)
5	0 (0.0)	16 (9.9)	16 (9.4)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =162)	Total (N ^a =170)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
>5	0 (0.0)	9 (5.6)	9 (5.3)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:47)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_wo_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =9)	Placebo (N ^a =169)	Total (N ^a =178)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (22.2)	77 (45.6)	79 (44.4)
New or increased cough	3 (33.3)	120 (71.0)	123 (69.1)
New or increased shortness of breath	0 (0.0)	26 (15.4)	26 (14.6)
Chills	2 (22.2)	58 (34.3)	60 (33.7)
New or increased muscle pain	1 (11.1)	82 (48.5)	83 (46.6)
New loss of taste or smell	5 (55.6)	46 (27.2)	51 (28.7)
Sore throat	4 (44.4)	71 (42.0)	75 (42.1)
Diarrhea	1 (11.1)	19 (11.2)	20 (11.2)
Vomiting	2 (22.2)	6 (3.6)	8 (4.5)
Subjects with specific number of signs and symptoms			
1	2 (22.2)	24 (14.2)	26 (14.6)
2	3 (33.3)	50 (29.6)	53 (29.8)
3	4 (44.4)	37 (21.9)	41 (23.0)
4	0 (0.0)	33 (19.5)	33 (18.5)
5	0 (0.0)	16 (9.5)	16 (9.0)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =9)	Placebo (N ^a =169)	Total (N ^a =178)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
>5	0 (0.0)	9 (5.3)	9 (5.1)

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:47)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =139)	Total (N ^a =147)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (25.0)	65 (46.8)	67 (45.6)
New or increased cough	3 (37.5)	101 (72.7)	104 (70.7)
New or increased shortness of breath	0 (0.0)	20 (14.4)	20 (13.6)
Chills	2 (25.0)	48 (34.5)	50 (34.0)
New or increased muscle pain	1 (12.5)	67 (48.2)	68 (46.3)
New loss of taste or smell	5 (62.5)	36 (25.9)	41 (27.9)
Sore throat	3 (37.5)	57 (41.0)	60 (40.8)
Diarrhea	1 (12.5)	17 (12.2)	18 (12.2)
Vomiting	2 (25.0)	5 (3.6)	7 (4.8)
Subjects with specific number of signs and symptoms			
1	1 (12.5)	21 (15.1)	22 (15.0)
2	3 (37.5)	39 (28.1)	42 (28.6)
3	4 (50.0)	31 (22.3)	35 (23.8)
4	0 (0.0)	29 (20.9)	29 (19.7)
5	0 (0.0)	10 (7.2)	10 (6.8)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =139)	Total (N ^a =147)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
>5	0 (0.0)	9 (6.5)	9 (6.1)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:47)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_wo_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =144)	Total (N ^a =152)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (25.0)	66 (45.8)	68 (44.7)
New or increased cough	3 (37.5)	105 (72.9)	108 (71.1)
New or increased shortness of breath	0 (0.0)	21 (14.6)	21 (13.8)
Chills	2 (25.0)	48 (33.3)	50 (32.9)
New or increased muscle pain	1 (12.5)	68 (47.2)	69 (45.4)
New loss of taste or smell	5 (62.5)	38 (26.4)	43 (28.3)
Sore throat	3 (37.5)	60 (41.7)	63 (41.4)
Diarrhea	1 (12.5)	17 (11.8)	18 (11.8)
Vomiting	2 (25.0)	5 (3.5)	7 (4.6)
Subjects with specific number of signs and symptoms			
1	1 (12.5)	21 (14.6)	22 (14.5)
2	3 (37.5)	42 (29.2)	45 (29.6)
3	4 (50.0)	33 (22.9)	37 (24.3)
4	0 (0.0)	29 (20.1)	29 (19.1)
5	0 (0.0)	10 (6.9)	10 (6.6)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =144)	Total (N ^a =152)
	n ^b (%)	n ^b (%)	n ^b (%)
>5	0 (0.0)	9 (6.3)	9 (5.9)
a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations. b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:47) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_eval			

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =165)	Total (N ^a =173)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (25.0)	78 (47.3)	80 (46.2)
New or increased cough	3 (37.5)	114 (69.1)	117 (67.6)
New or increased shortness of breath	0 (0.0)	25 (15.2)	25 (14.5)
Chills	2 (25.0)	58 (35.2)	60 (34.7)
New or increased muscle pain	1 (12.5)	82 (49.7)	83 (48.0)
New loss of taste or smell	5 (62.5)	44 (26.7)	49 (28.3)
Sore throat	3 (37.5)	70 (42.4)	73 (42.2)
Diarrhea	1 (12.5)	19 (11.5)	20 (11.6)
Vomiting	2 (25.0)	6 (3.6)	8 (4.6)
Subjects with specific number of signs and symptoms			
1	1 (12.5)	24 (14.5)	25 (14.5)
2	3 (37.5)	47 (28.5)	50 (28.9)
3	4 (50.0)	36 (21.8)	40 (23.1)
4	0 (0.0)	33 (20.0)	33 (19.1)
5	0 (0.0)	16 (9.7)	16 (9.2)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =165)	Total (N ^a =173)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
>5	0 (0.0)	9 (5.5)	9 (5.2)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:47)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_wo_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =9)	Placebo (N ^a =172)	Total (N ^a =181)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (22.2)	79 (45.9)	81 (44.8)
New or increased cough	3 (33.3)	120 (69.8)	123 (68.0)
New or increased shortness of breath	0 (0.0)	26 (15.1)	26 (14.4)
Chills	2 (22.2)	59 (34.3)	61 (33.7)
New or increased muscle pain	1 (11.1)	83 (48.3)	84 (46.4)
New loss of taste or smell	5 (55.6)	47 (27.3)	52 (28.7)
Sore throat	4 (44.4)	73 (42.4)	77 (42.5)
Diarrhea	1 (11.1)	20 (11.6)	21 (11.6)
Vomiting	2 (22.2)	6 (3.5)	8 (4.4)
Subjects with specific number of signs and symptoms			
1	2 (22.2)	24 (14.0)	26 (14.4)
2	3 (33.3)	51 (29.7)	54 (29.8)
3	4 (44.4)	39 (22.7)	43 (23.8)
4	0 (0.0)	33 (19.2)	33 (18.2)
5	0 (0.0)	16 (9.3)	16 (8.8)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Dose 2 All-Available Efficacy Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =9)	Placebo (N ^a =172)	Total (N ^a =181)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
>5	0 (0.0)	9 (5.2)	9 (5.0)

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:47)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =141)	Total (N ^a =149)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (25.0)	67 (47.5)	69 (46.3)
New or increased cough	3 (37.5)	101 (71.6)	104 (69.8)
New or increased shortness of breath	0 (0.0)	20 (14.2)	20 (13.4)
Chills	2 (25.0)	49 (34.8)	51 (34.2)
New or increased muscle pain	1 (12.5)	67 (47.5)	68 (45.6)
New loss of taste or smell	5 (62.5)	36 (25.5)	41 (27.5)
Sore throat	3 (37.5)	58 (41.1)	61 (40.9)
Diarrhea	1 (12.5)	18 (12.8)	19 (12.8)
Vomiting	2 (25.0)	5 (3.5)	7 (4.7)
Subjects with specific number of signs and symptoms			
1	1 (12.5)	21 (14.9)	22 (14.8)
2	3 (37.5)	40 (28.4)	43 (28.9)
3	4 (50.0)	32 (22.7)	36 (24.2)
4	0 (0.0)	29 (20.6)	29 (19.5)
5	0 (0.0)	10 (7.1)	10 (6.7)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =141)	Total (N ^a =149)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
>5	0 (0.0)	9 (6.4)	9 (6.0)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:47)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_wo_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =146)	Total (N ^a =154)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (25.0)	68 (46.6)	70 (45.5)
New or increased cough	3 (37.5)	105 (71.9)	108 (70.1)
New or increased shortness of breath	0 (0.0)	21 (14.4)	21 (13.6)
Chills	2 (25.0)	49 (33.6)	51 (33.1)
New or increased muscle pain	1 (12.5)	68 (46.6)	69 (44.8)
New loss of taste or smell	5 (62.5)	38 (26.0)	43 (27.9)
Sore throat	3 (37.5)	61 (41.8)	64 (41.6)
Diarrhea	1 (12.5)	18 (12.3)	19 (12.3)
Vomiting	2 (25.0)	5 (3.4)	7 (4.5)
Subjects with specific number of signs and symptoms			
1	1 (12.5)	21 (14.4)	22 (14.3)
2	3 (37.5)	43 (29.5)	46 (29.9)
3	4 (50.0)	34 (23.3)	38 (24.7)
4	0 (0.0)	29 (19.9)	29 (18.8)
5	0 (0.0)	10 (6.8)	10 (6.5)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Dose 2 All-Available Efficacy Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =8)	Placebo (N ^a =146)	Total (N ^a =154)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
>5	0 (0.0)	9 (6.2)	9 (5.8)

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:47)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_aai

Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 1 – Dose 1 All-Available Efficacy Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =50)	Placebo (N ^a =275)	Total (N ^a =325)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	20 (40.0)	122 (44.4)	142 (43.7)
New or increased cough	22 (44.0)	186 (67.6)	208 (64.0)
New or increased shortness of breath	4 (8.0)	44 (16.0)	48 (14.8)
Chills	10 (20.0)	86 (31.3)	96 (29.5)
New or increased muscle pain	12 (24.0)	121 (44.0)	133 (40.9)
New loss of taste or smell	24 (48.0)	91 (33.1)	115 (35.4)
Sore throat	18 (36.0)	111 (40.4)	129 (39.7)
Diarrhea	4 (8.0)	35 (12.7)	39 (12.0)
Vomiting	5 (10.0)	11 (4.0)	16 (4.9)
Subjects with specific number of signs and symptoms			
1	16 (32.0)	44 (16.0)	60 (18.5)
2	14 (28.0)	82 (29.8)	96 (29.5)
3	11 (22.0)	63 (22.9)	74 (22.8)
4	5 (10.0)	40 (14.5)	45 (13.8)
5	2 (4.0)	31 (11.3)	33 (10.2)
>5	2 (4.0)	15 (5.5)	17 (5.2)

Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 1 – Dose 1 All-Available Efficacy Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =50)	Placebo (N ^a =275)	Total (N ^a =325)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

a. N = number of subjects with COVID-19 occurrence after dose 1 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:47)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_d1_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =114)	Total (N ^a =119)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	0 (0.0)	52 (45.6)	52 (43.7)
New or increased cough	3 (60.0)	76 (66.7)	79 (66.4)
New or increased shortness of breath	0 (0.0)	19 (16.7)	19 (16.0)
Chills	1 (20.0)	37 (32.5)	38 (31.9)
New or increased muscle pain	0 (0.0)	61 (53.5)	61 (51.3)
New loss of taste or smell	4 (80.0)	36 (31.6)	40 (33.6)
Sore throat	3 (60.0)	53 (46.5)	56 (47.1)
Diarrhea	0 (0.0)	15 (13.2)	15 (12.6)
Vomiting	0 (0.0)	6 (5.3)	6 (5.0)
Subjects with specific number of signs and symptoms			
1	1 (20.0)	13 (11.4)	14 (11.8)
2	2 (40.0)	33 (28.9)	35 (29.4)
3	2 (40.0)	27 (23.7)	29 (24.4)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: 16-55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =114)	Total (N ^a =119)
	n ^b (%)	n ^b (%)	n ^b (%)
4	0 (0.0)	22 (19.3)	22 (18.5)
5	0 (0.0)	11 (9.6)	11 (9.2)
>5	0 (0.0)	8 (7.0)	8 (6.7)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_wo_age_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =48)	Total (N ^a =51)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (66.7)	24 (50.0)	26 (51.0)
New or increased cough	0 (0.0)	38 (79.2)	38 (74.5)
New or increased shortness of breath	0 (0.0)	6 (12.5)	6 (11.8)
Chills	1 (33.3)	20 (41.7)	21 (41.2)
New or increased muscle pain	1 (33.3)	20 (41.7)	21 (41.2)
New loss of taste or smell	1 (33.3)	7 (14.6)	8 (15.7)
Sore throat	0 (0.0)	15 (31.3)	15 (29.4)
Diarrhea	1 (33.3)	3 (6.3)	4 (7.8)
Vomiting	2 (66.7)	0 (0.0)	2 (3.9)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	11 (22.9)	11 (21.6)
2	1 (33.3)	13 (27.1)	14 (27.5)
3	2 (66.7)	7 (14.6)	9 (17.6)
4	0 (0.0)	11 (22.9)	11 (21.6)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =48)	Total (N ^a =51)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
5	0 (0.0)	5 (10.4)	5 (9.8)
>5	0 (0.0)	1 (2.1)	1 (2.0)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_wo_age_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =19)	Total (N ^a =20)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	1 (100.0)	6 (31.6)	7 (35.0)
New or increased cough	0 (0.0)	15 (78.9)	15 (75.0)
New or increased shortness of breath	0 (0.0)	1 (5.3)	1 (5.0)
Chills	1 (100.0)	6 (31.6)	7 (35.0)
New or increased muscle pain	0 (0.0)	4 (21.1)	4 (20.0)
Sore throat	0 (0.0)	6 (31.6)	6 (30.0)
Vomiting	1 (100.0)	0 (0.0)	1 (5.0)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	7 (36.8)	7 (35.0)
2	0 (0.0)	7 (36.8)	7 (35.0)
3	1 (100.0)	3 (15.8)	4 (20.0)
4	0 (0.0)	2 (10.5)	2 (10.0)
Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome			

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =19)	Total (N ^a =20)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
coronavirus 2. Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis. a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations. b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_wo_age_eval			

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =6)	Placebo (N ^a =120)	Total (N ^a =126)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	0 (0.0)	53 (44.2)	53 (42.1)
New or increased cough	3 (50.0)	81 (67.5)	84 (66.7)
New or increased shortness of breath	0 (0.0)	20 (16.7)	20 (15.9)
Chills	1 (16.7)	38 (31.7)	39 (31.0)
New or increased muscle pain	0 (0.0)	62 (51.7)	62 (49.2)
New loss of taste or smell	4 (66.7)	39 (32.5)	43 (34.1)
Sore throat	4 (66.7)	55 (45.8)	59 (46.8)
Diarrhea	0 (0.0)	16 (13.3)	16 (12.7)
Vomiting	0 (0.0)	6 (5.0)	6 (4.8)
Subjects with specific number of signs and symptoms			
1	2 (33.3)	13 (10.8)	15 (11.9)
2	2 (33.3)	36 (30.0)	38 (30.2)
3	2 (33.3)	30 (25.0)	32 (25.4)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =6)	Placebo (N ^a =120)	Total (N ^a =126)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
4	0 (0.0)	22 (18.3)	22 (17.5)
5	0 (0.0)	11 (9.2)	11 (8.7)
>5	0 (0.0)	8 (6.7)	8 (6.3)

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_age_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =49)	Total (N ^a =52)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (66.7)	24 (49.0)	26 (50.0)
New or increased cough	0 (0.0)	39 (79.6)	39 (75.0)
New or increased shortness of breath	0 (0.0)	6 (12.2)	6 (11.5)
Chills	1 (33.3)	20 (40.8)	21 (40.4)
New or increased muscle pain	1 (33.3)	20 (40.8)	21 (40.4)
New loss of taste or smell	1 (33.3)	7 (14.3)	8 (15.4)
Sore throat	0 (0.0)	16 (32.7)	16 (30.8)
Diarrhea	1 (33.3)	3 (6.1)	4 (7.7)
Vomiting	2 (66.7)	0 (0.0)	2 (3.8)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	11 (22.4)	11 (21.2)
2	1 (33.3)	14 (28.6)	15 (28.8)
3	2 (66.7)	7 (14.3)	9 (17.3)
4	0 (0.0)	11 (22.4)	11 (21.2)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =49)	Total (N ^a =52)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
5	0 (0.0)	5 (10.2)	5 (9.6)
>5	0 (0.0)	1 (2.0)	1 (1.9)

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_age_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: ≥65 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =19)	Total (N ^a =20)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	1 (100.0)	6 (31.6)	7 (35.0)
New or increased cough	0 (0.0)	15 (78.9)	15 (75.0)
New or increased shortness of breath	0 (0.0)	1 (5.3)	1 (5.0)
Chills	1 (100.0)	6 (31.6)	7 (35.0)
New or increased muscle pain	0 (0.0)	4 (21.1)	4 (20.0)
Sore throat	0 (0.0)	6 (31.6)	6 (30.0)
Vomiting	1 (100.0)	0 (0.0)	1 (5.0)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	7 (36.8)	7 (35.0)
2	0 (0.0)	7 (36.8)	7 (35.0)
3	1 (100.0)	3 (15.8)	4 (20.0)
4	0 (0.0)	2 (10.5)	2 (10.0)

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Evaluable Efficacy (7 Days) Population

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =19)	Total (N ^a =20)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_age_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =96)	Total (N ^a =101)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	0 (0.0)	45 (46.9)	45 (44.6)
New or increased cough	3 (60.0)	67 (69.8)	70 (69.3)
New or increased shortness of breath	0 (0.0)	14 (14.6)	14 (13.9)
Chills	1 (20.0)	31 (32.3)	32 (31.7)
New or increased muscle pain	0 (0.0)	51 (53.1)	51 (50.5)
New loss of taste or smell	4 (80.0)	29 (30.2)	33 (32.7)
Sore throat	3 (60.0)	44 (45.8)	47 (46.5)
Diarrhea	0 (0.0)	14 (14.6)	14 (13.9)
Vomiting	0 (0.0)	5 (5.2)	5 (5.0)
Subjects with specific number of signs and symptoms			
1	1 (20.0)	11 (11.5)	12 (11.9)
2	2 (40.0)	27 (28.1)	29 (28.7)
3	2 (40.0)	24 (25.0)	26 (25.7)

090177e195901438\Final\Final On: 19-Nov-2020 06:23 (GMT)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: 16-55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =96)	Total (N ^a =101)
	n ^b (%)	n ^b (%)	n ^b (%)
4	0 (0.0)	19 (19.8)	19 (18.8)
5	0 (0.0)	7 (7.3)	7 (6.9)
>5	0 (0.0)	8 (8.3)	8 (7.9)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_wo_age_eva

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =43)	Total (N ^a =46)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (66.7)	20 (46.5)	22 (47.8)
New or increased cough	0 (0.0)	34 (79.1)	34 (73.9)
New or increased shortness of breath	0 (0.0)	6 (14.0)	6 (13.0)
Chills	1 (33.3)	17 (39.5)	18 (39.1)
New or increased muscle pain	1 (33.3)	16 (37.2)	17 (37.0)
New loss of taste or smell	1 (33.3)	7 (16.3)	8 (17.4)
Sore throat	0 (0.0)	13 (30.2)	13 (28.3)
Diarrhea	1 (33.3)	3 (7.0)	4 (8.7)
Vomiting	2 (66.7)	0 (0.0)	2 (4.3)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	10 (23.3)	10 (21.7)
2	1 (33.3)	12 (27.9)	13 (28.3)
3	2 (66.7)	7 (16.3)	9 (19.6)
4	0 (0.0)	10 (23.3)	10 (21.7)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =43)	Total (N ^a =46)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
5	0 (0.0)	3 (7.0)	3 (6.5)
>5	0 (0.0)	1 (2.3)	1 (2.2)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_wo_age_eva

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =17)	Total (N ^a =18)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	1 (100.0)	5 (29.4)	6 (33.3)
New or increased cough	0 (0.0)	13 (76.5)	13 (72.2)
New or increased shortness of breath	0 (0.0)	1 (5.9)	1 (5.6)
Chills	1 (100.0)	5 (29.4)	6 (33.3)
New or increased muscle pain	0 (0.0)	3 (17.6)	3 (16.7)
Sore throat	0 (0.0)	6 (35.3)	6 (33.3)
Vomiting	1 (100.0)	0 (0.0)	1 (5.6)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	6 (35.3)	6 (33.3)
2	0 (0.0)	7 (41.2)	7 (38.9)
3	1 (100.0)	3 (17.6)	4 (22.2)
4	0 (0.0)	1 (5.9)	1 (5.6)
Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome			

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =17)	Total (N ^a =18)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_wo_age_eva

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =100)	Total (N ^a =105)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	0 (0.0)	46 (46.0)	46 (43.8)
New or increased cough	3 (60.0)	70 (70.0)	73 (69.5)
New or increased shortness of breath	0 (0.0)	15 (15.0)	15 (14.3)
Chills	1 (20.0)	31 (31.0)	32 (30.5)
New or increased muscle pain	0 (0.0)	52 (52.0)	52 (49.5)
New loss of taste or smell	4 (80.0)	31 (31.0)	35 (33.3)
Sore throat	3 (60.0)	46 (46.0)	49 (46.7)
Diarrhea	0 (0.0)	14 (14.0)	14 (13.3)
Vomiting	0 (0.0)	5 (5.0)	5 (4.8)
Subjects with specific number of signs and symptoms			
1	1 (20.0)	11 (11.0)	12 (11.4)
2	2 (40.0)	29 (29.0)	31 (29.5)
3	2 (40.0)	26 (26.0)	28 (26.7)

090177e195901438\Final\Final On: 19-Nov-2020 06:23 (GMT)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: 16-55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =100)	Total (N ^a =105)
	n ^b (%)	n ^b (%)	n ^b (%)
4	0 (0.0)	19 (19.0)	19 (18.1)
5	0 (0.0)	7 (7.0)	7 (6.7)
>5	0 (0.0)	8 (8.0)	8 (7.6)

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_age_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =44)	Total (N ^a =47)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (66.7)	20 (45.5)	22 (46.8)
New or increased cough	0 (0.0)	35 (79.5)	35 (74.5)
New or increased shortness of breath	0 (0.0)	6 (13.6)	6 (12.8)
Chills	1 (33.3)	17 (38.6)	18 (38.3)
New or increased muscle pain	1 (33.3)	16 (36.4)	17 (36.2)
New loss of taste or smell	1 (33.3)	7 (15.9)	8 (17.0)
Sore throat	0 (0.0)	14 (31.8)	14 (29.8)
Diarrhea	1 (33.3)	3 (6.8)	4 (8.5)
Vomiting	2 (66.7)	0 (0.0)	2 (4.3)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	10 (22.7)	10 (21.3)
2	1 (33.3)	13 (29.5)	14 (29.8)
3	2 (66.7)	7 (15.9)	9 (19.1)
4	0 (0.0)	10 (22.7)	10 (21.3)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: >55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =44)	Total (N ^a =47)
	n ^b (%)	n ^b (%)	n ^b (%)
5	0 (0.0)	3 (6.8)	3 (6.4)
>5	0 (0.0)	1 (2.3)	1 (2.1)

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_age_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: ≥65 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =17)	Total (N ^a =18)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	1 (100.0)	5 (29.4)	6 (33.3)
New or increased cough	0 (0.0)	13 (76.5)	13 (72.2)
New or increased shortness of breath	0 (0.0)	1 (5.9)	1 (5.6)
Chills	1 (100.0)	5 (29.4)	6 (33.3)
New or increased muscle pain	0 (0.0)	3 (17.6)	3 (16.7)
Sore throat	0 (0.0)	6 (35.3)	6 (33.3)
Vomiting	1 (100.0)	0 (0.0)	1 (5.6)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	6 (35.3)	6 (33.3)
2	0 (0.0)	7 (41.2)	7 (38.9)
3	1 (100.0)	3 (17.6)	4 (22.2)
4	0 (0.0)	1 (5.9)	1 (5.6)
a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the			

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Evaluable Efficacy (14 Days) Population

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =17)	Total (N ^a =18)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_age_eval

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =117)	Total (N ^a =122)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	0 (0.0)	54 (46.2)	54 (44.3)
New or increased cough	3 (60.0)	76 (65.0)	79 (64.8)
New or increased shortness of breath	0 (0.0)	19 (16.2)	19 (15.6)
Chills	1 (20.0)	38 (32.5)	39 (32.0)
New or increased muscle pain	0 (0.0)	62 (53.0)	62 (50.8)
New loss of taste or smell	4 (80.0)	37 (31.6)	41 (33.6)
Sore throat	3 (60.0)	55 (47.0)	58 (47.5)
Diarrhea	0 (0.0)	16 (13.7)	16 (13.1)
Vomiting	0 (0.0)	6 (5.1)	6 (4.9)
Subjects with specific number of signs and symptoms			
1	1 (20.0)	13 (11.1)	14 (11.5)
2	2 (40.0)	34 (29.1)	36 (29.5)
3	2 (40.0)	29 (24.8)	31 (25.4)

090177e195901438\Final\Final On: 19-Nov-2020 06:23 (GMT)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =117)	Total (N ^a =122)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
4	0 (0.0)	22 (18.8)	22 (18.0)
5	0 (0.0)	11 (9.4)	11 (9.0)
>5	0 (0.0)	8 (6.8)	8 (6.6)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_wo_age_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =48)	Total (N ^a =51)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (66.7)	24 (50.0)	26 (51.0)
New or increased cough	0 (0.0)	38 (79.2)	38 (74.5)
New or increased shortness of breath	0 (0.0)	6 (12.5)	6 (11.8)
Chills	1 (33.3)	20 (41.7)	21 (41.2)
New or increased muscle pain	1 (33.3)	20 (41.7)	21 (41.2)
New loss of taste or smell	1 (33.3)	7 (14.6)	8 (15.7)
Sore throat	0 (0.0)	15 (31.3)	15 (29.4)
Diarrhea	1 (33.3)	3 (6.3)	4 (7.8)
Vomiting	2 (66.7)	0 (0.0)	2 (3.9)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	11 (22.9)	11 (21.6)
2	1 (33.3)	13 (27.1)	14 (27.5)
3	2 (66.7)	7 (14.6)	9 (17.6)
4	0 (0.0)	11 (22.9)	11 (21.6)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =48)	Total (N ^a =51)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
5	0 (0.0)	5 (10.4)	5 (9.8)
>5	0 (0.0)	1 (2.1)	1 (2.0)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_wo_age_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =19)	Total (N ^a =20)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	1 (100.0)	6 (31.6)	7 (35.0)
New or increased cough	0 (0.0)	15 (78.9)	15 (75.0)
New or increased shortness of breath	0 (0.0)	1 (5.3)	1 (5.0)
Chills	1 (100.0)	6 (31.6)	7 (35.0)
New or increased muscle pain	0 (0.0)	4 (21.1)	4 (20.0)
Sore throat	0 (0.0)	6 (31.6)	6 (30.0)
Vomiting	1 (100.0)	0 (0.0)	1 (5.0)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	7 (36.8)	7 (35.0)
2	0 (0.0)	7 (36.8)	7 (35.0)
3	1 (100.0)	3 (15.8)	4 (20.0)
4	0 (0.0)	2 (10.5)	2 (10.0)
Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome			

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =19)	Total (N ^a =20)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
coronavirus 2. Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis. a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations. b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_wo_age_aai			

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =6)	Placebo (N ^a =123)	Total (N ^a =129)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	0 (0.0)	55 (44.7)	55 (42.6)
New or increased cough	3 (50.0)	81 (65.9)	84 (65.1)
New or increased shortness of breath	0 (0.0)	20 (16.3)	20 (15.5)
Chills	1 (16.7)	39 (31.7)	40 (31.0)
New or increased muscle pain	0 (0.0)	63 (51.2)	63 (48.8)
New loss of taste or smell	4 (66.7)	40 (32.5)	44 (34.1)
Sore throat	4 (66.7)	57 (46.3)	61 (47.3)
Diarrhea	0 (0.0)	17 (13.8)	17 (13.2)
Vomiting	0 (0.0)	6 (4.9)	6 (4.7)
Subjects with specific number of signs and symptoms			
1	2 (33.3)	13 (10.6)	15 (11.6)
2	2 (33.3)	37 (30.1)	39 (30.2)
3	2 (33.3)	32 (26.0)	34 (26.4)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =6)	Placebo (N ^a =123)	Total (N ^a =129)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
4	0 (0.0)	22 (17.9)	22 (17.1)
5	0 (0.0)	11 (8.9)	11 (8.5)
>5	0 (0.0)	8 (6.5)	8 (6.2)

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_age_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: >55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =49)	Total (N ^a =52)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (66.7)	24 (49.0)	26 (50.0)
New or increased cough	0 (0.0)	39 (79.6)	39 (75.0)
New or increased shortness of breath	0 (0.0)	6 (12.2)	6 (11.5)
Chills	1 (33.3)	20 (40.8)	21 (40.4)
New or increased muscle pain	1 (33.3)	20 (40.8)	21 (40.4)
New loss of taste or smell	1 (33.3)	7 (14.3)	8 (15.4)
Sore throat	0 (0.0)	16 (32.7)	16 (30.8)
Diarrhea	1 (33.3)	3 (6.1)	4 (7.7)
Vomiting	2 (66.7)	0 (0.0)	2 (3.8)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	11 (22.4)	11 (21.2)
2	1 (33.3)	14 (28.6)	15 (28.8)
3	2 (66.7)	7 (14.3)	9 (17.3)
4	0 (0.0)	11 (22.4)	11 (21.2)

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =49)	Total (N ^a =52)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
5	0 (0.0)	5 (10.2)	5 (9.6)
>5	0 (0.0)	1 (2.0)	1 (1.9)

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_age_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: ≥65 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =19)	Total (N ^a =20)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	1 (100.0)	6 (31.6)	7 (35.0)
New or increased cough	0 (0.0)	15 (78.9)	15 (75.0)
New or increased shortness of breath	0 (0.0)	1 (5.3)	1 (5.0)
Chills	1 (100.0)	6 (31.6)	7 (35.0)
New or increased muscle pain	0 (0.0)	4 (21.1)	4 (20.0)
Sore throat	0 (0.0)	6 (31.6)	6 (30.0)
Vomiting	1 (100.0)	0 (0.0)	1 (5.0)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	7 (36.8)	7 (35.0)
2	0 (0.0)	7 (36.8)	7 (35.0)
3	1 (100.0)	3 (15.8)	4 (20.0)
4	0 (0.0)	2 (10.5)	2 (10.0)

a. N = number of subjects with COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage

Summary of Signs and Symptoms for COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: ≥65 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =19)	Total (N ^a =20)
	n ^b (%)	n ^b (%)	n ^b (%)

calculations.
b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.
PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)
(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_7d2_age_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: 16-55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =98)	Total (N ^a =103)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	0 (0.0)	47 (48.0)	47 (45.6)
New or increased cough	3 (60.0)	67 (68.4)	70 (68.0)
New or increased shortness of breath	0 (0.0)	14 (14.3)	14 (13.6)
Chills	1 (20.0)	32 (32.7)	33 (32.0)
New or increased muscle pain	0 (0.0)	51 (52.0)	51 (49.5)
New loss of taste or smell	4 (80.0)	29 (29.6)	33 (32.0)
Sore throat	3 (60.0)	45 (45.9)	48 (46.6)
Diarrhea	0 (0.0)	15 (15.3)	15 (14.6)
Vomiting	0 (0.0)	5 (5.1)	5 (4.9)
Subjects with specific number of signs and symptoms			
1	1 (20.0)	11 (11.2)	12 (11.7)
2	2 (40.0)	28 (28.6)	30 (29.1)
3	2 (40.0)	25 (25.5)	27 (26.2)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: 16-55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =98)	Total (N ^a =103)
	n ^b (%)	n ^b (%)	n ^b (%)
4	0 (0.0)	19 (19.4)	19 (18.4)
5	0 (0.0)	7 (7.1)	7 (6.8)
>5	0 (0.0)	8 (8.2)	8 (7.8)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_wo_age_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: >55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =43)	Total (N ^a =46)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (66.7)	20 (46.5)	22 (47.8)
New or increased cough	0 (0.0)	34 (79.1)	34 (73.9)
New or increased shortness of breath	0 (0.0)	6 (14.0)	6 (13.0)
Chills	1 (33.3)	17 (39.5)	18 (39.1)
New or increased muscle pain	1 (33.3)	16 (37.2)	17 (37.0)
New loss of taste or smell	1 (33.3)	7 (16.3)	8 (17.4)
Sore throat	0 (0.0)	13 (30.2)	13 (28.3)
Diarrhea	1 (33.3)	3 (7.0)	4 (8.7)
Vomiting	2 (66.7)	0 (0.0)	2 (4.3)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	10 (23.3)	10 (21.7)
2	1 (33.3)	12 (27.9)	13 (28.3)
3	2 (66.7)	7 (16.3)	9 (19.6)
4	0 (0.0)	10 (23.3)	10 (21.7)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =43)	Total (N ^a =46)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
5	0 (0.0)	3 (7.0)	3 (6.5)
>5	0 (0.0)	1 (2.3)	1 (2.2)

Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome coronavirus 2.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_wo_age_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: ≥65 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =17)	Total (N ^a =18)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	1 (100.0)	5 (29.4)	6 (33.3)
New or increased cough	0 (0.0)	13 (76.5)	13 (72.2)
New or increased shortness of breath	0 (0.0)	1 (5.9)	1 (5.6)
Chills	1 (100.0)	5 (29.4)	6 (33.3)
New or increased muscle pain	0 (0.0)	3 (17.6)	3 (16.7)
Sore throat	0 (0.0)	6 (35.3)	6 (33.3)
Vomiting	1 (100.0)	0 (0.0)	1 (5.6)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	6 (35.3)	6 (33.3)
2	0 (0.0)	7 (41.2)	7 (38.9)
3	1 (100.0)	3 (17.6)	4 (22.2)
4	0 (0.0)	1 (5.9)	1 (5.6)
Abbreviations: N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; SARS-COV-2 = severe acute respiratory syndrome			

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =17)	Total (N ^a =18)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
coronavirus 2. Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis. a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations. b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_wo_age_aai			

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: 16-55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =102)	Total (N ^a =107)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	0 (0.0)	48 (47.1)	48 (44.9)
New or increased cough	3 (60.0)	70 (68.6)	73 (68.2)
New or increased shortness of breath	0 (0.0)	15 (14.7)	15 (14.0)
Chills	1 (20.0)	32 (31.4)	33 (30.8)
New or increased muscle pain	0 (0.0)	52 (51.0)	52 (48.6)
New loss of taste or smell	4 (80.0)	31 (30.4)	35 (32.7)
Sore throat	3 (60.0)	47 (46.1)	50 (46.7)
Diarrhea	0 (0.0)	15 (14.7)	15 (14.0)
Vomiting	0 (0.0)	5 (4.9)	5 (4.7)
Subjects with specific number of signs and symptoms			
1	1 (20.0)	11 (10.8)	12 (11.2)
2	2 (40.0)	30 (29.4)	32 (29.9)
3	2 (40.0)	27 (26.5)	29 (27.1)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: 16-55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =5)	Placebo (N ^a =102)	Total (N ^a =107)
	n ^b (%)	n ^b (%)	n ^b (%)
4	0 (0.0)	19 (18.6)	19 (17.8)
5	0 (0.0)	7 (6.9)	7 (6.5)
>5	0 (0.0)	8 (7.8)	8 (7.5)

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_age_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: >55 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =44)	Total (N ^a =47)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	2 (66.7)	20 (45.5)	22 (46.8)
New or increased cough	0 (0.0)	35 (79.5)	35 (74.5)
New or increased shortness of breath	0 (0.0)	6 (13.6)	6 (12.8)
Chills	1 (33.3)	17 (38.6)	18 (38.3)
New or increased muscle pain	1 (33.3)	16 (36.4)	17 (36.2)
New loss of taste or smell	1 (33.3)	7 (15.9)	8 (17.0)
Sore throat	0 (0.0)	14 (31.8)	14 (29.8)
Diarrhea	1 (33.3)	3 (6.8)	4 (8.5)
Vomiting	2 (66.7)	0 (0.0)	2 (4.3)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	10 (22.7)	10 (21.3)
2	1 (33.3)	13 (29.5)	14 (29.8)
3	2 (66.7)	7 (15.9)	9 (19.1)
4	0 (0.0)	10 (22.7)	10 (21.3)

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =3)	Placebo (N ^a =44)	Total (N ^a =47)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
5	0 (0.0)	3 (6.8)	3 (6.4)
>5	0 (0.0)	1 (2.3)	1 (2.1)

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_age_aai

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: ≥65 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =17)	Total (N ^a =18)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	1 (100.0)	5 (29.4)	6 (33.3)
New or increased cough	0 (0.0)	13 (76.5)	13 (72.2)
New or increased shortness of breath	0 (0.0)	1 (5.9)	1 (5.6)
Chills	1 (100.0)	5 (29.4)	6 (33.3)
New or increased muscle pain	0 (0.0)	3 (17.6)	3 (16.7)
Sore throat	0 (0.0)	6 (35.3)	6 (33.3)
Vomiting	1 (100.0)	0 (0.0)	1 (5.6)
Subjects with specific number of signs and symptoms			
1	0 (0.0)	6 (35.3)	6 (33.3)
2	0 (0.0)	7 (41.2)	7 (38.9)
3	1 (100.0)	3 (17.6)	4 (22.2)
4	0 (0.0)	1 (5.9)	1 (5.6)

a. N = number of subjects with COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the

Summary of Signs and Symptoms for COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – by Age Group – Dose 2 All-Available Efficacy Population

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =17)	Total (N ^a =18)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_14d2_age_aai

**Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 1 – by Age Group –
Dose 1 All-Available Efficacy Population**

Age Group: 12-15 Years

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =2)	Total (N ^a =3)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of COVID-19			
Fever	1 (100.0)	1 (50.0)	2 (66.7)
New or increased cough	0 (0.0)	2 (100.0)	2 (66.7)
Chills	0 (0.0)	2 (100.0)	2 (66.7)
New or increased muscle pain	0 (0.0)	1 (50.0)	1 (33.3)
New loss of taste or smell	0 (0.0)	2 (100.0)	2 (66.7)
Sore throat	0 (0.0)	2 (100.0)	2 (66.7)
Diarrhea	0 (0.0)	1 (50.0)	1 (33.3)
Subjects with specific number of signs and symptoms			
1	1 (100.0)	0 (0.0)	1 (33.3)
5	0 (0.0)	1 (50.0)	1 (33.3)
>5	0 (0.0)	1 (50.0)	1 (33.3)
a. N = number of subjects with COVID-19 occurrence after dose 1 in the specified group. This value is used as the denominator for the percentage calculations.			

**Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 1 – by Age Group –
Dose 1 All-Available Efficacy Population**

Age Group: 12-15 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =2)	Total (N ^a =3)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.
PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)
(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_d1_age_aai

**Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 1 – by Age Group –
Dose 1 All-Available Efficacy Population**

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =38)	Placebo (N ^a =200)	Total (N ^a =238)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	14 (36.8)	82 (41.0)	96 (40.3)
New or increased cough	16 (42.1)	125 (62.5)	141 (59.2)
New or increased shortness of breath	2 (5.3)	30 (15.0)	32 (13.4)
Chills	9 (23.7)	56 (28.0)	65 (27.3)
New or increased muscle pain	8 (21.1)	90 (45.0)	98 (41.2)
New loss of taste or smell	20 (52.6)	79 (39.5)	99 (41.6)
Sore throat	16 (42.1)	88 (44.0)	104 (43.7)
Diarrhea	3 (7.9)	25 (12.5)	28 (11.8)
Vomiting	3 (7.9)	10 (5.0)	13 (5.5)
Subjects with specific number of signs and symptoms			
1	14 (36.8)	31 (15.5)	45 (18.9)
2	8 (21.1)	58 (29.0)	66 (27.7)
3	8 (21.1)	52 (26.0)	60 (25.2)
4	5 (13.2)	28 (14.0)	33 (13.9)

**Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 1 – by Age Group –
Dose 1 All-Available Efficacy Population**

Age Group: 16-55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =38)	Placebo (N ^a =200)	Total (N ^a =238)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
5	1 (2.6)	20 (10.0)	21 (8.8)
>5	2 (5.3)	11 (5.5)	13 (5.5)

a. N = number of subjects with COVID-19 occurrence after dose 1 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_d1_age_aai

**Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 1 – by Age Group –
Dose 1 All-Available Efficacy Population**

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =11)	Placebo (N ^a =73)	Total (N ^a =84)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	5 (45.5)	39 (53.4)	44 (52.4)
New or increased cough	6 (54.5)	59 (80.8)	65 (77.4)
New or increased shortness of breath	2 (18.2)	14 (19.2)	16 (19.0)
Chills	1 (9.1)	28 (38.4)	29 (34.5)
New or increased muscle pain	4 (36.4)	30 (41.1)	34 (40.5)
New loss of taste or smell	4 (36.4)	10 (13.7)	14 (16.7)
Sore throat	2 (18.2)	21 (28.8)	23 (27.4)
Diarrhea	1 (9.1)	9 (12.3)	10 (11.9)
Vomiting	2 (18.2)	1 (1.4)	3 (3.6)
Subjects with specific number of signs and symptoms			
1	1 (9.1)	13 (17.8)	14 (16.7)
2	6 (54.5)	24 (32.9)	30 (35.7)
3	3 (27.3)	11 (15.1)	14 (16.7)
4	0 (0.0)	12 (16.4)	12 (14.3)

**Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 1 – by Age Group –
Dose 1 All-Available Efficacy Population**

Age Group: >55 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =11)	Placebo (N ^a =73)	Total (N ^a =84)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
5	1 (9.1)	10 (13.7)	11 (13.1)
>5	0 (0.0)	3 (4.1)	3 (3.6)

a. N = number of subjects with COVID-19 occurrence after dose 1 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_d1_age_aai

**Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 1 – by Age Group –
Dose 1 All-Available Efficacy Population**

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =6)	Placebo (N ^a =24)	Total (N ^a =30)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of COVID-19			
Fever	3 (50.0)	10 (41.7)	13 (43.3)
New or increased cough	5 (83.3)	18 (75.0)	23 (76.7)
New or increased shortness of breath	1 (16.7)	3 (12.5)	4 (13.3)
Chills	1 (16.7)	9 (37.5)	10 (33.3)
New or increased muscle pain	1 (16.7)	8 (33.3)	9 (30.0)
New loss of taste or smell	2 (33.3)	1 (4.2)	3 (10.0)
Sore throat	2 (33.3)	6 (25.0)	8 (26.7)
Diarrhea	0 (0.0)	3 (12.5)	3 (10.0)
Vomiting	1 (16.7)	0 (0.0)	1 (3.3)
Subjects with specific number of signs and symptoms			
1	1 (16.7)	7 (29.2)	8 (26.7)
2	2 (33.3)	8 (33.3)	10 (33.3)
3	2 (33.3)	4 (16.7)	6 (20.0)
4	0 (0.0)	2 (8.3)	2 (6.7)

**Summary of Signs and Symptoms for COVID-19 Occurrence After Dose 1 – by Age Group –
Dose 1 All-Available Efficacy Population**

Age Group: ≥65 Years

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =6)	Placebo (N ^a =24)	Total (N ^a =30)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
5	1 (16.7)	3 (12.5)	4 (13.3)

a. N = number of subjects with COVID-19 occurrence after dose 1 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:48)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_cov_d1_age_aai

Summary of Signs and Symptoms for Severe COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =3)	Total (N ^a =4)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of severe COVID-19			
Clinical signs at rest indicative of severe systemic illness (RR ≥30 breaths per minute, HR ≥125 beats per minute, SpO ₂ ≤93% on room air at sea level, or PaO ₂ /FiO ₂ <300 mm Hg)	1 (100.0)	3 (100.0)	4 (100.0)
Respiratory failure (defined as needing high-flow oxygen, noninvasive ventilation, mechanical ventilation, or ECMO)	0 (0.0)	1 (33.3)	1 (25.0)
Admission to an ICU	0 (0.0)	1 (33.3)	1 (25.0)
Subjects with specific number of signs and symptoms			
1	1 (100.0)	2 (66.7)	3 (75.0)
3	0 (0.0)	1 (33.3)	1 (25.0)

Abbreviations: DBP = diastolic blood pressure; ECMO = extracorporeal membrane oxygenation; FiO₂ = fraction of inspired oxygen; HR = heart rate; ICU = intensive care unit; N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; PaO₂ = partial pressure of oxygen, arterial; RR = respiratory rate; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; SBP = systolic blood pressure; SpO₂ = oxygen saturation as measured by pulse oximetry.

Note: Subjects who had no serological or virological evidence (prior to 7 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 7 days after Dose 2 were included in the analysis.

a. N = number of subjects with severe COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

Summary of Signs and Symptoms for Severe COVID-19 Occurrence From 7 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =3)	Total (N ^a =4)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:49)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_sev_cov_7d2_wo_eval

Summary of Signs and Symptoms for Severe COVID-19 Occurrence From 7 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 7 Days After Dose 2 – Evaluable Efficacy (7 Days) Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =3)	Total (N ^a =4)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of severe COVID-19			
Clinical signs at rest indicative of severe systemic illness (RR ≥30 breaths per minute, HR ≥125 beats per minute, SpO ₂ ≤93% on room air at sea level, or PaO ₂ /FiO ₂ <300 mm Hg)	1 (100.0)	3 (100.0)	4 (100.0)
Respiratory failure (defined as needing high-flow oxygen, noninvasive ventilation, mechanical ventilation, or ECMO)	0 (0.0)	1 (33.3)	1 (25.0)
Admission to an ICU	0 (0.0)	1 (33.3)	1 (25.0)
Subjects with specific number of signs and symptoms			
1	1 (100.0)	2 (66.7)	3 (75.0)
3	0 (0.0)	1 (33.3)	1 (25.0)

Abbreviations: DBP = diastolic blood pressure; ECMO = extracorporeal membrane oxygenation; FiO₂ = fraction of inspired oxygen; HR = heart rate; ICU = intensive care unit; PaO₂ = partial pressure of oxygen, arterial; RR = respiratory rate; SBP = systolic blood pressure; SpO₂ = oxygen saturation as measured by pulse oximetry.

a. N = number of subjects with severe COVID-19 occurrence from 7 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:49)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_sev_cov_7d2_eval

Summary of Signs and Symptoms for Severe COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =3)	Total (N ^a =4)
	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of severe COVID-19			
Clinical signs at rest indicative of severe systemic illness (RR ≥30 breaths per minute, HR ≥125 beats per minute, SpO ₂ ≤93% on room air at sea level, or PaO ₂ /FiO ₂ <300 mm Hg)	1 (100.0)	3 (100.0)	4 (100.0)
Respiratory failure (defined as needing high-flow oxygen, noninvasive ventilation, mechanical ventilation, or ECMO)	0 (0.0)	1 (33.3)	1 (25.0)
Admission to an ICU	0 (0.0)	1 (33.3)	1 (25.0)
Subjects with specific number of signs and symptoms			
1	1 (100.0)	2 (66.7)	3 (75.0)
3	0 (0.0)	1 (33.3)	1 (25.0)

Abbreviations: DBP = diastolic blood pressure; ECMO = extracorporeal membrane oxygenation; FiO₂ = fraction of inspired oxygen; HR = heart rate; ICU = intensive care unit; N-binding = SARS-CoV-2 nucleoprotein-binding; NAAT = nucleic acid amplification test; PaO₂ = partial pressure of oxygen, arterial; RR = respiratory rate; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; SBP = systolic blood pressure; SpO₂ = oxygen saturation as measured by pulse oximetry.

Note: Subjects who had no serological or virological evidence (prior to 14 days after receipt of the last dose) of past SARS-CoV-2 infection (ie, N-binding antibody [serum] negative at Visit 1 and SARS-CoV-2 not detected by NAAT [nasal swab] at Visits 1 and 2), and had negative NAAT (nasal swab) at any unscheduled visit prior to 14 days after Dose 2 were included in the analysis.

a. N = number of subjects with severe COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

Summary of Signs and Symptoms for Severe COVID-19 Occurrence From 14 Days After Dose 2 – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =3)	Total (N ^a =4)
	n ^b (%)	n ^b (%)	n ^b (%)
b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom. PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:49) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_sev_cov_14d2_wo_eva			

Summary of Signs and Symptoms for Severe COVID-19 Occurrence From 14 Days After Dose 2 – Subjects With or Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy (14 Days) Population

	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =3)	Total (N ^a =4)
	n ^b (%)	n ^b (%)	n ^b (%)
Signs and Symptoms			
Subjects with specific signs and symptoms of severe COVID-19			
Clinical signs at rest indicative of severe systemic illness (RR ≥30 breaths per minute, HR ≥125 beats per minute, SpO ₂ ≤93% on room air at sea level, or PaO ₂ /FiO ₂ <300 mm Hg)	1 (100.0)	3 (100.0)	4 (100.0)
Respiratory failure (defined as needing high-flow oxygen, noninvasive ventilation, mechanical ventilation, or ECMO)	0 (0.0)	1 (33.3)	1 (25.0)
Admission to an ICU	0 (0.0)	1 (33.3)	1 (25.0)
Subjects with specific number of signs and symptoms			
1	1 (100.0)	2 (66.7)	3 (75.0)
3	0 (0.0)	1 (33.3)	1 (25.0)

Abbreviations: DBP = diastolic blood pressure; ECMO = extracorporeal membrane oxygenation; FiO₂ = fraction of inspired oxygen; HR = heart rate; ICU = intensive care unit; PaO₂ = partial pressure of oxygen, arterial; RR = respiratory rate; SBP = systolic blood pressure; SpO₂ = oxygen saturation as measured by pulse oximetry.

a. N = number of subjects with severe COVID-19 occurrence from 14 days after dose 2 in the specified group. This value is used as the denominator for the percentage calculations.

b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.

PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:49)

(Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_sev_cov_14d2_eval

Summary of Signs and Symptoms for Severe COVID-19 Occurrence After Dose 1 – Dose 1 All-Available Efficacy Population			
	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =9)	Total (N ^a =10)
Signs and Symptoms	n ^b (%)	n ^b (%)	n ^b (%)
Subjects with specific signs and symptoms of severe COVID-19			
Clinical signs at rest indicative of severe systemic illness (RR ≥30 breaths per minute, HR ≥125 beats per minute, SpO ₂ ≤93% on room air at sea level, or PaO ₂ /FiO ₂ <300 mm Hg)	1 (100.0)	9 (100.0)	10 (100.0)
Respiratory failure (defined as needing high-flow oxygen, noninvasive ventilation, mechanical ventilation, or ECMO)	0 (0.0)	3 (33.3)	3 (30.0)
Significant acute renal, hepatic, or neurologic dysfunction	0 (0.0)	1 (11.1)	1 (10.0)
Admission to an ICU	0 (0.0)	3 (33.3)	3 (30.0)
Subjects with specific number of signs and symptoms			
1	1 (100.0)	5 (55.6)	6 (60.0)
2	0 (0.0)	2 (22.2)	2 (20.0)
3	0 (0.0)	1 (11.1)	1 (10.0)
5	0 (0.0)	1 (11.1)	1 (10.0)
Abbreviations: DBP = diastolic blood pressure; ECMO = extracorporeal membrane oxygenation; FiO ₂ = fraction of inspired oxygen; HR = heart rate; ICU = intensive care unit; PaO ₂ = partial pressure of oxygen, arterial; RR = respiratory rate; SBP = systolic blood pressure; SpO ₂ = oxygen saturation as measured by pulse oximetry.			
a. N = number of subjects with severe COVID-19 occurrence after dose 1 in the specified group. This value is used as the denominator for the percentage calculations.			
b. n = Number of subjects with the specific criteria meeting the definition. A subject can have more than 1 symptom.			

Summary of Signs and Symptoms for Severe COVID-19 Occurrence After Dose 1 – Dose 1 All-Available Efficacy Population			
Signs and Symptoms	Vaccine Group (as Randomized)		
	BNT162b2 (30 µg) (N ^a =1)	Placebo (N ^a =9)	Total (N ^a =10)
	n ^b (%)	n ^b (%)	n ^b (%)
PFIZER CONFIDENTIAL SDTM Creation: 17NOV2020 (09:54) Source Data: adc19ef Table Generation: 17NOV2020 (16:49) (Cutoff Date: 14NOV2020, Snapshot Date: 16NOV2020) Output File: ./nda2_unblinded/C4591001_Efficacy_FA_164/adsympt_symp_sev_cov_d1_aai			