ICES NWWG REPORT 2017 | 309

Table 12.2.2. Icelandic Capelin. Abundance of age-classes in numbers (109) measured in acoustic surveys in autumn.

Year	Mon	Day	Age1	Age1	Age2	Age2	Age3	Age3	Age4	Age5
			Imm.	Mat.	Imm.	Mat.	Imm.	Mat	Mat.	Mat.
1978	10	16				60.0		13.9	0.4	
1979	10	14	10.0			49.7		9.1	0.4	
1980	10	11	23.5			19.5		4.8		
1981	11	26	21.0		1.1	11.9		0.6		
1982	10	2	68.0		1.7	15.0		1.6		
1983	10	3	44.1		8.2	58.6		5.6	0.1	
1984	11	1	73.8		4.6	31.9		10.3	0.3	
1985	10	8	33.8		12.6	43.7		14.4	0.4	0.1
1986	10	4	58.6		1.4	19.9		29.8	0.3	
1987	11	18	21.3		2.5	52.0		13.5		
1988	10	6	43.9		6.7	53.0		17.0	0.4	
1989	10	26	29.2		1.8	2.9		0.6		
1990	11	8	24.9		1.3	16.4		2.7	0.1	
1991	11	15	60.0		5.3	44.7		4.2		
1992	10	13	104.6		2.3	54.5		4.3	0.1	
1993	11	18	100.4		9.8	55.1		4.9		
1994	11	25	119.0		6.9	29.2		4.4		
1995	11	30	165.0		30.1	84.6		7.0		
1996	11	27	111.9		16.4	70.0		15.9		
1997	11	1	66.8		30.8	52.5		8.5		
1998	11	13	121.0		5.9	20.5		3.3		
1999	11	15	89.8		4.4	18.1		0.9		
2000	11	10	103.7		10.9	11.6	0.1	0.6		
2001	11	12	101.8		2.4	22.1	0.0	0.7		
2002	11	12	1.0		0.5					
2003	11	6	4.9		3.1	1.7	0.1	0.2		
2004	11	22	7.9		0.1	7.3		0.8	0.0	
2005	11									
2006	11	6	44.7		0.3	5.2		0.4		
2007	11	7	5.7		0.1	1.3		0.0		
2008	11	17	7.5	5.1	0.4	12.1		1.8		
2009	11	24	13.0	2.4		5.0		0.7		
2010	10	1	91.6	9.6	6.3	25.8	0.1	0.8	0.02	
2011	11	29	9.0	0.6	3.6	19.9	0.05	2.1		
2012	10	3	18.5	0.9	2.0	21.2	0.07	11.4	0.1	
2013	9	17	60.1	0.6	6.9	25.0	1.3	6.9	0.1	
2014	9	16	57.0	1.0	3.3	26.5	0.2	7.6	0.1	
2015	9	16	5.0	0.4	1.2	21.2		6.7		
2016	9	10	8.7	0.5	0.7	4.5	0.0	0.9	0.01	

^{1987 -} The number at age 1 was from survey earlier in autumn.

²⁰⁰⁵ - Scouting vessels searched for capelin. r/s $\acute{A}F$ measured. No samples taken for age determination. Estimated to be < 50 thous. tonnes.

²⁰¹¹⁻Only limited coverage of the traditional capelin distribution area.

 $^{2001\}hbox{--}2009 \ and \ 2016-Not \ full \ coverage \ of \ stock.$