

# Bipartite Entanglement Entropy

---

Aditya Chincholi

June 28, 2021

# Quasiperiodic Kicked Rotor

---

## What I did earlier

- Constructing the full floquet operator in a single basis.

## What I did earlier

- Constructing the full floquet operator in a single basis.
- Using a density matrix the whole time for calculations.

## What I did earlier

- Constructing the full floquet operator in a single basis.
- Using a density matrix the whole time for calculations.
- This has the drawback of increasing computational complexity of each individual step and the memory used at any given time is large.

## What I have done now

- Separate the floquet operator into momentum space and position space parts.

## What I have done now

- Separate the floquet operator into momentum space and position space parts.
- Use a single pure state for calculations.

## What I have done now

- Separate the floquet operator into momentum space and position space parts.
- Use a single pure state for calculations.
- Fourier transform the state at each time step and apply both parts of the floquet operator in their resp. basis.



## What I have done now


- Separate the floquet operator into momentum space and position space parts.
- Use a single pure state for calculations.
- Fourier transform the state at each time step and apply both parts of the floquet operator in their resp. basis.
- This is better as the memory used is less but computation increases. Since fourier transforms are computationally cheap anyway, so it's fine.

## What I have done now


- Separate the floquet operator into momentum space and position space parts.
- Use a single pure state for calculations.
- Fourier transform the state at each time step and apply both parts of the floquet operator in their resp. basis.
- This is better as the memory used is less but computation increases. Since fourier transforms are computationally cheap anyway, so it's fine.
- Peak memory required scales the same way but we have reduced it by a constant factor and it is not used in all calculations.

We use  $\hbar = 2.85$ ,  $\omega_2 = 2\pi\sqrt{5}$ ,  $\omega_3 = 2\pi\sqrt{13}$ , the momentum ranges from -10 to 10


$$H = \frac{p_1^2}{2} + p_2\omega_2 + p_3\omega_3 + K\cos(\theta_1)(1 + \alpha\cos(\theta_2)\cos(\theta_3)) \sum_n \delta(t - n)$$



**Figure 1:** Precritical (Insulator):  $K = 4, \alpha = 0.2$



**Figure 2:** Critical:  $K = 6.36, \alpha = 0.4375$



**Figure 3:** Post-critical (Metal):  $K = 8, \alpha = 0.8$

- I don't see much of a trend here. The entanglement grows faster and higher with higher  $K$  values i.e. more diffusive the regime higher the entanglement for the same number of time steps but other than that, I don't see anything here.

## Bypassing The Density Matrix

- What I was doing:  $|\psi\rangle \rightarrow \rho \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$ . This is a memory bottleneck as the full density matrix  $\rho$  still requires the  $O(N^6)$  memory.



## Bypassing The Density Matrix

- What I was doing:  $|\psi\rangle \rightarrow \rho \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$ . This is a memory bottleneck as the full density matrix  $\rho$  still requires the  $O(N^6)$  memory.
- What I realized:  $|\psi\rangle \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$  This gives a really high speedup and allows me to check if localisation is happening properly or not.

## Bypassing The Density Matrix

- What I was doing:  $|\psi\rangle \rightarrow \rho \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$ . This is a memory bottleneck as the full density matrix  $\rho$  still requires the  $O(N^6)$  memory.
- What I realized:  $|\psi\rangle \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$  This gives a really high speedup and allows me to check if localisation is happening properly or not.
- I used these params to generate the following results: p-basis = [-100, 100], 80 timesteps,  $\omega_2 = 2\pi\sqrt{5}$ ,  $\omega_3 = 2\pi\sqrt{13}$ ,  $\hbar = 2.85$ .

## Bypassing The Density Matrix

- What I was doing:  $|\psi\rangle \rightarrow \rho \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$ . This is a memory bottleneck as the full density matrix  $\rho$  still requires the  $O(N^6)$  memory.
- What I realized:  $|\psi\rangle \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$  This gives a really high speedup and allows me to check if localisation is happening properly or not.
- I used these params to generate the following results: p-basis = [-100, 100], 80 timesteps,  $\omega_2 = 2\pi\sqrt{5}$ ,  $\omega_3 = 2\pi\sqrt{13}$ ,  $\hbar = 2.85$ .
- We plot the following quantities:

## Bypassing The Density Matrix

- What I was doing:  $|\psi\rangle \rightarrow \rho \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$ . This is a memory bottleneck as the full density matrix  $\rho$  still requires the  $O(N^6)$  memory.
- What I realized:  $|\psi\rangle \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$  This gives a really high speedup and allows me to check if localisation is happening properly or not.
- I used these params to generate the following results: p-basis = [-100, 100], 80 timesteps,  $\omega_2 = 2\pi\sqrt{5}$ ,  $\omega_3 = 2\pi\sqrt{13}$ ,  $\hbar = 2.85$ .
- We plot the following quantities:
  - $P(p_1 = m\hbar)$

## Bypassing The Density Matrix

- What I was doing:  $|\psi\rangle \rightarrow \rho \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$ . This is a memory bottleneck as the full density matrix  $\rho$  still requires the  $O(N^6)$  memory.
- What I realized:  $|\psi\rangle \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$  This gives a really high speedup and allows me to check if localisation is happening properly or not.
- I used these params to generate the following results: p-basis = [-100, 100], 80 timesteps,  $\omega_2 = 2\pi\sqrt{5}$ ,  $\omega_3 = 2\pi\sqrt{13}$ ,  $\hbar = 2.85$ .
- We plot the following quantities:
  - $P(p_1 = m\hbar)$
  - $E = p_1^2/2 + p_2\omega_2 + p_3\omega_3$

# Bypassing The Density Matrix

- What I was doing:  $|\psi\rangle \rightarrow \rho \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$ . This is a memory bottleneck as the full density matrix  $\rho$  still requires the  $O(N^6)$  memory.
- What I realized:  $|\psi\rangle \rightarrow \rho_1 = \text{Tr}_{2,3}[\rho]$  This gives a really high speedup and allows me to check if localisation is happening properly or not.
- I used these params to generate the following results: p-basis = [-100, 100], 80 timesteps,  $\omega_2 = 2\pi\sqrt{5}$ ,  $\omega_3 = 2\pi\sqrt{13}$ ,  $\hbar = 2.85$ .
- We plot the following quantities:
  - $P(p_1 = m\hbar)$
  - $E = p_1^2/2 + p_2\omega_2 + p_3\omega_3$
  - $S = -\rho_1 \ln(\rho_1)$

# Momentum ( $p_1$ ) distributions





Figure 4:  $K = 3$ ,  $\alpha = 0.1$


# Momentum ( $p_1$ ) distributions




**Figure 5:**  $K = 6.36$ ,  $\alpha = 0.4375$




# Momentum ( $p_1$ ) distributions




**Figure 6:**  $K = 7$ ,  $\alpha = 0.8$



**Figure 7:**  $K = 3$ ,  $\alpha = 0.1$



**Figure 8:**  $K = 6.36$ ,  $\alpha = 0.4375$



**Figure 9:**  $K = 7$ ,  $\alpha = 0.8$

# Entropy






Figure 10:  $K = 3$ ,  $\alpha = 0.1$



**Figure 11:**  $K = 6.36$ ,  $\alpha = 0.4375$



**Figure 12:**  $K = 7$ ,  $\alpha = 0.8$