

Spacing Ratio Distribution

Aditya Chincholi

28 June, 2021

First: A Minor Problem

- Is time reversal symmetry $t \rightarrow -t$?

First: A Minor Problem

- Is time reversal symmetry $t \rightarrow -t$?
- Or is it $t \rightarrow -t, \theta \rightarrow -\theta, p \rightarrow p$?

- Is time reversal symmetry $t \rightarrow -t$?
- Or is it $t \rightarrow -t, \theta \rightarrow -\theta, p \rightarrow p$?
- The issue is the second definition is the one used by Lemarie et al in the “Universality of the Anderson Transition” paper (Lemarié, Grémaud, and Delande 2009).

$$H = \frac{p^2}{2} + K \cos(\theta) \sum_n \delta(t - n\tau)$$

This hamiltonian is symmetric wrt $t \rightarrow -t$, $\theta \rightarrow -\theta$ i.e time reversal and wrt $\theta \rightarrow -\theta$, $p \rightarrow -p$ i.e. parity. The time reversal only holds if we consider only $\Delta t = N\tau$ but that is fine I guess.

- To remind you, I got very odd graphs for the standard kicked rotor system wherein the spacing and ratio distributions were primarily concentrated near zero even after mean normalization (i.e. unfolding).

Parity is a headache

- To remind you, I got very odd graphs for the standard kicked rotor system wherein the spacing and ratio distributions were primarily concentrated near zero even after mean normalization (i.e. unfolding).
- So I tried to study up Mehta's book better and try to understand the issue.

- To remind you, I got very odd graphs for the standard kicked rotor system wherein the spacing and ratio distributions were primarily concentrated near zero even after mean normalization (i.e. unfolding).
- So I tried to study up Mehta's book better and try to understand the issue.
- Since the Hamiltonian has parity symmetry, I think that is what is causing the problem. The eigenvectors have a parity quantum number which needs to be separated.

Parity is a headache

- To remind you, I got very odd graphs for the standard kicked rotor system wherein the spacing and ratio distributions were primarily concentrated near zero even after mean normalization (i.e. unfolding).
- So I tried to study up Mehta's book better and try to understand the issue.
- Since the Hamiltonian has parity symmetry, I think that is what is causing the problem. The eigenvectors have a parity quantum number which needs to be separated.
- ~~But even then, the distribution should be e^{-s} for the spacing, which is not the case.~~

Parity is a headache

- To remind you, I got very odd graphs for the standard kicked rotor system wherein the spacing and ratio distributions were primarily concentrated near zero even after mean normalization (i.e. unfolding).
- So I tried to study up Mehta's book better and try to understand the issue.
- Since the Hamiltonian has parity symmetry, I think that is what is causing the problem. The eigenvectors have a parity quantum number which needs to be separated.
- ~~But even then, the distribution should be e^{-s} for the spacing, which is not the case.~~
- The Izrailev and Atas papers I was talking about last time were indeed on kicked rotors on a torus as you suspected.

Parity is a headache (cont.)

- So I tried separating out the eigenvectors by determining their parity, but the eigenvectors I obtained were not of sufficient accuracy to distinguish the parity. They are good for a 100×100 matrix, but for a 1000×1000 matrix, the eigenvectors are not accurate enough for me to do this.

Parity is a headache (cont.)

- So I tried separating out the eigenvectors by determining their parity, but the eigenvectors I obtained were not of sufficient accuracy to distinguish the parity. They are good for a 100×100 matrix, but for a 1000×1000 matrix, the eigenvectors are not accurate enough for me to do this.
- I thought that maybe the eigenvectors I was getting were linear combos so I tried to understand how to simultaneously diagonalize two matrices.

Parity is a headache (cont.)

- So I tried separating out the eigenvectors by determining their parity, but the eigenvectors I obtained were not of sufficient accuracy to distinguish the parity. They are good for a 100×100 matrix, but for a 1000×1000 matrix, the eigenvectors are not accurate enough for me to do this.
- I thought that maybe the eigenvectors I was getting were linear combos so I tried to understand how to simultaneously diagonalize two matrices.
- Apparently, we just take an arbitrary linear combination of the two matrices and solve for the eigenvectors of this linear combination to get the required eigenvectors. I tried it, but it wasn't very successful.

Parity is a headache (cont.)

- So I tried separating out the eigenvectors by determining their parity, but the eigenvectors I obtained were not of sufficient accuracy to distinguish the parity. They are good for a 100×100 matrix, but for a 1000×1000 matrix, the eigenvectors are not accurate enough for me to do this.
- I thought that maybe the eigenvectors I was getting were linear combos so I tried to understand how to simultaneously diagonalize two matrices.
- Apparently, we just take an arbitrary linear combination of the two matrices and solve for the eigenvectors of this linear combination to get the required eigenvectors. I tried it, but it wasn't very successful.
- Maybe I'm missing something but apparently, eigenvector solving doesn't allow me to specify accuracy/tolerance anywhere.

Parity is a headache (cont.)

- So I tried separating out the eigenvectors by determining their parity, but the eigenvectors I obtained were not of sufficient accuracy to distinguish the parity. They are good for a 100×100 matrix, but for a 1000×1000 matrix, the eigenvectors are not accurate enough for me to do this.
- I thought that maybe the eigenvectors I was getting were linear combos so I tried to understand how to simultaneously diagonalize two matrices.
- Apparently, we just take an arbitrary linear combination of the two matrices and solve for the eigenvectors of this linear combination to get the required eigenvectors. I tried it, but it wasn't very successful.
- Maybe I'm missing something but apparently, eigenvector solving doesn't allow me to specify accuracy/tolerance anywhere.
- ~~All in all, I have no clue what is going wrong.~~

Finding the root

- In order to find out the main issue with my code, I began to try and generate random matrices from different classes as test cases.

- In order to find out the main issue with my code, I began to try and generate random matrices from different classes as test cases.
- As it turns out, people write a great deal of theoretical analysis but rarely write about how to generate them numerically. The symplectic ensembles are particularly hard to find in this case.

Finding the root

- In order to find out the main issue with my code, I began to try and generate random matrices from different classes as test cases.
- As it turns out, people write a great deal of theoretical analysis but rarely write about how to generate them numerically. The symplectic ensembles are particularly hard to find in this case.
- I found a few places where it was detailed and I managed to implement GUE, GOE, GSE, CUE, COE, CSE generators. (Mezzadri 2007; Edelman and Rao 2005; Sirca and Horvat 2012)

Finding the root

- In order to find out the main issue with my code, I began to try and generate random matrices from different classes as test cases.
- As it turns out, people write a great deal of theoretical analysis but rarely write about how to generate them numerically. The symplectic ensembles are particularly hard to find in this case.
- I found a few places where it was detailed and I managed to implement GUE, GOE, GSE, CUE, COE, CSE generators. (Mezzadri 2007; Edelman and Rao 2005; Sirca and Horvat 2012)
- It turned out that my program worked correctly for the unitary and orthogonal ensembles but failed for the symplectic ensembles because they show degeneracy.

- The symplectic ensembles show Kramer's degeneracy due to the half integral spin. Because of that, each energy level has at least a double degeneracy (up and down spin for spin $\frac{1}{2}$).

Finding the root (cont.)

- The symplectic ensembles show Kramer's degeneracy due to the half integral spin. Because of that, each energy level has at least a double degeneracy (up and down spin for spin $\frac{1}{2}$).
- So by writing code that kept only unique eigenvalues (or eigenphases), the issue got resolved.

Finding the root (cont.)

- The symplectic ensembles show Kramer's degeneracy due to the half integral spin. Because of that, each energy level has at least a double degeneracy (up and down spin for spin $\frac{1}{2}$).
- So by writing code that kept only unique eigenvalues (or eigenphases), the issue got resolved.
- And this worked for the earlier case of the quantum kicked rotor as well. Using the parity eigenvalue would be more accurate in some sense because we don't remove other symmetries that may exist that we don't know of, but it is not good numerically.

Finding the root (cont.)

- The symplectic ensembles show Kramer's degeneracy due to the half integral spin. Because of that, each energy level has at least a double degeneracy (up and down spin for spin $\frac{1}{2}$).
- So by writing code that kept only unique eigenvalues (or eigenphases), the issue got resolved.
- And this worked for the earlier case of the quantum kicked rotor as well. Using the parity eigenvalue would be more accurate in some sense because we don't remove other symmetries that may exist that we don't know of, but it is not good numerically.
- This method removes the degeneracies arising from all the symmetries, but if we are sure that there is only one symmetry this will work. By taking the ratio of number of unique eigenvalues to the original number of eigenvalues, we can find the mean degeneracy.




Figure 1: Standard kicked rotor with $K = 5$, p -basis $[-2000, 2000]$, 3991 eigenphases used and 10 discarded (tol = 0.1). Uniqueness ratio is 0.52 with tol = 10^{-8}

Quasiperiodic Kicked Rotor Systems

$$H = \frac{p_1^2}{2} + p_2\omega_2 + p_3\omega_3 + K\cos(\theta_1)(1 + \alpha\cos(\theta_2)\cos(\theta_3)) \sum_n \delta(t - n)$$

We use $\hbar = 2.85$, $\omega_2 = 2\pi\sqrt{5}$, $\omega_3 = 2\pi\sqrt{13}$ and the momentum basis is from -10 to 10. We consider a tolerance of 0.1 for discarding eigenvalues based on their modulus. We consider a tolerance of 10^{-8} for degeneracy.

Results (cont.)

For large K values, the system nicely fits into the poisson distribution.




Figure 2: Post-critical (Metallic): $K = 7$, $\alpha = 0.8$ with 6934 eigenphases used. No degeneracy, but 2327 phases were discarded as they were out of the $[0.9, 1.1]$ range

Results (cont.)

At critical point, things seem okay, but there is a noticeable difference at zero.




Figure 3: Critical: $K = 6.36$, $\alpha = 0.4375$ with 7874 eigenphases. No degeneracy, but 1387 phases were discarded for being out of range.

Results (cont.)

For small K values, things get weird...




Figure 4: Pre-critical (Insulator): $K = 4$, $\alpha = 0.1$ with 8379 eigenphases. No degeneracy, but 882 phases were discarded for being out of range.

- The program from Delande is a program that simulates the 1d version of the quasiperiodic kicked rotor. That doesn't help me much.

- The program from Delande is a program that simulates the 1d version of the quasiperiodic kicked rotor. That doesn't help me much.
- I tried plotting the energy, entropy, p_1 -distribution plots for the matrix. But the dimensionality is too low for me to actually see anything. It does look pretty similar to the corresponding plots made by the direct method, but there are slight differences.

- The program from Delande is a program that simulates the 1d version of the quasiperiodic kicked rotor. That doesn't help me much.
- I tried plotting the energy, entropy, p_1 -distribution plots for the matrix. But the dimensionality is too low for me to actually see anything. It does look pretty similar to the corresponding plots made by the direct method, but there are slight differences.
- Since I am trusting the direct method, I thought of generating the matrix using that and comparing the two. Now what I'm essentially doing in this is calculating $F\mathbf{e}_j$ and comparing it with the j th column of the matrix generated.

- The two are not as close as one would hope it to be but the error seems to decrease with the size of the matrix. The maximum size I could do on my laptop was $[-15, 15]$.
Unsurprisingly, the difference is maximum near the edges both row-wise and col-wise.

- The two are not as close as one would hope it to be but the error seems to decrease with the size of the matrix. The maximum size I could do on my laptop was $[-15, 15]$.
Unsurprisingly, the difference is maximum near the edges both row-wise and col-wise.
- I also tried generating random states and comparing the difference between the generated outcomes, but I thought the direct matrix comparison was a better metric.

- The two are not as close as one would hope it to be but the error seems to decrease with the size of the matrix. The maximum size I could do on my laptop was $[-15, 15]$.
Unsurprisingly, the difference is maximum near the edges both row-wise and col-wise.
- I also tried generating random states and comparing the difference between the generated outcomes, but I thought the direct matrix comparison was a better metric.
- Matching tolerances: $\text{atol} = 10^{-5}$, $\text{rtol} = 10^{-8}$




Figure 5: $K = 3$, $\alpha = 0.1$




Figure 6: $K = 3$, $\alpha = 0.1$




Figure 7: $K = 3$, $\alpha = 0.1$




Figure 8: $K = 3$, $\alpha = 0.1$

- Edelman, Alan, and N. Raj Rao. 2005. "Random Matrix Theory." *Acta Numerica* 14 (May): 233–97.
<https://doi.org/10.1017/S0962492904000236>.
- Lemarié, G., B. Grémaud, and D. Delande. 2009. "Universality of the Anderson Transition with the Quasiperiodic Kicked Rotor." *Europhys. Lett.* 87 (3): 37007.
<https://doi.org/10.1209/0295-5075/87/37007>.
- Mezzadri, Francesco. 2007. "How to Generate Random Matrices from the Classical Compact Groups." February 27, 2007. <http://arxiv.org/abs/math-ph/0609050>.
- Sirca, Simon, and Martin Horvat. 2012. *Computational Methods for Physicists: Compendium for Students*. Graduate Texts in Physics. Berlin Heidelberg: Springer-Verlag.
<https://www.springer.com/in/book/9783642324772>.