

Report on Bipartite Entanglement Entropy

Aditya Chincholi

June 28, 2021

Introduction

$$H = \frac{p_1^2}{2} + p_2\omega_2 + p_3\omega_3 + K\cos(\theta_1)(1 + \alpha\cos(\theta_2)\cos(\theta_3)) \sum_n \delta(t - n)$$

- We are calculating the entanglement entropy between \mathcal{H}_1 and \mathcal{H}_{23} .

$$H = \frac{p_1^2}{2} + p_2\omega_2 + p_3\omega_3 + K\cos(\theta_1)(1 + \alpha\cos(\theta_2)\cos(\theta_3)) \sum_n \delta(t - n)$$

- We are calculating the entanglement entropy between \mathcal{H}_1 and \mathcal{H}_{23} .
- The hypothesis was that the entanglement should increase rapidly at the critical point.

$$H = \frac{p_1^2}{2} + p_2\omega_2 + p_3\omega_3 + K\cos(\theta_1)(1 + \alpha\cos(\theta_2)\cos(\theta_3)) \sum_n \delta(t - n)$$

- We are calculating the entanglement entropy between \mathcal{H}_1 and \mathcal{H}_{23} .
- The hypothesis was that the entanglement should increase rapidly at the critical point.
- Calculated and plotted quantities are:

$$H = \frac{p_1^2}{2} + p_2\omega_2 + p_3\omega_3 + K\cos(\theta_1)(1 + \alpha\cos(\theta_2)\cos(\theta_3)) \sum_n \delta(t - n)$$

- We are calculating the entanglement entropy between \mathcal{H}_1 and \mathcal{H}_{23} .
- The hypothesis was that the entanglement should increase rapidly at the critical point.
- Calculated and plotted quantities are:
 1. Entanglement entropy: $S = -Tr(\rho_1 \ln(\rho_1))$

$$H = \frac{p_1^2}{2} + p_2\omega_2 + p_3\omega_3 + K\cos(\theta_1)(1 + \alpha\cos(\theta_2)\cos(\theta_3)) \sum_n \delta(t - n)$$

- We are calculating the entanglement entropy between \mathcal{H}_1 and \mathcal{H}_{23} .
- The hypothesis was that the entanglement should increase rapidly at the critical point.
- Calculated and plotted quantities are:
 1. Entanglement entropy: $S = -\text{Tr}(\rho_1 \ln(\rho_1))$
 2. Energy: $E = p_1^2/2 + p_2\omega_2 + p_3\omega_3$

$$H = \frac{p_1^2}{2} + p_2\omega_2 + p_3\omega_3 + K\cos(\theta_1)(1 + \alpha\cos(\theta_2)\cos(\theta_3)) \sum_n \delta(t - n)$$

- We are calculating the entanglement entropy between \mathcal{H}_1 and \mathcal{H}_{23} .
- The hypothesis was that the entanglement should increase rapidly at the critical point.
- Calculated and plotted quantities are:
 1. Entanglement entropy: $S = -Tr(\rho_1 \ln(\rho_1))$
 2. Energy: $E = p_1^2/2 + p_2\omega_2 + p_3\omega_3$
 3. Momentum: $P(p_1 = m\hbar)$

$$H = \frac{p_1^2}{2} + p_2\omega_2 + p_3\omega_3 + K\cos(\theta_1)(1 + \alpha\cos(\theta_2)\cos(\theta_3)) \sum_n \delta(t - n)$$

- We are calculating the entanglement entropy between \mathcal{H}_1 and \mathcal{H}_{23} .
- The hypothesis was that the entanglement should increase rapidly at the critical point.
- Calculated and plotted quantities are:
 1. Entanglement entropy: $S = -Tr(\rho_1 \ln(\rho_1))$
 2. Energy: $E = p_1^2/2 + p_2\omega_2 + p_3\omega_3$
 3. Momentum: $P(p_1 = m\hbar)$
 4. Entropy Change: $S_{K_{n+1}, \alpha_{n+1}} - S_{K_n, \alpha_n}$

$$H = \frac{p_1^2}{2} + p_2\omega_2 + p_3\omega_3 + K\cos(\theta_1)(1 + \alpha\cos(\theta_2)\cos(\theta_3)) \sum_n \delta(t - n)$$

- We are calculating the entanglement entropy between \mathcal{H}_1 and \mathcal{H}_{23} .
- The hypothesis was that the entanglement should increase rapidly at the critical point.
- Calculated and plotted quantities are:
 1. Entanglement entropy: $S = -Tr(\rho_1 \ln(\rho_1))$
 2. Energy: $E = p_1^2/2 + p_2\omega_2 + p_3\omega_3$
 3. Momentum: $P(p_1 = m\hbar)$
 4. Entropy Change: $S_{K_{n+1}, \alpha_{n+1}} - S_{K_n, \alpha_n}$
 5. Energy Change: $E_{K_{n+1}, \alpha_{n+1}} - E_{K_n, \alpha_n}$

- We use $\hbar = 1$, $\omega_2 = 2\pi\sqrt{5}$ and $\omega_3 = 2\pi\sqrt{13}$. The critical point of the metal insulator transition is known to occur at $K_c = 6.36 \pm 0.02$.

- We use $\hbar = 1$, $\omega_2 = 2\pi\sqrt{5}$ and $\omega_3 = 2\pi\sqrt{13}$. The critical point of the metal insulator transition is known to occur at $K_c = 6.36 \pm 0.02$.
- The value of α_c , however is not determined accurately, nor written in any literature as it doesn't form a part of the scaling function in a nice way, though it most certainly does matter as it generates the anisotropy. We determine the value of α_c roughly by using the fact that the authors of the cited papers traversed the $K - \alpha$ space along a straight line perpendicular to the transition parabola.^{1 2}


- We use $\hbar = 1$, $\omega_2 = 2\pi\sqrt{5}$ and $\omega_3 = 2\pi\sqrt{13}$. The critical point of the metal insulator transition is known to occur at $K_c = 6.36 \pm 0.02$.
- The value of α_c , however is not determined accurately, nor written in any literature as it doesn't form a part of the scaling function in a nice way, though it most certainly does matter as it generates the anisotropy. We determine the value of α_c roughly by using the fact that the authors of the cited papers traversed the $K - \alpha$ space along a straight line perpendicular to the transition parabola.^{1 2}
- TL;DR we used $K_c = 6.36$ and $\alpha_c = 0.4303$.

Parameters


- We use $\hbar = 1$, $\omega_2 = 2\pi\sqrt{5}$ and $\omega_3 = 2\pi\sqrt{13}$. The critical point of the metal insulator transition is known to occur at $K_c = 6.36 \pm 0.02$.
- The value of α_c , however is not determined accurately, nor written in any literature as it doesn't form a part of the scaling function in a nice way, though it most certainly does matter as it generates the anisotropy. We determine the value of α_c roughly by using the fact that the authors of the cited papers traversed the $K - \alpha$ space along a straight line perpendicular to the transition parabola.^{1 2}
- TL;DR we used $K_c = 6.36$ and $\alpha_c = 0.4303$.
- We used a basis size of 201 (-100 to 100) for each of the 3 coordinates and the simulations were done for 80 timesteps.


Varying K


Bipartite Entanglement Entropy




Energy Evolution








Energy Difference




Varying α


Bipartite Entanglement Entropy



Energy Evolution




Momentum Distribution




Entropy Difference

$$S_{K_{n+1}}(t) - S_{K_n}(t)$$

10^{-2}




Energy Difference




Varying K and α


Bipartite Entanglement Entropy




Energy Evolution




Momentum Distribution





Energy Difference



- We use α from 0.1 to 0.8 and $\alpha_c = 0.5$.

1.

Lemarié, G., Grémaud, B. & Delande, D. Universality of the Anderson transition with the quasiperiodic kicked rotor. *Europhys. Lett.* **87**, 37007 (2009).

2.

Lemarié, G. *et al.* Observation of the Anderson metal-insulator transition with atomic matter waves: Theory and experiment. *Phys. Rev. A* **80**, 043626 (2009).