

Spacing Ratio Distribution

Aditya Chincholi

28 June, 2021

First: A Minor Problem

- Is time reversal symmetry $t \rightarrow -t$?

First: A Minor Problem

- Is time reversal symmetry $t \rightarrow -t$?
- Or is it $t \rightarrow -t, \theta \rightarrow -\theta, p \rightarrow p$?

The issue is the second definition is the one used by Lemarie et al in the “Universality of the Anderson Transition” paper.

The Problem

$$H = \frac{p^2}{2} + K \cos(\theta) \sum_n \delta(t - n\tau)$$

This hamiltonian is symmetric wrt $t \rightarrow -t$, $\theta \rightarrow -\theta$ i.e time reversal and wrt $\theta \rightarrow -\theta$, $p \rightarrow -p$ i.e. parity. The time reversal only holds if we consider only $\Delta t = N\tau$ but that is fine I guess.

Parity is a headache

- To remind you, I got very odd graphs for the standard kicked rotor system wherein the spacing and ratio distributions were primarily concentrated near zero even after mean normalization (i.e. unfolding).

Parity is a headache

- To remind you, I got very odd graphs for the standard kicked rotor system wherein the spacing and ratio distributions were primarily concentrated near zero even after mean normalization (i.e. unfolding).
- So I tried to study up Mehta's book better and try to understand the issue.

Parity is a headache

- To remind you, I got very odd graphs for the standard kicked rotor system wherein the spacing and ratio distributions were primarily concentrated near zero even after mean normalization (i.e. unfolding).
- So I tried to study up Mehta's book better and try to understand the issue.
- Since the Hamiltonian has parity symmetry, I think that is what is causing the problem. The eigenvectors have a parity quantum number which needs to be separated.

Parity is a headache

- To remind you, I got very odd graphs for the standard kicked rotor system wherein the spacing and ratio distributions were primarily concentrated near zero even after mean normalization (i.e. unfolding).
- So I tried to study up Mehta's book better and try to understand the issue.
- Since the Hamiltonian has parity symmetry, I think that is what is causing the problem. The eigenvectors have a parity quantum number which needs to be separated.
- But even then, the distribution should be e^{-s} for the spacing. which is not the case.

Parity is a headache

- To remind you, I got very odd graphs for the standard kicked rotor system wherein the spacing and ratio distributions were primarily concentrated near zero even after mean normalization (i.e. unfolding).
- So I tried to study up Mehta's book better and try to understand the issue.
- Since the Hamiltonian has parity symmetry, I think that is what is causing the problem. The eigenvectors have a parity quantum number which needs to be separated.
- But even then, the distribution should be e^{-s} for the spacing. which is not the case.
- The Izrailev and Atas papers I was talking about last time were indeed on kicked rotors on a torus as you suspected.

- So I tried separating out the eigenvectors by determining their parity, but the eigenvectors I obtained were not of sufficient accuracy to distinguish the parity. They are good for a 100×100 matrix, but for a 1000×1000 matrix, the eigenvectors are not accurate enough for me to do this.

- So I tried separating out the eigenvectors by determining their parity, but the eigenvectors I obtained were not of sufficient accuracy to distinguish the parity. They are good for a 100×100 matrix, but for a 1000×1000 matrix, the eigenvectors are not accurate enough for me to do this.
- I thought that maybe the eigenvectors I was getting were linear combos so I tried to understand how to simultaneously diagonalize two matrices.

- So I tried separating out the eigenvectors by determining their parity, but the eigenvectors I obtained were not of sufficient accuracy to distinguish the parity. They are good for a 100×100 matrix, but for a 1000×1000 matrix, the eigenvectors are not accurate enough for me to do this.
- I thought that maybe the eigenvectors I was getting were linear combos so I tried to understand how to simultaneously diagonalize two matrices.
- Apparently, we just take an arbitrary linear combination of the two matrices and solve for the eigenvectors of this linear combination to get the required eigenvectors. I tried it, but it wasn't very successful.

- So I tried separating out the eigenvectors by determining their parity, but the eigenvectors I obtained were not of sufficient accuracy to distinguish the parity. They are good for a 100×100 matrix, but for a 1000×1000 matrix, the eigenvectors are not accurate enough for me to do this.
- I thought that maybe the eigenvectors I was getting were linear combos so I tried to understand how to simultaneously diagonalize two matrices.
- Apparently, we just take an arbitrary linear combination of the two matrices and solve for the eigenvectors of this linear combination to get the required eigenvectors. I tried it, but it wasn't very successful.
- Maybe I'm missing something but apparently, eigenvector solving doesn't allow me to specify accuracy/tolerance anywhere.

- So I tried separating out the eigenvectors by determining their parity, but the eigenvectors I obtained were not of sufficient accuracy to distinguish the parity. They are good for a 100×100 matrix, but for a 1000×1000 matrix, the eigenvectors are not accurate enough for me to do this.
- I thought that maybe the eigenvectors I was getting were linear combos so I tried to understand how to simultaneously diagonalize two matrices.
- Apparently, we just take an arbitrary linear combination of the two matrices and solve for the eigenvectors of this linear combination to get the required eigenvectors. I tried it, but it wasn't very successful.
- Maybe I'm missing something but apparently, eigenvector solving doesn't allow me to specify accuracy/tolerance anywhere.
- All in all, I have no clue what is going wrong.