

# 1. Pseudosolutions and its applications. Linear regression.

Let's repeat main possible situation for solving linear equations task, which one can be written by the next notation:


$$A\vec{x} = \vec{b},$$

where  $A \in M_{m \times n}(\mathbb{C})$ ,  $\vec{b} \in \mathbb{C}^m$ ,  $\vec{x} \in \mathbb{C}^n$ .

0. The first case is about square matrix  $A \in M_{n \times n}(\mathbb{C})$ ,  $\text{rank } A = n$ . In such situation we can easily obtain unique  $\vec{x}$  by inverting the matrix of initial coefficients:

$$\vec{x} = A^{-1}\vec{b}.$$

1. The next easy option is a definite system, when  $A \in M_{m \times n}(\mathbb{C})$ ,  $\text{rank } A = n$ . Then unique  $\hat{x}$  can be expressed by the following ideas.



Example of definite system.

Consider a system of the form:

$$\begin{cases} 2x + y = 3, \\ x + 2y = 3, \\ x - y = 0. \end{cases}$$

It is obviously that system have only one correct solution in the point A and it is a solution of a type:  $\hat{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . However, we would like to generalize the method of obtaining a solution in such a way that it looks similar to the first (zero) case, namely:

$$\hat{x} = ? \cdot \vec{b}.$$

And looking ahead we can obtain such a factor to express solution that way. But now let's get a broader generalization.

2. Also we can obtain an indefinite solution, that can provide us an infinite amount of solutions.


Consider a system of two equations:

$$\begin{cases} x + 2y = 3, \\ 2x + 4y = 5. \end{cases}$$

It is not so obvious to choose a specific solution here because a whole family of solutions of the following form  $\hat{x} = \begin{bmatrix} 3 - 2y \\ y \end{bmatrix}$  is suitable for us.

And now we need to get some understanding about which solution is a kind of optimum. We will discuss it a little bit later, now let's consider one more possible situation.

3.



Example of definite system.

Inconsistent system is a system of a kind:

$$\begin{cases} 2x + y = 3, \\ 2x + y = 6 \end{cases}$$


Need to remind, that we want to obtain solution in term of factors:

$$\hat{x} = ? \cdot \vec{b}.$$

There we have matrix and vector of initial values

$$A = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \text{ and } \vec{b} = \begin{bmatrix} 3 \\ 6 \end{bmatrix}.$$

Manually we can understand that the best solution will lie somewhere between two parallel lines, perhaps even exactly in the middle. But it is still a whole family of solutions that can be the answer to the request of the product or business problem. We need a general variant to find the best solution. For this we introduce the definition:



Example of inconsistent system.

### Definition

Consider a system of a linear equations  $A\vec{x} = \vec{b}$  ( $A \in M_{m \times n}(\mathbb{C})$ ). A vector  $\vec{u} \in \mathbb{C}^n$  is called a pseudosolution or a least square solution, if  $\forall \vec{x} \in \mathbb{C}^n$  the length of  $A\vec{u} - \vec{b}$  is less or equal to the length of  $A\vec{x} - \vec{b}$ :

$$|A\vec{u} - \vec{b}| \leq |A\vec{x} - \vec{b}|.$$

That is: if  $f_x = \begin{bmatrix} f_1 \\ \vdots \\ f_n \end{bmatrix} = A\vec{x} - \vec{b}$ , then  $|f_x|^2 = |f_1|^2 + \dots + |f_n|^2$  for  $\vec{x} = \vec{u}$  is minimal.

### Theorem

The vector  $\vec{u} = A^+ \vec{b}$  is a pseudosolution of the system of linear equations  $A\vec{x} = \vec{b}$ . Moreover, among all pseudosolutions,  $\vec{u}$  has the minimal length.

**Prop** If  $\hat{x}$  is a solution, then it is a pseudosolution.

**Proof:**  $A\hat{x} - \vec{b} = 0 \implies |A\hat{x} - \vec{b}| = 0 = \min |f_x|^2.$

□

Example 1:

Type of a system	Solution
definite	$\vec{u} = \hat{x}$ is the solution
indefinite	$\vec{u} = \hat{x}$ is the solution of minimal length
inconsistent	$\vec{u} = \hat{x}$ is the pseudosolution of minimal length


**Proof:** (Of the theorem) In proof we will use

**Theorem: (Pythagoras)**

Suppose  $\vec{a} \perp \vec{b}$ , that is  $(\vec{a}, \vec{b}) = 0$ . Then for  $\vec{c} = \vec{a} + \vec{b}$ :  $|\vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2$ . In particular  $|\vec{c}| \geq |\vec{a}|$ . The equality holds only for  $\vec{b} = \vec{0}$ .

**Lemma**

$\text{Im}A \perp \text{Im}B$ , where  $B = AA^+ - I$ .



We should prove: each column  $A^j$  of  $A$  is orthogonal to the one  $B^l$  of  $B$ .

$$\text{OR: } \forall l \ (B^l, A^j) \stackrel{?}{=} 0, \text{ or } B^{l*} \cdot A^j \stackrel{?}{=} 0,$$

$$\text{or } (B^*)^l \cdot A^j \stackrel{?}{=} 0, \text{ or } B^* A \stackrel{?}{=} 0.$$

$$((AA^+)^* - I^*) A = (AA^+ - I) A = AA^+ A - A = 0.$$

□