

Exploring Flood Risk Asheville, NC

https://github.com/addienavarro/Diamond_Navarro_Von_Turkovich_ENV872_EDA_FinalProject

Sarah Diamond, Addie Navaro, Natalie von Turkovich

Contents

0.1	Rationale and Research Questions	5
0.2	Dataset Information	6
0.2.1	Discharge Data	6
0.2.2	Precipitation Data	6
0.2.3	Combined Data	6
0.3	Exploratory Analysis	7
0.3.1	Discharge Exploration	7
0.3.2	Precipitation Exploration	9
0.4	Analysis	10
0.4.1	Discharge Analysis	10
0.4.2	Precipitation Analysis	14
0.4.3	Discharge and Precipitation Relationship	16
0.5	Summary and Conclusions	18
0.6	References:	19

List of Tables

1 French Broad River USGS Discharge Data.	7
2 Asheville NOAA Precipitation Data.	7
3 Discharge and Precipitation Combined Data.	7
4 Data types of combined discharge and precipitation data.	8
5 Significant rainfall events in Asheville, NC.	14

List of Figures

1	Flooding in Asheville.	5
2	French Broad Discharge over time.	8
3	Asheville precipitation over time.	9
4	French Broad River discharge, time series decompostion.	10
5	French Broad River Discharge, trend over time with trend line.	11
6	French Broad River Discharge, trend over time with seasonal variation line.	12
7	French Broad River Discharge, trend over time with line of best fit.	13
8	Frequency and magnitude of precipitation events over time.	15
9	French Broad River discharge and Asheville precipitation, linear model results.	16
10	French Broad River discharge by Asheville precipitation, linear relationship.	17
11	Flooding in Asheville.	19

0.1 Rationale and Research Questions

The earth's climate is changing and is resulting in higher seas, new weather patterns and stronger storms (Floodfactor, 2022). The warming atmosphere is causing more evaporation, which leads to more water available for precipitation (Floodfactor, 2022). This is resulting in more extreme weather events. Both the frequency and magnitude of weather events is increasing worldwide. North Carolina has not been spared these new, more extreme weather patterns. Intense rains in North Carolina have been causing flooding. Devastating flooding as occurred recently in the mountainous, western portion of the state. Flooding in Asheville, located in the western North Carolina along the French Broad River, resulted in two fatalities in the fall of 2022 (Harris, 2021). The orographic rains that this mountainous region is prone to, along with the topography of the terrain, contributes to this area's vulnerability to flooding.

In light of the recent flooding in Asheville, we are interested in exploring if flood risk in the mountain city is increasing over time. To do this, we will analyze both precipitation data in Asheville as well as river discharge data on the French Broad River. The data sets we will look at include NOAA precipitation data and USGS stream gage data. We will be asking the following questions:

1. Is discharge increasing over time?
2. What trends exist in the discharge data over time?
3. Is precipitation increasing over time?
4. Are the frequency of significant precipitation events increasing over time?
5. Is the magnitude of significant rainfall events increasing over time?
6. Does precipitation have a significant effect on rain fall?



Figure 1: Flooding in Asheville.

0.2 Dataset Information

0.2.1 Discharge Data

To understand the discharge in Asheville, North Carolina, we used data from the United States Geographical Survey's (USGS) National Water Information System (NWIS). We were able to pull this dataset into R using the data retrieval function which lets you simply put the specific USGS code for the area we were interested in looking more closely at, ours being the French Broad River. We chose stream gauge station 03451500 which is close to the city center of Asheville. There were multiple parameters available for this site including discharge, precipitation, pH, stream level, etc. By identifying the USGS code as well as the specific codes for the parameters we wanted to look at (i.e., discharge data) we were able to pull in corresponding data for the last 60 years. Because discharge data is recorded daily, we had records of every day from 1963 to 2021. The pulled dataset included the agency (USGS), the site number, the date, and the amount of discharge in cubic feet per second (Table 1).

Because we were able to pull in exactly which columns we wanted, there was not much to wrangle for this specific dataset. We did however decide that for all the parameters we were looking at that we would only include the last forty years (1981-2021). In order to do this, we filtered the dataset to only include those specific years. We also used the lubridate package to change the date column to have a date class format.

0.2.2 Precipitation Data

For precipitation data, we felt it was important to pull daily precipitation data to better understand how extreme precipitation could affect flash flooding of the French Broad River (Table 2). The dataset we pulled is a csv file from the NOAA Global Historical Climatology Network. When downloading the data, we were able to specify dates from 1981-2021 in order to obtain 40 years' worth of data so we could compare earlier and later 20 year segments. This also helped us to match the daily precipitation data to the daily discharge data we obtained from the USGS National Water Information System.

In order to better understand what significant precipitation looked like in Asheville, North Carolina, we read in another csv file from the NOAA Precipitation Frequency Estimate Database. This table gave us information about the 24-hour 1-year rainfall event, 2-year rainfall event, and 5-year rainfall event (Table 5). With this knowledge, we were able to explore whether extreme precipitation events (daily precipitation over the 1-year, 2-year, and 5-year thresholds) had increased over time and whether this impacted river discharge and flood risk in the city.

0.2.3 Combined Data

For our linear model dataset, we combined both the discharge and the precipitation datasets into one using the leftjoin function (Table 3). Doing so, we had both the date, the amount of discharge in cubic feet per second, and the precipitation in milliliters from the past 40 years. For some reason, some of the precipitation entries were negative numbers so we filtered to make sure the final dataset included only precipitation values greater than 0 milliliters.

French Broad River USGS Discharge Data			
Agency Code	Site Number	Date	Discharge
USGS	03451500	1981-01-02	867
USGS	03451500	1981-01-03	840
USGS	03451500	1981-01-04	825
USGS	03451500	1981-01-05	751
USGS	03451500	1981-01-06	790
USGS	03451500	1981-01-07	800

Table 1:
French Broad River USGS Discharge Data Table.

Asheville NOAA Precipitation Data			
Date	Precip.mm	Month	Year
1981-01-01	0.3	1	1981
1981-01-02	0.0	1	1981
1981-01-03	0.0	1	1981
1981-01-04	0.0	1	1981
1981-01-05	0.0	1	1981
1981-01-06	0.0	1	1981

Table 2:
Asheville NOAA Precipitation Data Table.

0.3 Exploratory Analysis

0.3.1 Discharge Exploration

In order to better understand the flood risk in Asheville, North Carolina, we were interested in understanding the daily discharge data throughout time (Figure 2) as well as the overall relationship between discharge and precipitation. To measure this we wanted to run a time series of the discharge data (Figure 4) as well as a seasonal decomposition to understand both the seasonality of the data and the trend throughout time. We also wanted to understand what in fact was the trend over time and how this information can inform the flood risk for Asheville.

As for the relationship between discharge and precipitation, we felt the best way to see this was through a general linear model (Figure 9). By looking at the dependence that discharge may have over precipitation, we were confident that this too would help us to answer our original research question of how these two parameters and the relationship between them have impacted the flood risk over time.

Discharge and Precipitation Combined Data				
Date	Precip.mm	Month	Year	Discharge
1981-01-01	0.3	1	1981	NA
1981-01-07	0.3	1	1981	800
1981-01-15	0.5	1	1981	800
1981-01-20	5.6	1	1981	770
1981-01-21	0.3	1	1981	770
1981-01-27	2.3	1	1981	703

Table 3:
Discharge and Precipitation Combined Data.

Data Set Structure	
Data	Data Types
Date	Date
Percip mm	Numeric
Month	Numeric
Year	Numeric
Discharge	Numeric

Table 4:

Data types of combined discharge and precipitation data.




Figure 2: French Broad Discharge over time.

0.3.2 Precipitation Exploration

One of the first things we did with precipitation data was plot it over time (Figure 3) to see if we could visually see an increase in precipitation over time. This plot did not give us much valuable information, so we decided to create filtered datasets to better see the intense precipitation events above the 1-year, 2-year, and 5-year thresholds (Table 5).




Figure 3: Asheville precipitation over time.

0.4 Analysis

Through our analysis we became more familiar with the data and the relationship between precipitation and discharge in Asheville.

The following analysis can be divided into three parts:

1. How has discharge changed over time?
2. How has precipitation changed over time?
3. What is the relationship of discharge to precipitation?

0.4.1 Discharge Analysis

To first look at the trends of discharge data over time, we ran a simple timeseries that began on January 1, 1981, and ran through December 31, 2021, with a frequency of 365 days (Figure 4). Because this data had a seasonal component, we examined the decomposition of this timeseries data to get a better visual of the trend and seasonality throughout time (Figure 6).




Figure 4: French Broad River discharge, time series decomposition.

After this, we created an “Observed” column in the dataframe to show the discharge with the corresponding date so we could visualize both the trend and the seasonality throughout time (Figure 5).

French Broad Discharge with Trend




Figure 5: French Broad River Discharge, trend over time with trend line.

French Broad Discharge with Seasonal Variation




Figure 6: French Broad River Discharge, trend over time with seasonal variation line.

Lastly, we used the Seasonal Mann-Kendall test to know whether the trend over time was positive or negative. Our results of this test showed that we could reject the null hypothesis that there is no trend in the seasonal data and that there is a positive trend over time in the discharge data ($p\text{value} > 0.05$, $\tau = 0.0622$). The final plot of the trend can be seen below.




Figure 7: French Broad River Discharge, trend over time with line of best fit.

Significant Rainfall in Asheville, NC										
Duration	1 year	2 year	5 year	10 year	25 year	50 year	100 year	200 year	500 year	1000 year
5-min:	8	10	12	14	16	17	19	20	22	24
10-min:	14	16	19	22	25	28	30	32	35	38
15-min:	17	20	25	28	32	35	38	41	44	47
30-min:	23	28	35	40	47	52	58	63	71	77
60-min:	29	35	45	52	63	71	80	89	102	112
2-hr:	33	40	51	59	71	81	91	102	117	129
3-hr:	35	41	52	61	74	85	96	108	125	139
6-hr:	41	48	60	70	84	96	109	123	143	159
12-hr:	50	59	73	84	99	111	124	136	153	166
24-hr:	55	66	82	95	112	125	139	153	171	184
2-day:	66	79	97	111	130	145	160	175	194	209
3-day:	70	84	102	116	136	151	165	180	199	213
4-day:	74	89	107	122	141	156	171	185	204	217
7-day:	87	104	125	141	163	180	196	213	234	249
10-day:	100	118	141	158	181	199	216	234	255	271
20-day:	138	162	189	210	237	258	278	297	321	338
30-day:	172	201	230	252	279	299	318	335	357	371
45-day:	218	255	286	310	338	358	376	393	412	424
60-day:	262	305	341	366	398	420	439	457	477	490

Table 5:

Significant rainfall events in Asheville, NC.

0.4.2 Precipitation Analysis

Filtering the data for the 1-year 24-hour precipitation events revealed that more precipitation events over the 1-year 55mm threshold occurred in the decades following 2000 than the decades preceding 2000.

This trend became more obvious as we increased the threshold of 24-hour rainfall events to the 2-year 66mm level. The scatterplot and frequency histogram revealed that the more intense rainfall events above the 2-year recurrence interval threshold were increasing in frequency.

Finally, we looked at the 5-year recurrence interval 24-hour rainfall event and saw that of the 7 extreme precipitation events over the 5-year 82mm threshold, 6 occurred after 2000 and only one occurred before 2000. Notably, the most extreme 24-hour precipitation event of the last 40 years occurred only last year in the fall of 2021.

Examining the precipitation data at the 1-year, 2-year, and 5-year 24-hour recurrence interval thresholds helped us to see that while there wasn't a significant increase in overall precipitation over the last 40 years, more extreme rainfall events are occurring more frequently in the last 20 years than in the previous two decades before 2000 (Figure 7).




Figure 8: Frequency and magnitude of precipitation events over time.

0.4.3 Discharge and Precipitation Relationship

In our next analysis, we looked more at the relationship between discharge and the precipitation data. As mentioned previously, we used a general linear model to look more closely at this relationship. Our results of this linear model regression show that there was in fact a significant relationship between the two variables, and we were confidently able to reject the null hypothesis that there was no relationship between discharge and precipitation. It can be seen below in the QQ-Plot that this data is not really normally distributed and that it would be advantageous to potentially log-transform the data to view the relationship better (Figure 8).




Figure 9: French Broad River discharge and Asheville precipitation, linear model results.

Looking closer at the linear model results, we were able to determine that this was a positive relationship and that for every one unit of precipitation, discharge increased by 48.3 cubic feet per second ($p\text{-value} < 0.05$, $F\text{-statistic} = 324.3$ on 1 and 525 degrees of freedom, $R^2 = 0.05805$). Our R^2 value however only explained around 6% of the variability and this could be due to the fact that discharge and precipitation isn't usually modelled linearly which could account for the lower R^2 and could be something to investigate further. A detailed visualization of the relationship can be seen in the plot below (Figure 9).

Both of these analyses can be helpful to inform our original research question of how these variables and the relationship between them have changed over time and will be discussed at greater length in our summary and conclusion sections.




Figure 10: French Broad River discharge by Asheville precipitation, linear relationship.

0.5 Summary and Conclusions

Through our analysis we observed how precipitation and river discharge are changing in Asheville over time. Even though there may be an increase of flooding events that, does not always correlate to an increases in precipitation and discharge as large events could be balanced by period of lower than normal rainfall. Our analysis did allow us to answer our research questions and gain a better understand of hydrologic events in Asheville in the past 40 years.

Through our analysis of discharge in the French Broad river we were able to answer our first research question of what trends exist in discharge data over time. We found that although small, there is indeed an increase in overall discharge moving through the French Broad river in the past 40 years. After decomposing our time series analysis we found that the trend of the overall data was not clearly increasing. We did however find that there was a slight visual correspondence between the seasonal data and the overall shape of the time series plot. This finding was confirmed with the results of the Seasonal Man Kendall test. Our results of this test showed that we could reject the null hypothesis that there is no trend in the seasonal data and that there is a positive trend over time in the discharge data, answering our second research question.

Analysis of the precipitation data in Asheville resulted in a result of a positive increase in the frequency of extreme precipitation events over time, answering research questions number three (Figure 3). Additionally, we looked at the frequency of the 24-hour rainfall event at the one-year, two-year, and five-year recurrence interval. We found that over the past 40 years, more intense precipitation events over the 1-year, 2-year, and 5-year thresholds have been increased in frequency. At the most extreme end of our analysis, the five-year rainfall event, we found that daily rainfall has exceeded this threshold almost exclusively in the past 20 years. This analysis allowed us to answer our fourth and fifth research questions. We can conclude that intense rainfall events are increasing over time in Asheville (Figure 7).

An analysis of the relationship between precipitation and river discharge gave us a significant result answering our sixth research question. As one would suspect, precipitation has a positive effect on river discharge. This result provides further evidence that the increases measured in precipitation event magnitude and frequency will also have an increase in magnitude and frequency of high discharge in the river (Figure 9).

The changing climate puts to the test the systems that have been built around data from the past. Acknowledging these increases in hydrologic activity, which lead to an increase in flood risk, requires municipalities to assess their current storm water system and flood plain boundaries. Awareness of the changing conditions is necessary for communities to be able to adapt and build resilience to climate risks in their areas.



Figure 11: Flooding in Asheville.

0.6 References:

Harris, S. (2021, August 20). Asheville and Buncombe flooding related road, park closures, Help Line. The Asheville Citizen Times. Retrieved April 16, 2022, from <https://www.citizen-times.com/story/news/2021/08/20/asheville-buncombe-nc-area-flooding-road-park-closures-help-line/8211528002/>

FloodFactor (2022). Asheville, North Carolina. Flood Factor. https://floodfactor.com/city/asheville-northcarolina/3702140_fsid

Figure 1: Photo credit: <https://www.ashevillenc.gov/department/public-works/stormwater-services-utility/flood-information/>

Figure 2: Photo credit: Amy Westmoreland