

# An Introduction into Building Theory

(Using Abstract Simplicial Complexes and Coxeter Groups)

Malachi Alexander

UC Santa Cruz

May 20, 2021

# Table of Contents

- 1 Motivation
- 2 Basics
  - Finite Reflection Groups
  - Coxeter Complex
- 3 Examples

# Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field.

# Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

# Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).

# Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.

# Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.
- (3) The apartments in the first perspective are viewed as metric spaces rather than simplicial complexes with specific geometric properties.

# Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.
- (3) The apartments in the first perspective are viewed as metric spaces rather than simplicial complexes with specific geometric properties.

The perspectives differ in how one views the chambers:



# Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.
- (3) The apartments in the first perspective are viewed as metric spaces rather than simplicial complexes with specific geometric properties.

The perspectives differ in how one views the chambers:

- (1) Chambers are maximal simplices.

# Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.
- (3) The apartments in the first perspective are viewed as metric spaces rather than simplicial complexes with specific geometric properties.

The perspectives differ in how one views the chambers:

- (1) Chambers are maximal simplices.
- (2) Chambers are elements of an abstract set.

# Motivation and Perspectives

Buildings were introduced by Jacques Tits in order to provide a unified geometric framework for understanding semisimple complex Lie groups and semisimple algebraic groups over an arbitrary field. There are three perspectives on buildings:

- (1) Simplicial complex with a family of subcomplexes (called apartments) composed of top-dimensional simplices (called chambers).
- (2) A collection of chambers together with a Weyl-group valued distance function.
- (3) The apartments in the first perspective are viewed as metric spaces rather than simplicial complexes with specific geometric properties.

The perspectives differ in how one views the chambers:

- (1) Chambers are maximal simplices.
- (2) Chambers are elements of an abstract set.
- (3) Chambers are metric spaces.

# Motivation and Perspectives

In each perspective, the notion of Coxeter groups are required.

# Motivation and Perspectives

In each perspective, the notion of Coxeter groups are required.

- We will use finite reflection groups (which characterize finite Coxeter groups).

# Motivation and Perspectives

In each perspective, the notion of Coxeter groups are required.

- We will use finite reflection groups (which characterize finite Coxeter groups).
- We will present from the simplicial complex perspective.

# Motivation and Perspectives

In each perspective, the notion of Coxeter groups are required.

- We will use finite reflection groups (which characterize finite Coxeter groups).
- We will present from the simplicial complex perspective.

## Definition of a Building

A (*weak*) *building* is a simplicial complex  $\Delta$  that can be expressed as the union of subcomplexes  $\Sigma$  (called *apartments*) satisfying axioms:

- (**B0**) Each apartment  $\Sigma$  is a Coxeter complex.
- (**B1**) For any two simplices  $A, B \in \Delta$ , there is an apartment  $\Sigma$  containing both of them.
- (**B2**) If  $\Sigma$  and  $\Sigma'$  are two apartments containing  $A$  and  $B$ , then there is an isomorphism  $\Sigma \rightarrow \Sigma'$  fixing  $A$  and  $B$  pointwise.

# Finite Reflection Groups


Let  $V$  be a Euclidean vector space, i.e. a finite-dimensional  $\mathbb{R}$ -vector space with an inner product and let  $V$  have dimension  $n$ .



# Finite Reflection Groups

Let  $V$  be a Euclidean vector space, i.e. a finite-dimensional  $\mathbb{R}$ -vector space with an inner product and let  $V$  have dimension  $n$ .

**Example:** Let  $V = \mathbb{R}^3$  equipped with the dot product.



## Inner Product

$$\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_1 + u_2 v_2 + u_3 v_3$$


# Finite Reflection Groups

**Definition:** A hyperplane  $H$  in  $V$  is a subspace of codimension 1 (we are assuming  $\dim_{\mathbb{R}}(V) = n$ , thus  $\dim_{\mathbb{R}}(H) = n - 1$ ).

# Finite Reflection Groups

**Definition:** A hyperplane  $H$  in  $V$  is a subspace of codimension 1 (we are assuming  $\dim_{\mathbb{R}}(V) = n$ , thus  $\dim_{\mathbb{R}}(H) = n - 1$ ).

**Example:** Let  $V = \mathbb{R}^3$  and  $H_1 = \{\mathbf{u} \in \mathbb{R}^3 : \mathbf{u} = (u_1, 0, u_3)\}$ .





# Finite Reflection Groups

**Definition:** A reflection with respect to  $H$  is a linear transformation  $s_H : V \rightarrow V$  such that  $s_H$  is the identity on  $H$  and multiplication by  $-1$  along the orthogonal complement.

# Finite Reflection Groups

**Definition:** A reflection with respect to  $H$  is a linear transformation  $s_H : V \rightarrow V$  such that  $s_H$  is the identity on  $H$  and multiplication by  $-1$  along the orthogonal complement.

**Example:** Let  $u_1 = (1, 0, 1)$  and  $u_2 = (1, 1, 1)$ .



We obtain  $s_{H_1}(1, 0, 1) = (1, 0, 1)$  and  $s_{H_1}(1, 1, 1) = (1, -1, 1)$ .

# Finite Reflection Groups




**Definition:** Let  $\mathcal{H}$  be a set of hyperplanes in  $V$ , we call this a hyperplane arrangement. A (finite) reflection group  $W$  is a (finite) group generated by reflections  $s_H$  for  $H \in \mathcal{H}$ .

# Finite Reflection Groups




**Definition:** Let  $\mathcal{H}$  be a set of hyperplanes in  $V$ , we call this a hyperplane arrangement. A (finite) reflection group  $W$  is a (finite) group generated by reflections  $s_H$  for  $H \in \mathcal{H}$ .

**Example:** Let  $V = \mathbb{R}^3$ , and

- $H_1 = \{\mathbf{u} \in \mathbb{R}^3 : \mathbf{u} = (u_1, 0, u_2)\}$
- $H_2 = \{\mathbf{u} \in \mathbb{R}^3 : \mathbf{u} = (u_1, \sqrt{3}u_2, u_2)\}$
- $H_3 = \{\mathbf{u} \in \mathbb{R}^3 : \mathbf{u} = (u_1, \sqrt{3}u_2, -u_2)\}$




# Finite Reflection Groups






# Finite Reflection Groups



We obtain the group:  $W = \langle s_{H_1}, s_{H_2}, s_{H_3} \rangle$ .


# Finite Reflection Groups



We obtain the group:  $W = \langle s_{H_1}, s_{H_2}, s_{H_3} \rangle$ .

(1) For all  $i$ ,  $s_{H_i}^2 = \text{id}_V$


# Finite Reflection Groups



We obtain the group:  $W = \langle s_{H_1}, s_{H_2}, s_{H_3} \rangle$ .

- (1) For all  $i$ ,  $s_{H_i}^2 = \text{id}_V$
- (2) For  $i \neq j$ ,  $(s_{H_i} \circ s_{H_j})^3 = 1$


# Finite Reflection Groups



We obtain the group:  $W = \langle s_{H_1}, s_{H_2}, s_{H_3} \rangle$ .

- (1) For all  $i$ ,  $s_{H_i}^2 = \text{id}_V$
- (2) For  $i \neq j$ ,  $(s_{H_i} \circ s_{H_j})^3 = 1$
- (3) This group can be generated by two elements ( $s_{H_3} = s_{H_1} \circ s_{H_2} \circ s_{H_1}$ ).

# Finite Reflection Groups



We obtain the group:  $W = \langle s_{H_1}, s_{H_2}, s_{H_3} \rangle$ .

- (1) For all  $i$ ,  $s_{H_i}^2 = \text{id}_V$
- (2) For  $i \neq j$ ,  $(s_{H_i} \circ s_{H_j})^3 = 1$
- (3) This group can be generated by two elements ( $s_{H_3} = s_{H_1} \circ s_{H_2} \circ s_{H_1}$ ).
- (4) This group is isomorphic to  $S_3$ .

# Finite Reflection Groups

**Remarks:**

# Finite Reflection Groups

## Remarks:

- (1) In general, each  $s_H$  has order 2 in  $W$ .

# Finite Reflection Groups

## Remarks:

- (1) In general, each  $s_H$  has order 2 in  $W$ .
- (2) Each hyperplane  $H$  is codimension 1, and thus  $H^\perp$  is spanned by some nonzero vector  $\alpha$ .




# Finite Reflection Groups

## Remarks:

- (1) In general, each  $s_H$  has order 2 in  $W$ .
- (2) Each hyperplane  $H$  is codimension 1, and thus  $H^\perp$  is spanned by some nonzero vector  $\alpha$ .

**Example:** For the hyperplane  $H_1$ , we can choose  $\alpha_1 = (0, 1.25, 0)$ .



# Finite Reflection Groups

**Remarks:**

# Finite Reflection Groups

## Remarks:

- (3) For each  $H \in \mathcal{H}$ , associate some  $0 \neq \alpha \in H^\perp$ , the collection of such vectors  $\Phi$  is the (generalized) root system associated to the (Weyl) group  $W =: W_\Phi$ .

# Finite Reflection Groups

## Remarks:

- (3) For each  $H \in \mathcal{H}$ , associate some  $0 \neq \alpha \in H^\perp$ , the collection of such vectors  $\Phi$  is the (generalized) root system associated to the (Weyl) group  $W =: W_\Phi$ .
- (4) For each  $H \in \mathcal{H}$ , we can explicitly write down  $s_H$  in terms of  $\alpha$  (and thus we denote  $s_H$  by  $s_\alpha$ ):

$$s_\alpha(x) = x - 2 \frac{\langle \alpha, x \rangle}{\langle \alpha, \alpha \rangle} \alpha$$

# Finite Reflection Groups

## Remarks:

- (3) For each  $H \in \mathcal{H}$ , associate some  $0 \neq \alpha \in H^\perp$ , the collection of such vectors  $\Phi$  is the (generalized) root system associated to the (Weyl) group  $W =: W_\Phi$ .
- (4) For each  $H \in \mathcal{H}$ , we can explicitly write down  $s_H$  in terms of  $\alpha$  (and thus we denote  $s_H$  by  $s_\alpha$ ):

$$s_\alpha(x) = x - 2 \frac{\langle \alpha, x \rangle}{\langle \alpha, \alpha \rangle} \alpha$$

**Example:** Let  $0 \neq \alpha \in H_1^\perp$ , then

$$s_{\alpha_1}(\mathbf{u}) = (u_1, -u_2, u_3)$$

The function  $s_H$  is well-defined for any choice of  $0 \neq \alpha \in H^\perp$ .

# Finite Reflection Groups

## Remarks:

(5) Let  $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$ , then  $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$ .

# Finite Reflection Groups

## Remarks:

(5) Let  $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$ , then  $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$ . This has a few properties:

# Finite Reflection Groups

## Remarks:

(5) Let  $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$ , then  $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$ . This has a few properties:

$$- \langle \alpha^\vee, \alpha \rangle = \left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \alpha \right\rangle = \frac{2}{\langle \alpha, \alpha \rangle} \langle \alpha, \alpha \rangle = 2$$



# Finite Reflection Groups

## Remarks:

(5) Let  $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$ , then  $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$ . This has a few properties:

$$\begin{aligned} - \langle \alpha^\vee, \alpha \rangle &= \left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \alpha \right\rangle = \frac{2}{\langle \alpha, \alpha \rangle} \langle \alpha, \alpha \rangle = 2 \\ - (\alpha^\vee)^\vee &= \frac{2\left(\frac{2\alpha}{\langle \alpha, \alpha \rangle}\right)}{\left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \frac{2\alpha}{\langle \alpha, \alpha \rangle} \right\rangle} = \frac{\frac{4}{\langle \alpha, \alpha \rangle} \alpha}{\frac{4}{\langle \alpha, \alpha \rangle^2} \langle \alpha, \alpha \rangle} = \alpha. \end{aligned}$$

# Finite Reflection Groups

## Remarks:

(5) Let  $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$ , then  $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$ . This has a few properties:

$$- \langle \alpha^\vee, \alpha \rangle = \left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \alpha \right\rangle = \frac{2}{\langle \alpha, \alpha \rangle} \langle \alpha, \alpha \rangle = 2$$

$$- (\alpha^\vee)^\vee = \frac{2\left(\frac{2\alpha}{\langle \alpha, \alpha \rangle}\right)}{\left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \frac{2\alpha}{\langle \alpha, \alpha \rangle} \right\rangle} = \frac{\frac{4}{\langle \alpha, \alpha \rangle} \alpha}{\frac{4}{\langle \alpha, \alpha \rangle^2} \langle \alpha, \alpha \rangle} = \alpha.$$

A crystallographic root system is a (generalized) root system if  $\Phi$  satisfies:

$$\langle \alpha^\vee, \beta \rangle \in \mathbb{Z}$$

for all  $\alpha, \beta \in \Phi$ . In Lie algebra theory, this condition arises naturally.

# Finite Reflection Groups

## Remarks:

(5) Let  $\alpha^\vee := \frac{2\alpha}{\langle \alpha, \alpha \rangle}$ , then  $s_\alpha(x) = x - \langle \alpha^\vee, x \rangle \alpha$ . This has a few properties:

$$\begin{aligned} - \langle \alpha^\vee, \alpha \rangle &= \left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \alpha \right\rangle = \frac{2}{\langle \alpha, \alpha \rangle} \langle \alpha, \alpha \rangle = 2 \\ - (\alpha^\vee)^\vee &= \frac{2\left(\frac{2\alpha}{\langle \alpha, \alpha \rangle}\right)}{\left\langle \frac{2\alpha}{\langle \alpha, \alpha \rangle}, \frac{2\alpha}{\langle \alpha, \alpha \rangle} \right\rangle} = \frac{\frac{4}{\langle \alpha, \alpha \rangle} \alpha}{\frac{4}{\langle \alpha, \alpha \rangle^2} \langle \alpha, \alpha \rangle} = \alpha. \end{aligned}$$

A crystallographic root system is a (generalized) root system if  $\Phi$  satisfies:

$$\langle \alpha^\vee, \beta \rangle \in \mathbb{Z}$$

for all  $\alpha, \beta \in \Phi$ . In Lie algebra theory, this condition arises naturally. We define  $\Phi^\vee := \{\alpha^\vee : \alpha \in \Phi\}$  to be the coroot system of  $W$ , in particular, a Weyl group can have multiple root systems (the choice of  $\Phi$  is important to the structure).

# Finite Reflection Groups

We have an orthogonal decomposition of  $V$  via the action  $(W, \Phi)$ .

# Finite Reflection Groups

We have an orthogonal decomposition of  $V$  via the action  $(W, \Phi)$ .

$$V_0 := \bigcap_{\alpha \in \Phi} \alpha^\perp$$

# Finite Reflection Groups

We have an orthogonal decomposition of  $V$  via the action  $(W, \Phi)$ .

$$V_0 := \bigcap_{\alpha \in \Phi} \alpha^\perp$$

This is the fixed-point set  $V^W = \{v \in V : wv = v \text{ for all } w \in W\}$ . We call this the *inessential part* of  $V$ .

# Finite Reflection Groups

We have an orthogonal decomposition of  $V$  via the action  $(W, \Phi)$ .

$$V_0 := \bigcap_{\alpha \in \Phi} \alpha^\perp$$

This is the fixed-point set  $V^W = \{v \in V : wv = v \text{ for all } w \in W\}$ . We call this the *inessential part* of  $V$ .

$$V_1 := \text{span}_{\mathbb{R}}(\Phi)$$

# Finite Reflection Groups

We have an orthogonal decomposition of  $V$  via the action  $(W, \Phi)$ .

$$V_0 := \bigcap_{\alpha \in \Phi} \alpha^\perp$$

This is the fixed-point set  $V^W = \{v \in V : wv = v \text{ for all } w \in W\}$ . We call this the *inessential part* of  $V$ .

$$V_1 := \text{span}_{\mathbb{R}}(\Phi)$$

The action of  $W$  is completely determined by  $V_1$ . We call this the *essential part* of  $V$ .



# Finite Reflection Groups

We have an orthogonal decomposition of  $V$  via the action  $(W, \Phi)$ .

$$V_0 := \bigcap_{\alpha \in \Phi} \alpha^\perp$$

This is the fixed-point set  $V^W = \{v \in V : wv = v \text{ for all } w \in W\}$ . We call this the *inessential part* of  $V$ .




$$V_1 := \text{span}_{\mathbb{R}}(\Phi)$$

The action of  $W$  is completely determined by  $V_1$ . We call this the *essential part* of  $V$ .

$$V = V_0 \oplus V_1$$


# Finite Reflection Groups

**Example:** Let  $V = \mathbb{R}^3$ , and  $H_1$ ,  $H_2$  and  $H_3$  as before and choose orthogonal vectors  $\alpha_1 = (0, 1, 0)$ ,  $\alpha_2 = (0, -1, \sqrt{3})$ , and  $\alpha_3 = (0, 1, \sqrt{3})$ , respectively.



# Finite Reflection Groups


**Example:** Let  $V = \mathbb{R}^3$ , and  $H_1$ ,  $H_2$  and  $H_3$  as before and choose orthogonal vectors  $\alpha_1 = (0, 1, 0)$ ,  $\alpha_2 = (0, -1, \sqrt{3})$ , and  $\alpha_3 = (0, 1, \sqrt{3})$ , respectively.




The  $x$ -axis is contained in each hyperplane and thus is orthogonal to each  $\alpha_i$  chosen, or equivalently, is fixed by the reflections across each  $H_i$ :

$$V_0 := (0, 1, 0)^\perp \cap (0, -1, \sqrt{3})^\perp \cap (0, 1, \sqrt{3})^\perp$$


# Finite Reflection Groups



# Finite Reflection Groups



By contracting  $V_0$  to a point, we see the essential part of  $(W, \Phi)$ :



# Finite Reflection Groups

## Remarks:

- (6) One can restrict themselves to studying essential reflection groups, i.e. for  $V = V_0 \oplus V_1$  as above, we require  $V_0 = 0$ .

# Finite Reflection Groups

## Remarks:

- (6) One can restrict themselves to studying essential reflection groups, i.e. for  $V = V_0 \oplus V_1$  as above, we require  $V_0 = 0$ .
- (7) We can define a product  $(W', V') \times (W'', V'') := (W' \times W'', V' \oplus V'')$ . If  $(W, V)$  cannot be expressed as a product, then we say that  $(W, V)$  is irreducible. One can restrict themselves to studying irreducible reflection groups.

# Finite Reflection Groups

## Remarks:

- (6) One can restrict themselves to studying essential reflection groups, i.e. for  $V = V_0 \oplus V_1$  as above, we require  $V_0 = 0$ .
- (7) We can define a product  $(W', V') \times (W'', V'') := (W' \times W'', V' \oplus V'')$ . If  $(W, V)$  cannot be expressed as a product, then we say that  $(W, V)$  is irreducible. One can restrict themselves to studying irreducible reflection groups.
- (8) There is a classification of non-trivial, essential, irreducible, finite reflection groups.



# Finite Reflection Groups

## Remarks:

- (6) One can restrict themselves to studying essential reflection groups, i.e. for  $V = V_0 \oplus V_1$  as above, we require  $V_0 = 0$ .
- (7) We can define a product  $(W', V') \times (W'', V'') := (W' \times W'', V' \oplus V'')$ . If  $(W, V)$  cannot be expressed as a product, then we say that  $(W, V)$  is irreducible. One can restrict themselves to studying irreducible reflection groups.
- (8) There is a classification of non-trivial, essential, irreducible, finite reflection groups.

Among these are: group of permutations on  $n$  letters, group of signed permutations on  $n$  letters, dihedral groups, symmetries of regular solids (see Section 1.3 in Buildings - Theory and Applications, Abramenko and Brown, 2008).

# Finite Reflection Groups

We have introduced quite a bit of notation:

- $V$  is a Euclidean vector space.
- $n$  is the dimension of  $V$ .
- $H$  is a hyperplane, there exists  $0 \neq \alpha \in H^\perp$  called a root.
- $\mathcal{H}$  is a hyperplane arrangement,  $\Phi$  is a generalized root system.
- $s_H$  is the reflection across  $H$ , also denoted  $s_\alpha$ , where  $\alpha$  is a root (determined by  $H$ ).
- $W$  is the Weyl group.
- $V_1 = \text{span}_{\mathbb{R}}(\Phi)$  is the essential part of  $V$ .
- $V_0 = V^W$  is the unessential part of  $V$ .

# Coxeter Complex

**Definition:** A (finite) abstract simplicial complex is a (finite) set  $A$  together with a collection  $\Delta$  of finite subsets of  $A$  such that if  $X \in \Delta$  and  $Y \subseteq X$ , then  $Y \in \Delta$ .

# Coxeter Complex

**Definition:** A (finite) abstract simplicial complex is a (finite) set  $A$  together with a collection  $\Delta$  of finite subsets of  $A$  such that if  $X \in \Delta$  and  $Y \subseteq X$ , then  $Y \in \Delta$ . An abstract simplicial complex is denoted by the collection  $\Delta$ .

# Coxeter Complex

**Definition:** A (finite) abstract simplicial complex is a (finite) set  $A$  together with a collection  $\Delta$  of finite subsets of  $A$  such that if  $X \in \Delta$  and  $Y \subseteq X$ , then  $Y \in \Delta$ . An abstract simplicial complex is denoted by the collection  $\Delta$ .

(1) A vertex is a  $v \in A$  such that  $\{v\} \in \Delta$ .

# Coxeter Complex

**Definition:** A (finite) abstract simplicial complex is a (finite) set  $A$  together with a collection  $\Delta$  of finite subsets of  $A$  such that if  $X \in \Delta$  and  $Y \subseteq X$ , then  $Y \in \Delta$ . An abstract simplicial complex is denoted by the collection  $\Delta$ .

- (1) A vertex is a  $v \in A$  such that  $\{v\} \in \Delta$ .
- (2) The set of vertices is denoted  $V(\Delta)$ .

# Coxeter Complex

**Definition:** A (finite) abstract simplicial complex is a (finite) set  $A$  together with a collection  $\Delta$  of finite subsets of  $A$  such that if  $X \in \Delta$  and  $Y \subseteq X$ , then  $Y \in \Delta$ . An abstract simplicial complex is denoted by the collection  $\Delta$ .

- (1) A vertex is a  $v \in A$  such that  $\{v\} \in \Delta$ .
- (2) The set of vertices is denoted  $V(\Delta)$ .
- (3) The elements of  $\Delta$  are called simplices.

# Coxeter Complex

**Definition:** A (finite) abstract simplicial complex is a (finite) set  $A$  together with a collection  $\Delta$  of finite subsets of  $A$  such that if  $X \in \Delta$  and  $Y \subseteq X$ , then  $Y \in \Delta$ . An abstract simplicial complex is denoted by the collection  $\Delta$ .

- (1) A vertex is a  $v \in A$  such that  $\{v\} \in \Delta$ .
- (2) The set of vertices is denoted  $V(\Delta)$ .
- (3) The elements of  $\Delta$  are called simplices.
- (4) The dimension of a simplex  $\delta \in \Delta$  is  $|\delta| - 1$ , i.e. vertices are zero dimensional, edges are one-dimensional, etc.



# Coxeter Complex

**Definition:** A (finite) abstract simplicial complex is a (finite) set  $A$  together with a collection  $\Delta$  of finite subsets of  $A$  such that if  $X \in \Delta$  and  $Y \subseteq X$ , then  $Y \in \Delta$ . An abstract simplicial complex is denoted by the collection  $\Delta$ .

- (1) A vertex is a  $v \in A$  such that  $\{v\} \in \Delta$ .
- (2) The set of vertices is denoted  $V(\Delta)$ .
- (3) The elements of  $\Delta$  are called simplices.
- (4) The dimension of a simplex  $\delta \in \Delta$  is  $|\delta| - 1$ , i.e. vertices are zero dimensional, edges are one-dimensional, etc.

To simplify notation with subsets, I will denote subsets  $X = \{a_1, a_2, \dots, a_k\}$  by  $X = a_1 a_2 \dots a_k$ .



# Coxeter Complex

**Example:** Let  $A = \{1, 2, 3, 4\}$  and consider

$\Delta_1 = \{\emptyset, 1, 2, 3, 12, 13, 23, 123\}$  and  $\Delta_2 = \{\emptyset, 1, 2, 4, 12, 14\}$ .



# Coxeter Complex

**Example:** Let  $A = \{1, 2, 3, 4\}$  and consider  $\Delta_1 = \{\emptyset, 1, 2, 3, 12, 13, 23, 123\}$  and  $\Delta_2 = \{\emptyset, 1, 2, 4, 12, 14\}$ .




# Coxeter Complex

**Example:** Let  $A = \{1, 2, 3, 4\}$  and consider  $\Delta_1 = \{\emptyset, 1, 2, 3, 12, 13, 23, 123\}$  and  $\Delta_2 = \{\emptyset, 1, 2, 4, 12, 14\}$ .



However,  $\Delta_3 = \{\emptyset, 1, 2, 23, 123\}$  is not an abstract simplicial complex.



# Coxeter Complex

**Definition:** Let  $\Delta_1$  and  $\Delta_2$  be two abstract simplicial complexes. A simplicial map is a function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that for all  $\delta \in \Delta_1$ ,  $f(\delta) \in \Delta_2$ .

# Coxeter Complex


**Definition:** Let  $\Delta_1$  and  $\Delta_2$  be two abstract simplicial complexes. A simplicial map is a function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that for all  $\delta \in \Delta_1$ ,  $f(\delta) \in \Delta_2$ .

**Example:** Let  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  be defined by  $f(1) = 1$  and  $f(2) = 2$  and  $f(3) = 1$ .

# Coxeter Complex

**Definition:** Let  $\Delta_1$  and  $\Delta_2$  be two abstract simplicial complexes. A simplicial map is a function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that for all  $\delta \in \Delta_1$ ,  $f(\delta) \in \Delta_2$ .


**Example:** Let  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  be defined by  $f(1) = 1$  and  $f(2) = 2$  and  $f(3) = 1$ .



# Coxeter Complex

**Definition:** Let  $\Delta_1$  and  $\Delta_2$  be two abstract simplicial complexes. A simplicial map is a function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that for all  $\delta \in \Delta_1$ ,  $f(\delta) \in \Delta_2$ .

**Example:** Let  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  be defined by  $f(1) = 1$  and  $f(2) = 2$  and  $f(3) = 1$ .






# Coxeter Complex

**Definition:** Let  $\Delta_1$  and  $\Delta_2$  be two abstract simplicial complexes. A simplicial map is a function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that for all  $\delta \in \Delta_1$ ,  $f(\delta) \in \Delta_2$ .


**Example:** Let  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  be defined by  $f(1) = 1$  and  $f(2) = 2$  and  $f(3) = 1$ .



# Coxeter Complex

**Definition:** Let  $\Delta_1$  and  $\Delta_2$  be two abstract simplicial complexes. A simplicial map is a function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that for all  $\delta \in \Delta_1$ ,  $f(\delta) \in \Delta_2$ .


**Example:** Let  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  be defined by  $f(1) = 1$  and  $f(2) = 2$  and  $f(3) = 1$ .



# Coxeter Complex

**Definition:** Let  $\Delta_1$  and  $\Delta_2$  be two abstract simplicial complexes. A simplicial map is a function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that for all  $\delta \in \Delta_1$ ,  $f(\delta) \in \Delta_2$ .

**Example:** Let  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  be defined by  $f(1) = 1$  and  $f(2) = 2$  and  $f(3) = 1$ .



# Coxeter Complex

(1) The identity map  $\text{id}_\Delta : \Delta \rightarrow \Delta$  is a simplicial map.

# Coxeter Complex

- (1) The identity map  $\text{id}_\Delta : \Delta \rightarrow \Delta$  is a simplicial map.
- (2) The composition of simplicial maps is simplicial.


# Coxeter Complex

- (1) The identity map  $\text{id}_\Delta : \Delta \rightarrow \Delta$  is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes  $\Delta_1$  and  $\Delta_2$  is a bijective function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that  $f^{-1}$  is simplicial.

# Coxeter Complex

- (1) The identity map  $\text{id}_\Delta : \Delta \rightarrow \Delta$  is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes  $\Delta_1$  and  $\Delta_2$  is a bijective function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that  $f^{-1}$  is simplicial.


**Example:** Let  $f : V(\Delta_2) \rightarrow V(\Delta_1)$  be defined by  $f(1) = 1$ ,  $f(2) = 2$ , and  $f(3) = 4$ .



# Coxeter Complex

- (1) The identity map  $\text{id}_\Delta : \Delta \rightarrow \Delta$  is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes  $\Delta_1$  and  $\Delta_2$  is a bijective function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that  $f^{-1}$  is simplicial.

**Example:** Let  $f : V(\Delta_2) \rightarrow V(\Delta_1)$  be defined by  $f(1) = 1$ ,  $f(2) = 2$ , and  $f(4) = 3$ .






# Coxeter Complex

- (1) The identity map  $\text{id}_\Delta : \Delta \rightarrow \Delta$  is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes  $\Delta_1$  and  $\Delta_2$  is a bijective function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that  $f^{-1}$  is simplicial.


**Example:** Let  $f : V(\Delta_2) \rightarrow V(\Delta_1)$  be defined by  $f(1) = 1$ ,  $f(2) = 2$ , and  $f(4) = 3$ .



# Coxeter Complex

- (1) The identity map  $\text{id}_\Delta : \Delta \rightarrow \Delta$  is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes  $\Delta_1$  and  $\Delta_2$  is a bijective function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that  $f^{-1}$  is simplicial.


**Example:** Let  $f : V(\Delta_2) \rightarrow V(\Delta_1)$  be defined by  $f(1) = 1$ ,  $f(2) = 2$ , and  $f(4) = 3$ .



# Coxeter Complex

- (1) The identity map  $\text{id}_\Delta : \Delta \rightarrow \Delta$  is a simplicial map.
- (2) The composition of simplicial maps is simplicial.
- (3) An isomorphism of simplicial complexes  $\Delta_1$  and  $\Delta_2$  is a bijective function  $f : V(\Delta_1) \rightarrow V(\Delta_2)$  such that  $f^{-1}$  is simplicial.

**Example:** Let  $f : V(\Delta_2) \rightarrow V(\Delta_1)$  be defined by  $f(1) = 1$ ,  $f(2) = 2$ , and  $f(4) = 3$ .




# Coxeter Complex

Our next goal is to construct to a chamber complex associated to an essential finite reflection group  $(W, V)$ , which we will denote by  $\Sigma := \Sigma(W, S)$ .

# Coxeter Complex

Our next goal is to construct a chamber complex associated to an essential finite reflection group  $(W, V)$ , which we will denote by  $\Sigma := \Sigma(W, S)$ . Let's return to our running example:



where  $V = \mathbb{R}^3$  with the usual inner product. We can reformulate this example in terms of the essential part of the action.

# Coxeter Complex

Let  $V = \mathbb{R}^2$ , with the usual inner product. Let  $\mathcal{H} = \{H_1, H_2, H_3\}$ .

- $H_1 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (0, u_2)\}$
- $H_2 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, u_2)\}$
- $H_3 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, -u_2)\}$

# Coxeter Complex

Let  $V = \mathbb{R}^2$ , with the usual inner product. Let  $\mathcal{H} = \{H_1, H_2, H_3\}$ .

- $H_1 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (0, u_2)\}$
- $H_2 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, u_2)\}$
- $H_3 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, -u_2)\}$

Set  $s := s_{H_1}$  and  $t := s_{H_2}$ , then  $s_{H_3} = s \circ t \circ s$ . We will denote the *set of fundamental reflections* by  $S = \{s, t\}$ .

# Coxeter Complex

Let  $V = \mathbb{R}^2$ , with the usual inner product. Let  $\mathcal{H} = \{H_1, H_2, H_3\}$ .

- $H_1 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (0, u_2)\}$
- $H_2 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, u_2)\}$
- $H_3 = \{\mathbf{u} \in \mathbb{R}^2 : \mathbf{u} = (\sqrt{3}u_2, -u_2)\}$

Set  $s := s_{H_1}$  and  $t := s_{H_2}$ , then  $s_{H_3} = s \circ t \circ s$ . We will denote the *set of fundamental reflections* by  $S = \{s, t\}$ . To each hyperplane, we will associate a linear function  $f_i : V \rightarrow \mathbb{R}$  such that  $f_i|_{H_i} = 0$  and  $f_i \neq 0$ .

- $f_1(x, y) = x$
- $f_2(x, y) = -x + \sqrt{3}y$
- $f_3(x, y) = x + \sqrt{3}y$

I will refer to the pair  $(W, S)$  as a Coxeter group.



# Coxeter Complex

**Definition:** A *cell* in  $V$  with respect to  $\mathcal{H} = \{H_i\}_{i \in I}$  is a nonempty set  $A$  obtained by choosing for each  $i$  a sign  $\sigma_i \in \{+, -, 0\}$  and specifying  $f_i = \sigma_i$ .

# Coxeter Complex

**Definition:** A *cell* in  $V$  with respect to  $\mathcal{H} = \{H_i\}_{i \in I}$  is a nonempty set  $A$  obtained by choosing for each  $i$  a sign  $\sigma_i \in \{+, -, 0\}$  and specifying  $f_i = \sigma_i$ . In other words,  $A$  is defined by a set of linear equalities and strict inequalities, one for each hyperplane in  $\mathcal{H}$ .

$$A = \bigcap_{i \in I} U_i$$

for  $U_i$  is either  $H_i$  or an open half-space with  $H_i$  as a boundary. We denote the set of cells  $\Sigma(\mathcal{H})$ .

# Coxeter Complex

**Definition:** A *cell* in  $V$  with respect to  $\mathcal{H} = \{H_i\}_{i \in I}$  is a nonempty set  $A$  obtained by choosing for each  $i$  a sign  $\sigma_i \in \{+, -, 0\}$  and specifying  $f_i = \sigma_i$ . In other words,  $A$  is defined by a set of linear equalities and strict inequalities, one for each hyperplane in  $\mathcal{H}$ .

$$A = \bigcap_{i \in I} U_i$$

for  $U_i$  is either  $H_i$  or an open half-space with  $H_i$  as a boundary. We denote the set of cells  $\Sigma(\mathcal{H})$ .

**Definition:** If each  $\sigma_i \neq 0$ , then we call the cell a *chamber*. We denote the set of chamber by  $\mathcal{C}(\mathcal{H})$ .

# Coxeter Complex

**Definition:** A *cell* in  $V$  with respect to  $\mathcal{H} = \{H_i\}_{i \in I}$  is a nonempty set  $A$  obtained by choosing for each  $i$  a sign  $\sigma_i \in \{+, -, 0\}$  and specifying  $f_i = \sigma_i$ . In other words,  $A$  is defined by a set of linear equalities and strict inequalities, one for each hyperplane in  $\mathcal{H}$ .








$$A = \bigcap_{i \in I} U_i$$

for  $U_i$  is either  $H_i$  or an open half-space with  $H_i$  as a boundary. We denote the set of cells  $\Sigma(\mathcal{H})$ .

**Definition:** If each  $\sigma_i \neq 0$ , then we call the cell a *chamber*. We denote the set of chamber by  $\mathcal{C}(\mathcal{H})$ .

**Definition:** If  $\sigma_i = 0$  for exactly one  $i \in I$ , then we call the cell a *panel*.


# Coxeter Complex




# Coxeter Complex




$$f_1 = 0$$




$$f_1 = +$$




$$f_1 = -$$




$$f_2 = 0$$



$$f_2 = +$$



$$f_2 = -$$



$$f_3 = 0$$




$$f_3 = +$$











$$f_3 = -$$

# Coxeter Complex




# Coxeter Complex





# Coxeter Complex




# Coxeter Complex

**Definition:** Given cells  $A, B \in \Sigma(\mathcal{H})$ ,  $B$  is a face of  $A$ , denoted  $B \leq A$ , if for each  $i \in I$ , either  $\sigma_i(B) = 0$  or  $\sigma_i(B) = \sigma_i(A)$ .

# Coxeter Complex

**Definition:** Given cells  $A, B \in \Sigma(\mathcal{H})$ ,  $B$  is a face of  $A$ , denoted  $B \leq A$ , if for each  $i \in I$ , either  $\sigma_i(B) = 0$  or  $\sigma_i(B) = \sigma_i(A)$ .


**Example:** In our previous examples of cells,  $B \leq A$ .



We say that  $B$  is a panel of  $A$ , and the hyperplane  $H_1$  containing  $B$  is the *wall* of  $A$ .

# Coxeter Complex

In this example, there are 13 cells: 6 chambers, 6 open rays and the origin.



# Coxeter Complex


We distinguish an arbitrary chamber  $C$  in the cell decomposition.

# Coxeter Complex

We distinguish an arbitrary chamber  $C$  in the cell decomposition. Recall  $s = s_{H_1}$  and  $t = s_{H_2}$ .


# Coxeter Complex

We distinguish an arbitrary chamber  $C$  in the cell decomposition. Recall  $s = s_{H_1}$  and  $t = s_{H_2}$ .



# Coxeter Complex

We distinguish an arbitrary chamber  $C$  in the cell decomposition. Recall  $s = s_{H_1}$  and  $t = s_{H_2}$ .



Note  $w_0 = sts = tst$ . By this definition, it is clear  $W$  acts transitively on the chambers and  $|\mathcal{C}(\mathcal{H})| = |W|$ .



# Coxeter Complex

Let  $\Sigma_{\leq C}$  be the subcomplex of faces of  $C$ . For  $A \leq C$ , let  $W_A$  be the stabilizer of  $A$ .

# Coxeter Complex

Let  $\Sigma_{\leq C}$  be the subcomplex of faces of  $C$ . For  $A \leq C$ , let  $W_A$  be the stabilizer of  $A$ .


**Example:** Let  $C$  be the fundamental chamber,  $C$ ,  $R_1^+$ ,  $R_2^+$ , and  $0$  are the faces of  $C$ .

$$W_C = \{e\}$$

$$W_{R_1^+} = \{e, s\}$$

$$W_{R_2^+} = \{e, t\}$$

$$W_0 = W$$



# Coxeter Complex

Note that  $W_{R_1^+} = \langle s \rangle$ ,  $W_{R_2^+} = \langle t \rangle$  and  $W_0 = \langle s, t \rangle$ .

# Coxeter Complex

Note that  $W_{R_1^+} = \langle s \rangle$ ,  $W_{R_2^+} = \langle t \rangle$  and  $W_0 = \langle s, t \rangle$ .


The stabilizers are generated by subsets of the set of fundamental reflections  $S$ , which are called *parabolic subgroups* of  $(W, S)$ .

# Coxeter Complex

Note that  $W_{R_1^+} = \langle s \rangle$ ,  $W_{R_2^+} = \langle t \rangle$  and  $W_0 = \langle s, t \rangle$ .

The stabilizers are generated by subsets of the set of fundamental reflections  $S$ , which are called *parabolic subgroups* of  $(W, S)$ . There is an order-reversing isomorphism of posets:

$$\phi : \Sigma_{\leq C} \xrightarrow{\sim} (\text{parabolic subgroups})^{\text{op}}$$




# Coxeter Complex

Note that  $W_{R_1^+} = \langle s \rangle$ ,  $W_{R_2^+} = \langle t \rangle$  and  $W_0 = \langle s, t \rangle$ .


The stabilizers are generated by subsets of the set of fundamental reflections  $S$ , which are called *parabolic subgroups* of  $(W, S)$ . There is an order-reversing isomorphism of posets:

$$\phi : \Sigma_{\leq C} \xrightarrow{\sim} (\text{parabolic subgroups})^{\text{op}}$$



We can extend this isomorphism (of posets) to the whole poset  $\Sigma$  by including cosets of parabolic subgroups  $\hat{\phi} : \Sigma \xrightarrow{\sim} (\text{parabolic cosets})^{\text{op}}$ .

# Coxeter Complex



# Coxeter Complex

Therefore, using the isomorphism of posets  $\hat{\phi}$ , we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on  $\Sigma(W, S)$ . Via this isomorphism,  $W$  is the underlying set of  $\Sigma$ .



# Coxeter Complex

Therefore, using the isomorphism of posets  $\hat{\phi}$ , we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on  $\Sigma(W, S)$ . Via this isomorphism,  $W$  is the underlying set of  $\Sigma$ .

- The vertices of  $\Sigma$  are  $\{w\}$ , which are the chambers.

# Coxeter Complex

Therefore, using the isomorphism of posets  $\hat{\phi}$ , we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on  $\Sigma(W, S)$ . Via this isomorphism,  $W$  is the underlying set of  $\Sigma$ .

- The vertices of  $\Sigma$  are  $\{w\}$ , which are the chambers.
- The edges of  $\Sigma$  are  $w\langle u \rangle = \{w, ws\}$  which is a panel of  $\{w\}$  (the parabolic cosets of order 2). We say  $w$  and  $wu$  are  $u$ -adjacent.

# Coxeter Complex

Therefore, using the isomorphism of posets  $\hat{\phi}$ , we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on  $\Sigma(W, S)$ . Via this isomorphism,  $W$  is the underlying set of  $\Sigma$ .

- The vertices of  $\Sigma$  are  $\{w\}$ , which are the chambers.
- The edges of  $\Sigma$  are  $w\langle u \rangle = \{w, ws\}$  which is a panel of  $\{w\}$  (the parabolic cosets of order 2). We say  $w$  and  $wu$  are  $u$ -adjacent.

**Remarks:**

# Coxeter Complex

Therefore, using the isomorphism of posets  $\hat{\phi}$ , we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on  $\Sigma(W, S)$ . Via this isomorphism,  $W$  is the underlying set of  $\Sigma$ .

- The vertices of  $\Sigma$  are  $\{w\}$ , which are the chambers.
- The edges of  $\Sigma$  are  $w\langle u \rangle = \{w, ws\}$  which is a panel of  $\{w\}$  (the parabolic cosets of order 2). We say  $w$  and  $wu$  are  $u$ -adjacent.

## Remarks:

- (1)  $\Sigma$  is *thin*, meaning each panel is the face of exactly two chambers.

# Coxeter Complex

Therefore, using the isomorphism of posets  $\hat{\phi}$ , we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on  $\Sigma(W, S)$ . Via this isomorphism,  $W$  is the underlying set of  $\Sigma$ .

- The vertices of  $\Sigma$  are  $\{w\}$ , which are the chambers.
- The edges of  $\Sigma$  are  $w\langle u \rangle = \{w, ws\}$  which is a panel of  $\{w\}$  (the parabolic cosets of order 2). We say  $w$  and  $wu$  are  $u$ -adjacent.

## Remarks:

- (1)  $\Sigma$  is *thin*, meaning each panel is the face of exactly two chambers.
- (2)  $\Sigma$  is *colorable*, meaning we can assign a labeling which distinguishes types of vertices.

# Coxeter Complex

Therefore, using the isomorphism of posets  $\hat{\phi}$ , we can identify

$$\Sigma \cong (\text{parabolic cosets})^{\text{op}}$$

This isomorphism defines a simplicial structure on  $\Sigma(W, S)$ . Via this isomorphism,  $W$  is the underlying set of  $\Sigma$ .

- The vertices of  $\Sigma$  are  $\{w\}$ , which are the chambers.
- The edges of  $\Sigma$  are  $w\langle u \rangle = \{w, ws\}$  which is a panel of  $\{w\}$  (the parabolic cosets of order 2). We say  $w$  and  $wu$  are  $u$ -adjacent.

## Remarks:

- (1)  $\Sigma$  is *thin*, meaning each panel is the face of exactly two chambers.
- (2)  $\Sigma$  is *colorable*, meaning we can assign a labeling which distinguishes types of vertices.
- (3) The action of  $W$  on  $\Sigma$  is *type-preserving*, meaning that the coloring is preserved under the action.

# Coxeter Complex

**Definition:** Any simplicial complex  $\Delta$  isomorphic to  $\Sigma(W, S)$  for some  $(W, S)$  Coxeter group (i.e. a finite reflection group) is a Coxeter complex.

# Coxeter Complex

**Definition:** Any simplicial complex  $\Delta$  isomorphic to  $\Sigma(W, S)$  for some  $(W, S)$  Coxeter group (i.e. a finite reflection group) is a Coxeter complex.

## Definition of a Building

A (*weak*) *building* is a simplicial complex  $\Delta$  that can be expressed as the union of subcomplexes  $\Sigma$  (called *apartments*) satisfying axioms:

- (B0) Each apartment  $\Sigma$  is a Coxeter complex.
- (B1) For any two simplices  $A, B \in \Delta$ , there is an apartment  $\Sigma$  containing both of them.
- (B2) If  $\Sigma$  and  $\Sigma'$  are two apartments containing  $A$  and  $B$ , then there is an isomorphism  $\Sigma \rightarrow \Sigma'$  fixing  $A$  and  $B$  pointwise.



# Coxeter Complex

**Definition:** Any simplicial complex  $\Delta$  isomorphic to  $\Sigma(W, S)$  for some  $(W, S)$  Coxeter group (i.e. a finite reflection group) is a Coxeter complex.

## Definition of a Building

A (weak) *building* is a simplicial complex  $\Delta$  that can be expressed as the union of subcomplexes  $\Sigma$  (called *apartments*) satisfying axioms:

- (B0) Each apartment  $\Sigma$  is a Coxeter complex.
- (B1) For any two simplices  $A, B \in \Delta$ , there is an apartment  $\Sigma$  containing both of them.
- (B2) If  $\Sigma$  and  $\Sigma'$  are two apartments containing  $A$  and  $B$ , then there is an isomorphism  $\Sigma \rightarrow \Sigma'$  fixing  $A$  and  $B$  pointwise.

**Remark:** Coxeter complexes characterize thin buildings with a single apartment.

# Buildings

For a building  $\Delta$ , let  $\mathcal{A}$  be the collection of apartments satisfying the axioms of a building (this is called a *system of apartments* for  $\Delta$ ).

# Buildings

For a building  $\Delta$ , let  $\mathcal{A}$  be the collection of apartments satisfying the axioms of a building (this is called a *system of apartments* for  $\Delta$ ).

## Remarks:

- (1) We do not require a specific system of apartments; however, every building admits a canonical system of apartments.

# Buildings

For a building  $\Delta$ , let  $\mathcal{A}$  be the collection of apartments satisfying the axioms of a building (this is called a *system of apartments* for  $\Delta$ ).

## Remarks:

- (1) We do not require a specific system of apartments; however, every building admits a canonical system of apartments.
- (2) Furthermore, when the apartments are finite Coxeter complexes, there is a unique system of apartments.

# Buildings

For a building  $\Delta$ , let  $\mathcal{A}$  be the collection of apartments satisfying the axioms of a building (this is called a *system of apartments* for  $\Delta$ ).

## Remarks:

- (1) We do not require a specific system of apartments; however, every building admits a canonical system of apartments.
- (2) Furthermore, when the apartments are finite Coxeter complexes, there is a unique system of apartments.
- (3) A building is colorable, and isomorphisms between buildings can be taken to be type-preserving.

# Buildings

For a building  $\Delta$ , let  $\mathcal{A}$  be the collection of apartments satisfying the axioms of a building (this is called a *system of apartments* for  $\Delta$ ).

## Remarks:

- (1) We do not require a specific system of apartments; however, every building admits a canonical system of apartments.
- (2) Furthermore, when the apartments are finite Coxeter complexes, there is a unique system of apartments.
- (3) A building is colorable, and isomorphisms between buildings can be taken to be type-preserving.
- (4) All apartments are Coxeter complexes for the same Coxeter group.

# Buildings

**Definition:** Let  $P$  be a set with a binary relation  $\mathcal{I}$  called “incidence” such that  $\mathcal{I}$  is reflexive and symmetric. A *flag* in  $P$  is a set of pairwise incident elements of  $P$ .

# Buildings

**Definition:** Let  $P$  be a set with a binary relation  $\mathcal{I}$  called “incidence” such that  $\mathcal{I}$  is reflexive and symmetric. A *flag* in  $P$  is a set of pairwise incident elements of  $P$ .

**Note:** If  $P$  is a poset, then a flag is a linearly ordered subset of  $P$ .

We will assume that  $P$  is partitioned into nonempty subsets  $P_0, P_1, \dots, P_{n-1}$ , where  $p \in P_i$  is said to have type  $i$ .



# Buildings

**Example:** Let  $V$  be a finite dimension  $F$ -vector space (dimension  $n \geq 2$ ).

# Buildings

**Example:** Let  $V$  be a finite dimension  $F$ -vector space (dimension  $n \geq 2$ ). Let  $P$  be the collection of nonzero proper subspaces of  $V$  and let two subspaces be “incident” if one is contained in the other.

# Buildings

**Example:** Let  $V$  be a finite dimension  $F$ -vector space (dimension  $n \geq 2$ ). Let  $P$  be the collection of nonzero proper subspaces of  $V$  and let two subspaces be “incident” if one is contained in the other. Let  $\Delta(V)$  be the flag complex consisting of simplices as chains of subspaces

$$V_1 < V_2 < \cdots < V_k$$

where  $\dim_F(V_i) = i$ . *Chambers* are chains:

$$V_1 < V_2 < \cdots < V_{n-1}$$

# Buildings

**Example:** Let  $V$  be a finite dimension  $F$ -vector space (dimension  $n \geq 2$ ). Let  $P$  be the collection of nonzero proper subspaces of  $V$  and let two subspaces be “incident” if one is contained in the other. Let  $\Delta(V)$  be the flag complex consisting of simplices as chains of subspaces

$$V_1 < V_2 < \cdots < V_k$$

where  $\dim_F(V_i) = i$ . *Chambers* are chains:

$$V_1 < V_2 < \cdots < V_{n-1}$$

Let  $\mathcal{F} = \{L_1, \dots, L_n\}$  be a collection of 1-dimensional subspaces which span  $V$ . The subcomplex  $\Sigma(\mathcal{F})$  of flags is an *apartment*.

# Buildings

**Example:** Let  $V$  be a finite dimension  $F$ -vector space (dimension  $n \geq 2$ ). Let  $P$  be the collection of nonzero proper subspaces of  $V$  and let two subspaces be “incident” if one is contained in the other. Let  $\Delta(V)$  be the flag complex consisting of simplices as chains of subspaces

$$V_1 < V_2 < \cdots < V_k$$

where  $\dim_F(V_i) = i$ . *Chambers* are chains:

$$V_1 < V_2 < \cdots < V_{n-1}$$

Let  $\mathcal{F} = \{L_1, \dots, L_n\}$  be a collection of 1-dimensional subspaces which span  $V$ . The subcomplex  $\Sigma(\mathcal{F})$  of flags is an *apartment*.

**Proof:**  $S_n$  acts on each  $\Sigma(\mathcal{F})$ , each chain is contained in a composition series, and apply Jordan-Hölder theorem to obtain canonical isomorphisms and projections.

# Buildings

Consider  $V = \mathbb{F}_2^3$ .

# Buildings

Consider  $V = \mathbb{F}_2^3$ . Set

$$\begin{array}{llll} L_0 := \mathbb{F}_2(0, 0, 0) & L_1 := \mathbb{F}_2(1, 0, 0) & L_2 := \mathbb{F}_2(0, 1, 0) & L_3 := \mathbb{F}_2(0, 0, 1) \\ L_4 := \mathbb{F}_2(1, 1, 0) & L_5 := \mathbb{F}_2(1, 0, 1) & L_6 := \mathbb{F}_2(0, 1, 1) & L_7 := \mathbb{F}_2(1, 1, 1) \end{array}$$

# Buildings

Consider  $V = \mathbb{F}_2^3$ . Set

$$\begin{aligned} L_0 &:= \mathbb{F}_2(0, 0, 0) & L_1 &:= \mathbb{F}_2(1, 0, 0) & L_2 &:= \mathbb{F}_2(0, 1, 0) & L_3 &:= \mathbb{F}_2(0, 0, 1) \\ L_4 &:= \mathbb{F}_2(1, 1, 0) & L_5 &:= \mathbb{F}_2(1, 0, 1) & L_6 &:= \mathbb{F}_2(0, 1, 1) & L_7 &:= \mathbb{F}_2(1, 1, 1) \end{aligned}$$

Note that  $L_0$  is not a 1-dimensional subspace. Therefore, we have seven 1-dimensional subspaces, which we will call “type 1” vertices.



# Buildings

Consider  $V = \mathbb{F}_2^3$ . Set

$$\begin{aligned} L_0 &:= \mathbb{F}_2(0, 0, 0) & L_1 &:= \mathbb{F}_2(1, 0, 0) & L_2 &:= \mathbb{F}_2(0, 1, 0) & L_3 &:= \mathbb{F}_2(0, 0, 1) \\ L_4 &:= \mathbb{F}_2(1, 1, 0) & L_5 &:= \mathbb{F}_2(1, 0, 1) & L_6 &:= \mathbb{F}_2(0, 1, 1) & L_7 &:= \mathbb{F}_2(1, 1, 1) \end{aligned}$$

Note that  $L_0$  is not a 1-dimensional subspace. Therefore, we have seven 1-dimensional subspaces, which we will call “type 1” vertices. Set

$$\begin{aligned} V_1 &:= L_1 + L_2 & V_2 &:= L_1 + L_3 & V_3 &:= L_1 + L_4 & V_4 &:= L_2 + L_3 \\ V_5 &:= L_2 + L_5 & V_6 &:= L_3 + L_4 & V_7 &:= L_4 + L_5 \end{aligned}$$

Therefore, we have seven 2-dimensional subspaces, which we will call “type 2” vertices.

# Buildings

Consider  $V = \mathbb{F}_2^3$ . Set

$$\begin{aligned} L_0 &:= \mathbb{F}_2(0, 0, 0) & L_1 &:= \mathbb{F}_2(1, 0, 0) & L_2 &:= \mathbb{F}_2(0, 1, 0) & L_3 &:= \mathbb{F}_2(0, 0, 1) \\ L_4 &:= \mathbb{F}_2(1, 1, 0) & L_5 &:= \mathbb{F}_2(1, 0, 1) & L_6 &:= \mathbb{F}_2(0, 1, 1) & L_7 &:= \mathbb{F}_2(1, 1, 1) \end{aligned}$$

Note that  $L_0$  is not a 1-dimensional subspace. Therefore, we have seven 1-dimensional subspaces, which we will call “type 1” vertices. Set


$$\begin{aligned} V_1 &:= L_1 + L_2 & V_2 &:= L_1 + L_3 & V_3 &:= L_1 + L_4 & V_4 &:= L_2 + L_3 \\ V_5 &:= L_2 + L_5 & V_6 &:= L_3 + L_4 & V_7 &:= L_4 + L_5 \end{aligned}$$

Therefore, we have seven 2-dimensional subspaces, which we will call “type 2” vertices. Note that each  $V_i$  contains exactly three subspaces.

$$V_1 = L_1 + L_2 = L_1 + L_4 = L_2 + L_4$$


# Buildings

For each  $\Sigma(\{L_i, L_j, L_k\})$  such that  $V = L_i \oplus L_j \oplus L_k$ , we obtain an apartment:



# Buildings

For each  $\Sigma(\{L_i, L_j, L_k\})$  such that  $V = L_i \oplus L_j \oplus L_k$ , we obtain an apartment:




Chambers are edges with a type 1 and type 2 vertex,  $S_3$  acts by permuting the labels  $\{i, j, k\}$ .

# Buildings

The axioms **B1** and **B2** of a building give us criteria of how to glue.


# Buildings

The axioms **B1** and **B2** of a building give us criteria of how to glue.



# Buildings

The axioms **B1** and **B2** of a building give us criteria of how to glue.



The apartment  $\Sigma(\{L_1, L_2, L_3\})$  is shown above.

## Important Question:



# Buildings

**Important Question:** What exactly was Jacques Tits trying to construct?

# Buildings

**Important Question:** What exactly was Jacques Tits trying to construct? Consider the group  $\mathrm{SL}(2, \mathbb{C})$ , it has a maximal compact subgroup  $\mathrm{SU}(2)$ .

**Important Question:** What exactly was Jacques Tits trying to construct? Consider the group  $SL(2, \mathbb{C})$ , it has a maximal compact subgroup  $SU(2)$ . Note that  $SL(2, \mathbb{C})/SU(2) \cong \mathbb{H} = \{z \in \mathbb{C} : \text{Im}(z) > 0\}$  via the map

$$SL(2, \mathbb{C})/SU(2) \rightarrow \mathbb{H}$$
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \frac{ai + b}{ci + d}$$

**Important Question:** What exactly was Jacques Tits trying to construct? Consider the group  $SL(2, \mathbb{C})$ , it has a maximal compact subgroup  $SU(2)$ . Note that  $SL(2, \mathbb{C})/SU(2) \cong \mathbb{H} = \{z \in \mathbb{C} : \text{Im}(z) > 0\}$  via the map

$$SL(2, \mathbb{C})/SU(2) \rightarrow \mathbb{H}$$
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \frac{ai + b}{ci + d}$$

And in general, if we take a semisimple group  $G$  over  $\mathbb{R}$  and quotient by its maximal compact subgroup  $K$ ,  $G/K$  is a nice geometric space.

# Buildings

**Important Question:** What exactly was Jacques Tits trying to construct? Consider the group  $\mathrm{SL}(2, \mathbb{C})$ , it has a maximal compact subgroup  $\mathrm{SU}(2)$ . Note that  $\mathrm{SL}(2, \mathbb{C})/\mathrm{SU}(2) \cong \mathbb{H} = \{z \in \mathbb{C} : \mathrm{Im}(z) > 0\}$  via the map

$$\mathrm{SL}(2, \mathbb{C})/\mathrm{SU}(2) \rightarrow \mathbb{H}$$
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \frac{ai + b}{ci + d}$$

And in general, if we take a semisimple group  $G$  over  $\mathbb{R}$  and quotient by its maximal compact subgroup  $K$ ,  $G/K$  is a nice geometric space.

However, if we consider a local field  $F = \mathbb{Q}_p$  and the algebraic group  $\mathrm{SL}(2, F)$ , its maximal compact subgroup is  $\mathrm{SL}(2, \mathcal{O}_F)$  and  $\mathrm{SL}(2, F)/\mathrm{SL}(2, \mathcal{O}_F)$ , it is a totally disconnected topological space.

# Buildings

**Important Question:** What exactly was Jacques Tits trying to construct? Consider the group  $\mathrm{SL}(2, \mathbb{C})$ , it has a maximal compact subgroup  $\mathrm{SU}(2)$ . Note that  $\mathrm{SL}(2, \mathbb{C})/\mathrm{SU}(2) \cong \mathbb{H} = \{z \in \mathbb{C} : \mathrm{Im}(z) > 0\}$  via the map

$$\mathrm{SL}(2, \mathbb{C})/\mathrm{SU}(2) \rightarrow \mathbb{H}$$
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \frac{ai + b}{ci + d}$$

And in general, if we take a semisimple group  $G$  over  $\mathbb{R}$  and quotient by its maximal compact subgroup  $K$ ,  $G/K$  is a nice geometric space.

However, if we consider a local field  $F = \mathbb{Q}_p$  and the algebraic group  $\mathrm{SL}(2, F)$ , its maximal compact subgroup is  $\mathrm{SL}(2, \mathcal{O}_F)$  and  $\mathrm{SL}(2, F)/\mathrm{SL}(2, \mathcal{O}_F)$ , it is a totally disconnected topological space. It is a *pretty lame space*.

# Buildings

This occurs as a property of non-archimedean fields, i.e. the natural numbers are not bounded under the absolute value on the field.

# Buildings

This occurs as a property of non-archimedean fields, i.e. the natural numbers are not bounded under the absolute value on the field.

In the case of  $SL(2, \mathbb{C})$ , we obtain a Riemannian symmetric space and we want an analogue to this when the field is non-archimedean.



# Buildings

This occurs as a property of non-archimedean fields, i.e. the natural numbers are not bounded under the absolute value on the field.

In the case of  $SL(2, \mathbb{C})$ , we obtain a Riemannian symmetric space and we want an analogue to this when the field is non-archimedean. A Bruhat-Tits building is the correct analogue. Historically, the following building appeared as a way to prove that every torsion-free subgroup of  $SL(2, \mathbb{Q}_p)$  is a free group (Ihara, 1966).

# Buildings

This occurs as a property of non-archimedean fields, i.e. the natural numbers are not bounded under the absolute value on the field.

In the case of  $SL(2, \mathbb{C})$ , we obtain a Riemannian symmetric space and we want an analogue to this when the field is non-archimedean. A Bruhat-Tits building is the correct analogue. Historically, the following building appeared as a way to prove that every torsion-free subgroup of  $SL(2, \mathbb{Q}_p)$  is a free group (Ihara, 1966). To make the construction concrete, we'll use  $SL(2, \mathbb{Q}_p)$  as our algebraic group.

# Buildings

This occurs as a property of non-archimedean fields, i.e. the natural numbers are not bounded under the absolute value on the field.

In the case of  $SL(2, \mathbb{C})$ , we obtain a Riemannian symmetric space and we want an analogue to this when the field is non-archimedean. A Bruhat-Tits building is the correct analogue. Historically, the following building appeared as a way to prove that every torsion-free subgroup of  $SL(2, \mathbb{Q}_p)$  is a free group (Ihara, 1966). To make the construction concrete, we'll use  $SL(2, \mathbb{Q}_p)$  as our algebraic group.

**Note:** Any element of  $\mathbb{Q}_p$  is of the form:

$$x = \sum_{i=k}^{\infty} a_i p^i$$

for  $0 \leq a_i < p$ . The valuation of  $x$ , is  $\nu(x) = k$ , where  $k$  is the smallest index such that  $a_k \neq 0$ .

**Definition:** A non-archimedean absolute value on a field  $F$  is a map

$$| - | : F \rightarrow \mathbb{R}$$

such that

- (1)  $|x| \geq 0$  for all  $x \in F$ , and  $|x| = 0$  if and only if  $x = 0$ .
- (2)  $|xy| = |x||y|$  for all  $x, y \in F$
- (3)  $|x + y| \leq \max\{|x|, |y|\}$  for all  $x, y \in F$ .

**Definition:** A non-archimedean absolute value on a field  $F$  is a map

$$| - | : F \rightarrow \mathbb{R}$$

such that

- (1)  $|x| \geq 0$  for all  $x \in F$ , and  $|x| = 0$  if and only if  $x = 0$ .
- (2)  $|xy| = |x||y|$  for all  $x, y \in F$
- (3)  $|x + y| \leq \max\{|x|, |y|\}$  for all  $x, y \in F$ .

**Example:** For  $\mathbb{Q}_p$ , the absolute value is defined by  $|x|_p = p^{-\nu(x)}$ .

# Buildings

**Definition:** A non-archimedean absolute value on a field  $F$  is a map

$$| - | : F \rightarrow \mathbb{R}$$

such that

- (1)  $|x| \geq 0$  for all  $x \in F$ , and  $|x| = 0$  if and only if  $x = 0$ .
- (2)  $|xy| = |x||y|$  for all  $x, y \in F$
- (3)  $|x + y| \leq \max\{|x|, |y|\}$  for all  $x, y \in F$ .

**Example:** For  $\mathbb{Q}_p$ , the absolute value is defined by  $|x|_p = p^{-\nu(x)}$ .

**Important Note:** The image of  $\nu(x)$  is an integer and thus  $\nu(\mathbb{Q}_p^\times) = \mathbb{Z}$  is discrete. Or, in particular importance for us,  $\log(|\mathbb{Q}_p^\times|)$  is discrete. And,  $\mathcal{O} = \{x \in \mathbb{Q}_p : |x|_p \leq 1\}$ .

# Buildings

Instead of choosing the maximal compact subgroup, we look at tori. The torus will be

$$T = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} : a \in \mathbb{Q}_p^\times \right\}$$

# Buildings


Instead of choosing the maximal compact subgroup, we look at tori. The torus will be

$$T = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} : a \in \mathbb{Q}_p^\times \right\}$$

Let  $A(T)$  be the apartment corresponding to  $T$ , it is an  $\mathbb{R}$  affine space with simplicial structure defined by the map

$$T \rightarrow \mathbb{R}$$

$$\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \mapsto \log(|a|_p)$$






# Buildings

Instead of choosing the maximal compact subgroup, we look at tori. The torus will be

$$T = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} : a \in \mathbb{Q}_p^\times \right\}$$

Let  $A(T)$  be the apartment corresponding to  $T$ , it is an  $\mathbb{R}$  affine space with simplicial structure defined by the map

$$\begin{aligned} T &\rightarrow \mathbb{R} \\ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} &\mapsto \log(|a|_p) \end{aligned}$$


Our collection of vertices are of the form:  $n \log(p)$  for some  $n \in \mathbb{Z}$ .

# Buildings

We then take the collection of tori  $\{gTg^{-1} : g \in \mathrm{SL}_2(\mathbb{Q}_p)\}$  and glue together the corresponding system of apartments

$$\mathcal{A} = \{A(gTg^{-1}) : g \in \mathrm{SL}_2(\mathbb{Q}_p)\}$$

# Buildings

We then take the collection of tori  $\{gTg^{-1} : g \in \mathrm{SL}_2(\mathbb{Q}_p)\}$  and glue together the corresponding system of apartments

$$\mathcal{A} = \{A(gTg^{-1}) : g \in \mathrm{SL}_2(\mathbb{Q}_p)\}$$


The stabilizers of each vertex is of the form

$$\mathrm{Stab}(\log(|a|_p)) = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \cdot \mathrm{SL}_2(\mathcal{O}) \cdot \begin{pmatrix} a^{-1} & 0 \\ 0 & a \end{pmatrix}$$

and for any two apartments  $A(T)$  and  $g \cdot A(T) = A(gTg^{-1})$ , we glue at  $n \log(p)$  if and only if  $g \in \mathrm{Stab}(n \log(p))$ .


# Buildings

**Example:** Let  $g = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ . For  $a = up^{-n}$ , note that  $g \in \text{Stab}(n \log(p))$  if and only if  $|a|_p \geq 1$  if and only if  $n \log(p) \geq 0$ .




# Buildings

**Example:** Let  $g = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ . For  $a = up^{-n}$ , note that  $g \in \text{Stab}(n \log(p))$  if and only if  $|a|_p \geq 1$  if and only if  $n \log(p) \geq 0$ .




# Buildings

Assume that  $p = 2$ , then the building is given by:



# Buildings


Assume that  $p = 2$ , then the building is given by:



An apartment  $A(gTg^{-1})$  is given by an infinite path along the tree.

# Buildings


Similarly for  $p = 3$ , then the building is given by:





# Buildings

Similarly for  $p = 3$ , then the building is given by:



Again, an apartment  $A(gTg^{-1})$  is given by an infinite path along the tree.

# Bibliography

-  Peter Abramenko, Kenneth S. Brown. *Buildings - Theory and Applications*. Springer Science+Business Media, LLC, 2008.
-  Jean-Pierre Serre. *Trees*. Springer-Verlag Berlin Heidelberg, 1980.  
(Translated by John Stillwell)
-  Annette Werner. *Buildings and tropical geometry [Video]*. Mathematical Sciences Research Institute, 2009.
-  Annette Werner. *A tropical view on Bruhat-Tits buildings and their compactifications*. Central European Journal of Mathematics, 2010.
-  Bertrand Rémy, Amaury Thuillier, Annette Werner. *Bruhat-Tits Buildings and Analytic Geometry* Springer International Publishing Switzerland, 2015.
-  Federico Ardila. *Coxeter groups [Online Lecture Series]*. San Francisco State University, 2008.
-  Dmitry Kozlov. *Combinatorial Algebraic Topology*. Springer-Verlag Berlin Heidelberg, 2008.