

# Fundamental Groups

Topological, Étale, Tannakian

UCSC Graduate Colloquium

---

Deewang Bhamidipati

11<sup>th</sup> October 2021

# Table of contents

## 1. Galois Theory

Quick Review

Infinite Galois Extensions

Grothendieck's formulation of Galois Theory

## 2. Topological Fundamental Group

Galois Covers

Topological Fundamental Group

## 3. Étale Fundamental Group

Étale Covers

Étale Fundamental Group

## 4. Tannakian Fundamental Group

Representations of Affine Group Schemes

Tensor Categories

Tannakian Fundamental Group

# Galois Theory

× × ×

# Quick Review: Field Extensions

## Algebraic Extensions

Let  $k$  be a field. An extension  $L|k$ , i.e. a field  $L$  with  $k \rightarrow L$ , is called *algebraic* if every element  $\alpha$  of  $L$  is a root of some polynomial with coefficients in  $k$ .

# Quick Review: Field Extensions

## Algebraic Extensions

Let  $k$  be a field. An extension  $L|k$ , i.e. a field  $L$  with  $k \rightarrow L$ , is called *algebraic* if every element  $\alpha$  of  $L$  is a root of some polynomial with coefficients in  $k$ . If this polynomial is monic and irreducible over  $k$ , it is called the minimal polynomial of  $\alpha$ .

# Quick Review: Field Extensions

## Algebraic Extensions

Let  $k$  be a field. An extension  $L|k$ , i.e. a field  $L$  with  $k \rightarrow L$ , is called *algebraic* if every element  $\alpha$  of  $L$  is a root of some polynomial with coefficients in  $k$ . If this polynomial is monic and irreducible over  $k$ , it is called the minimal polynomial of  $\alpha$ .

## Degree

The degree of an extension  $L|k$ , denoted by  $[L : k]$ , is its dimension as a  $k$ -vector space.

# Quick Review: Field Extensions

## Algebraic Extensions

Let  $k$  be a field. An extension  $L|k$ , i.e. a field  $L$  with  $k \rightarrow L$ , is called *algebraic* if every element  $\alpha$  of  $L$  is a root of some polynomial with coefficients in  $k$ . If this polynomial is monic and irreducible over  $k$ , it is called the minimal polynomial of  $\alpha$ .

## Degree

The degree of an extension  $L|k$ , denoted by  $[L : k]$ , is its dimension as a  $k$ -vector space. A finite extension  $L$  of  $k$ , that is  $[L : k] < \infty$ , is algebraic.

# Quick Review: Field Extensions

## Algebraic Extensions

Let  $k$  be a field. An extension  $L|k$ , i.e. a field  $L$  with  $k \rightarrow L$ , is called *algebraic* if every element  $\alpha$  of  $L$  is a root of some polynomial with coefficients in  $k$ . If this polynomial is monic and irreducible over  $k$ , it is called the minimal polynomial of  $\alpha$ .

## Degree

The degree of an extension  $L|k$ , denoted by  $[L : k]$ , is its dimension as a  $k$ -vector space. A finite extension  $L$  of  $k$ , that is  $[L : k] < \infty$ , is algebraic.

## Separable Extensions

A polynomial  $f \in k[x]$  is *separable* if its roots, in some algebraic closure of  $k$ , are distinct.



# Quick Review: Field Extensions

## Algebraic Extensions

Let  $k$  be a field. An extension  $L|k$ , i.e. a field  $L$  with  $k \rightarrow L$ , is called *algebraic* if every element  $\alpha$  of  $L$  is a root of some polynomial with coefficients in  $k$ . If this polynomial is monic and irreducible over  $k$ , it is called the minimal polynomial of  $\alpha$ .

## Degree

The degree of an extension  $L|k$ , denoted by  $[L : k]$ , is its dimension as a  $k$ -vector space. A finite extension  $L$  of  $k$ , that is  $[L : k] < \infty$ , is algebraic.

## Separable Extensions

A polynomial  $f \in k[x]$  is *separable* if its roots, in some algebraic closure of  $k$ , are distinct.

An element of an algebraic extension  $L|k$  is *separable* over  $k$  if its minimal polynomial is separable; the extension itself is called *separable* if every element of  $L$  is separable over  $k$ .

# Quick Review: Field Extensions

## Algebraic Closure

A field is *algebraically closed* if it has no algebraic extensions other than itself. An *algebraic closure* of  $k$  is an algebraic extension  $\bar{k}$  that is algebraically closed.

# Quick Review: Field Extensions

## Algebraic Closure

A field is *algebraically closed* if it has no algebraic extensions other than itself. An *algebraic closure* of  $k$  is an algebraic extension  $\bar{k}$  that is algebraically closed. There exists an algebraic closure  $\bar{k}$  of  $k$ . It is unique up to (non-unique) isomorphism.

# Quick Review: Field Extensions

## Algebraic Closure

A field is *algebraically closed* if it has no algebraic extensions other than itself. An *algebraic closure* of  $k$  is an algebraic extension  $\bar{k}$  that is algebraically closed. There exists an algebraic closure  $\bar{k}$  of  $k$ . It is unique up to (non-unique) isomorphism.

## Separable Closure

The compositum of all finite separable subextensions of  $\bar{k}$  is a separable extension  $k_s|k$  containing each finite separable subextension of  $\bar{k}|k$ .

# Quick Review: Field Extensions

## Algebraic Closure

A field is *algebraically closed* if it has no algebraic extensions other than itself. An *algebraic closure* of  $k$  is an algebraic extension  $\bar{k}$  that is algebraically closed. There exists an algebraic closure  $\bar{k}$  of  $k$ . It is unique up to (non-unique) isomorphism.

## Separable Closure

The compositum of all finite separable subextensions of  $\bar{k}$  is a separable extension  $k_s|k$  containing each finite separable subextension of  $\bar{k}|k$ .

The extension  $k_s$  is called the separable closure of  $k$  in  $\bar{k}$ .

# Quick Review: Field Extensions

## Algebraic Closure

A field is *algebraically closed* if it has no algebraic extensions other than itself. An *algebraic closure* of  $k$  is an algebraic extension  $\bar{k}$  that is algebraically closed. There exists an algebraic closure  $\bar{k}$  of  $k$ . It is unique up to (non-unique) isomorphism.

## Separable Closure

The compositum of all finite separable subextensions of  $\bar{k}$  is a separable extension  $k_s|k$  containing each finite separable subextension of  $\bar{k}|k$ .

The extension  $k_s$  is called the separable closure of  $k$  in  $\bar{k}$ .

From now on by "a separable closure of  $k$ " we shall mean its separable closure in some chosen algebraic closure.

# Quick Review: Galois Extensions

## Galois Extensions

Given an extension  $L$  of  $k$ , denote by  $\text{Aut}(L|k)$  the group of field automorphisms of  $L$  fixing  $k$  elementwise.

# Quick Review: Galois Extensions

## Galois Extensions

Given an extension  $L$  of  $k$ , denote by  $\text{Aut}(L|k)$  the group of field automorphisms of  $L$  fixing  $k$  elementwise.

The set of elements of  $L$  that are fixed by the action of  $\text{Aut}(L|k)$ , denoted  $L^{\text{Aut}(L|k)}$ , forms a field extension of  $k$ . So  $k \rightarrow L$  factors as

$$\begin{array}{ccc} L^{\text{Aut}(L|k)} & \xrightarrow{\quad} & L \\ & \nwarrow \quad \nearrow & \\ & k & \end{array}$$



# Quick Review: Galois Extensions

## Galois Extensions

Given an extension  $L$  of  $k$ , denote by  $\text{Aut}(L|k)$  the group of field automorphisms of  $L$  fixing  $k$  elementwise.

The set of elements of  $L$  that are fixed by the action of  $\text{Aut}(L|k)$ , denoted  $L^{\text{Aut}(L|k)}$ , forms a field extension of  $k$ . So  $k \rightarrow L$  factors as

$$\begin{array}{ccc} L^{\text{Aut}(L|k)} & \xrightarrow{\quad} & L \\ & \nwarrow \quad \nearrow & \\ & k & \end{array}$$

An algebraic extension  $L$  of  $k$  is called a *Galois extension* of  $k$  if  $L^{\text{Aut}(L|k)} = k$ .

# Quick Review: Galois Extensions

## Galois Extensions

Given an extension  $L$  of  $k$ , denote by  $\text{Aut}(L|k)$  the group of field automorphisms of  $L$  fixing  $k$  elementwise.

The set of elements of  $L$  that are fixed by the action of  $\text{Aut}(L|k)$ , denoted  $L^{\text{Aut}(L|k)}$ , forms a field extension of  $k$ . So  $k \rightarrow L$  factors as

$$\begin{array}{ccc} L^{\text{Aut}(L|k)} & \xrightarrow{\quad} & L \\ & \nwarrow \quad \nearrow & \\ & k & \end{array}$$

An algebraic extension  $L$  of  $k$  is called a *Galois extension* of  $k$  if  $L^{\text{Aut}(L|k)} = k$ .

In this case  $\text{Aut}(L|k)$  is denoted by  $\text{Gal}(L|k)$ , and called the *Galois group* of  $L$  over  $k$ .

# Quick Review: Galois Extensions

## Galois Extensions

Given an extension  $L$  of  $k$ , denote by  $\text{Aut}(L|k)$  the group of field automorphisms of  $L$  fixing  $k$  elementwise.

The set of elements of  $L$  that are fixed by the action of  $\text{Aut}(L|k)$ , denoted  $L^{\text{Aut}(L|k)}$ , forms a field extension of  $k$ . So  $k \rightarrow L$  factors as

$$\begin{array}{ccc} L^{\text{Aut}(L|k)} & \xrightarrow{\quad} & L \\ & \nwarrow \quad \nearrow & \\ & k & \end{array}$$

An algebraic extension  $L$  of  $k$  is called a *Galois extension* of  $k$  if  $L^{\text{Aut}(L|k)} = k$ .

In this case  $\text{Aut}(L|k)$  is denoted by  $\text{Gal}(L|k)$ , and called the *Galois group* of  $L$  over  $k$ .

A Galois extension  $L|k$  is separable.

# Quick Review: Galois Extensions

## Galois Extensions

Given an extension  $L$  of  $k$ , denote by  $\text{Aut}(L|k)$  the group of field automorphisms of  $L$  fixing  $k$  elementwise.

The set of elements of  $L$  that are fixed by the action of  $\text{Aut}(L|k)$ , denoted  $L^{\text{Aut}(L|k)}$ , forms a field extension of  $k$ . So  $k \rightarrow L$  factors as

$$\begin{array}{ccc} L^{\text{Aut}(L|k)} & \xrightarrow{\quad} & L \\ & \nwarrow \quad \nearrow & \\ & k & \end{array}$$

An algebraic extension  $L$  of  $k$  is called a *Galois extension* of  $k$  if  $L^{\text{Aut}(L|k)} = k$ .

In this case  $\text{Aut}(L|k)$  is denoted by  $\text{Gal}(L|k)$ , and called the *Galois group* of  $L$  over  $k$ .

A Galois extension  $L|k$  is separable.

## Absolute Galois group

A separable closure  $k_s$  of a field  $k$  is always a Galois extension.

# Quick Review: Galois Extensions

## Galois Extensions

Given an extension  $L$  of  $k$ , denote by  $\text{Aut}(L|k)$  the group of field automorphisms of  $L$  fixing  $k$  elementwise.

The set of elements of  $L$  that are fixed by the action of  $\text{Aut}(L|k)$ , denoted  $L^{\text{Aut}(L|k)}$ , forms a field extension of  $k$ . So  $k \rightarrow L$  factors as

$$\begin{array}{ccc} L^{\text{Aut}(L|k)} & \xrightarrow{\quad} & L \\ & \nwarrow \quad \nearrow & \\ & k & \end{array}$$

An algebraic extension  $L$  of  $k$  is called a *Galois extension* of  $k$  if  $L^{\text{Aut}(L|k)} = k$ .

In this case  $\text{Aut}(L|k)$  is denoted by  $\text{Gal}(L|k)$ , and called the *Galois group* of  $L$  over  $k$ .

A Galois extension  $L|k$  is separable.

## Absolute Galois group

A separable closure  $k_s$  of a field  $k$  is always a Galois extension.

The group  $\text{Gal}(k_s|k)$  is called the *absolute Galois group* of  $k$ .

# Galois Correspondence for Finite Extensions

## Galois Correspondence for Finite Extensions

Let  $L|k$  be a finite Galois extension with Galois group  $G$ . The maps

$$\begin{array}{ccc} \{\text{Subfields of } L \text{ containing } k\} & & \\ & \uparrow \quad \downarrow & \\ L^H \leftrightarrow H & & M \mapsto \text{Aut}(L|M) \\ & \downarrow \quad \uparrow & \\ \{\text{Subgroups of } G\} & & \end{array}$$

yields an inclusion-reversing bijection.

# Galois Correspondence for Finite Extensions

## Galois Correspondence for Finite Extensions

Let  $L|k$  be a finite Galois extension with Galois group  $G$ . The maps

$$\begin{array}{ccc} \{\text{Subfields of } L \text{ containing } k\} & & \\ & \uparrow \quad \downarrow & \\ L^H \leftrightarrow H & & M \mapsto \text{Aut}(L|M) \\ & \downarrow \quad \uparrow & \\ \{\text{Subgroups of } G\} & & \end{array}$$

yields an inclusion-reversing bijection.

The extension  $L|M$  is always Galois.

# Galois Correspondence for Finite Extensions

## Galois Correspondence for Finite Extensions

Let  $L|k$  be a finite Galois extension with Galois group  $G$ . The maps

$$\begin{array}{ccc} \{\text{Subfields of } L \text{ containing } k\} & & \\ & \uparrow \quad \downarrow & \\ L^H \leftrightarrow H & & M \mapsto \text{Aut}(L|M) \\ & \downarrow \quad \uparrow & \\ \{\text{Subgroups of } G\} & & \end{array}$$

yields an inclusion-reversing bijection.

The extension  $L|M$  is always Galois.

The extension  $M|k$  is Galois if and only if  $H := \text{Gal}(L|M)$  is a normal subgroup of  $G$ ; in this case we have  $\text{Gal}(M|k) \cong G/H$ .



## Galois Group of Infinite Galois Extensions

Let  $K|k$  be a Galois extension of fields. The Galois groups of *finite* Galois subextensions of  $K|k$  together with the homomorphisms  $\phi_{ML} : \text{Gal}(M|k) \rightarrow \text{Gal}(L|k)$  form an inverse system whose inverse limit is isomorphic to  $\text{Gal}(K|k)$ .

## Galois Group of Infinite Galois Extensions

Let  $K|k$  be a Galois extension of fields. The Galois groups of *finite* Galois subextensions of  $K|k$  together with the homomorphisms  $\phi_{ML} : \text{Gal}(M|k) \rightarrow \text{Gal}(L|k)$  form an inverse system whose inverse limit is isomorphic to  $\text{Gal}(K|k)$ .

In particular,  $\text{Gal}(K|k)$  is a profinite group.

## Galois Group of Infinite Galois Extensions

Let  $K|k$  be a Galois extension of fields. The Galois groups of *finite* Galois subextensions of  $K|k$  together with the homomorphisms  $\phi_{ML} : \text{Gal}(M|k) \rightarrow \text{Gal}(L|k)$  form an inverse system whose inverse limit is isomorphic to  $\text{Gal}(K|k)$ .

In particular,  $\text{Gal}(K|k)$  is a profinite group.

## Profinite Groups

Profinite groups are endowed with a natural topology. They are compact and totally disconnected. Moreover, the open subgroups are precisely the closed subgroups of finite index.

## Galois Correspondence for Infinite Extensions (Krull)

Let  $L$  be a subextension of the Galois extension  $K|k$ . Then  $\text{Gal}(K|L)$  is a closed subgroup of  $G = \text{Gal}(K|k)$ .

## Galois Correspondence for Infinite Extensions (Krull)

Let  $L$  be a subextension of the Galois extension  $K|k$ . Then  $\text{Gal}(K|L)$  is a closed subgroup of  $G = \text{Gal}(K|k)$ . The maps

$$\begin{array}{ccc} \{\text{Subfields of } K \text{ containing } k\} & & \\ & \begin{array}{c} \uparrow \\ K^H \leftarrow H \\ \downarrow \end{array} & \\ & & \{ \text{Closed subgroups of } G \} \end{array} \quad \begin{array}{c} \\ L \mapsto \text{Gal}(K|L) \\ \end{array}$$

yield an inclusion-reversing bijection between subfields  $K \supseteq L \supseteq k$  and closed subgroups  $H \leq G$ .

## Galois Correspondence for Infinite Extensions (Krull)

Let  $L$  be a subextension of the Galois extension  $K|k$ . Then  $\text{Gal}(K|L)$  is a closed subgroup of  $G = \text{Gal}(K|k)$ . The maps

$$\begin{array}{ccc} \{\text{Subfields of } K \text{ containing } k\} & & \\ & \begin{array}{c} \uparrow \\ K^H \leftarrow H \\ \downarrow \end{array} & \\ & & \{ \text{Closed subgroups of } G \} \end{array} \quad \begin{array}{c} \\ L \mapsto \text{Gal}(K|L) \\ \end{array}$$

yield an inclusion-reversing bijection between subfields  $K \supseteq L \supseteq k$  and closed subgroups  $H \leq G$ .

A subextension  $L|k$  is Galois over  $k$  if and only if  $H := \text{Gal}(K|L)$  is a normal subgroup of  $G$ ; in this case we have  $\text{Gal}(L|k) \cong G/H$ .

## Galois Correspondence for Infinite Extensions (Krull)

Let  $L$  be a subextension of the Galois extension  $K|k$ . Then  $\text{Gal}(K|L)$  is a closed subgroup of  $G = \text{Gal}(K|k)$ . The maps

$$\begin{array}{ccc} \{\text{Subfields of } K \text{ containing } k\} & & \\ & \uparrow \quad \downarrow & \\ K^H \leftarrow H & & L \mapsto \text{Gal}(K|L) \\ & \downarrow \quad \uparrow & \\ \{\text{Closed subgroups of } G\} & & \end{array}$$

yield an inclusion-reversing bijection between subfields  $K \supseteq L \supseteq k$  and closed subgroups  $H \leq G$ .

A subextension  $L|k$  is Galois over  $k$  if and only if  $H := \text{Gal}(K|L)$  is a normal subgroup of  $G$ ; in this case we have  $\text{Gal}(L|k) \cong G/H$ .

In particular, these all apply to  $k_s|k$  and the absolute Galois group of  $k$ .

# Grothendieck's formulation of Galois Theory

We start with a base field  $k$ , of which we fix separable and algebraic closures  $k_s \subseteq \bar{k}$ . Let  $\text{Gal}(k) := \text{Gal}(k_s|k)$ .



# Grothendieck's formulation of Galois Theory

We start with a base field  $k$ , of which we fix separable and algebraic closures  $k_s \subseteq \bar{k}$ . Let  $\text{Gal}(k) := \text{Gal}(k_s|k)$ .

Let  $L$  be a finite separable extension of  $k$ ; *here we do not consider  $L$  as a subextension of  $k_s$ .*

# Grothendieck's formulation of Galois Theory

We start with a base field  $k$ , of which we fix separable and algebraic closures  $k_s \subseteq \bar{k}$ . Let  $\text{Gal}(k) := \text{Gal}(k_s|k)$ .

Let  $L$  be a finite separable extension of  $k$ ; *here we do not consider  $L$  as a subextension of  $k_s$* .  $L$  has only finitely many  $k$ -algebra homomorphisms into  $k_s$ , the number of these is equal to  $[L : k]$ .

# Grothendieck's formulation of Galois Theory

We start with a base field  $k$ , of which we fix separable and algebraic closures  $k_s \subseteq \bar{k}$ . Let  $\text{Gal}(k) := \text{Gal}(k_s|k)$ .

Let  $L$  be a finite separable extension of  $k$ ; *here we do not consider  $L$  as a subextension of  $k_s$* .  $L$  has only finitely many  $k$ -algebra homomorphisms into  $k_s$ , the number of these is equal to  $[L : k]$ .

So we may consider the finite set  $\text{Hom}_k(L, k_s)$  which is endowed by a natural left action of  $\text{Gal}(k)$  given by  $(g, \phi) \mapsto g \circ \phi$  for  $g \in \text{Gal}(k)$ ,  $\phi \in \text{Hom}_k(L, k_s)$ .

# Grothendieck's formulation of Galois Theory

We start with a base field  $k$ , of which we fix separable and algebraic closures  $k_s \subseteq \bar{k}$ . Let  $\text{Gal}(k) := \text{Gal}(k_s|k)$ .

Let  $L$  be a finite separable extension of  $k$ ; *here we do not consider  $L$  as a subextension of  $k_s$* .  $L$  has only finitely many  $k$ -algebra homomorphisms into  $k_s$ , the number of these is equal to  $[L : k]$ .

So we may consider the finite set  $\text{Hom}_k(L, k_s)$  which is endowed by a natural left action of  $\text{Gal}(k)$  given by  $(g, \phi) \mapsto g \circ \phi$  for  $g \in \text{Gal}(k)$ ,  $\phi \in \text{Hom}_k(L, k_s)$ . This action is continuous.

# Grothendieck's formulation of Galois Theory

We start with a base field  $k$ , of which we fix separable and algebraic closures  $k_s \subseteq \bar{k}$ . Let  $\text{Gal}(k) := \text{Gal}(k_s|k)$ .

Let  $L$  be a finite separable extension of  $k$ ; *here we do not consider  $L$  as a subextension of  $k_s$* .  $L$  has only finitely many  $k$ -algebra homomorphisms into  $k_s$ , the number of these is equal to  $[L : k]$ .

So we may consider the finite set  $\text{Hom}_k(L, k_s)$  which is endowed by a natural left action of  $\text{Gal}(k)$  given by  $(g, \phi) \mapsto g \circ \phi$  for  $g \in \text{Gal}(k)$ ,  $\phi \in \text{Hom}_k(L, k_s)$ . This action is continuous.

## Main Theorem of Galois Theory, version I

Let  $k$  be a field with fixed separable closure  $k_s$ .

# Grothendieck's formulation of Galois Theory

We start with a base field  $k$ , of which we fix separable and algebraic closures  $k_s \subseteq \bar{k}$ . Let  $\text{Gal}(k) := \text{Gal}(k_s|k)$ .

Let  $L$  be a finite separable extension of  $k$ ; *here we do not consider  $L$  as a subextension of  $k_s$* .  $L$  has only finitely many  $k$ -algebra homomorphisms into  $k_s$ , the number of these is equal to  $[L : k]$ .

So we may consider the finite set  $\text{Hom}_k(L, k_s)$  which is endowed by a natural left action of  $\text{Gal}(k)$  given by  $(g, \phi) \mapsto g \circ \phi$  for  $g \in \text{Gal}(k)$ ,  $\phi \in \text{Hom}_k(L, k_s)$ . This action is continuous.

## Main Theorem of Galois Theory, version I

Let  $k$  be a field with fixed separable closure  $k_s$ . The contravariant functor

$$\text{Hom}_k(-, k_s) : \left\{ \begin{array}{c} \text{category of finite} \\ \text{separable} \\ \text{extensions of } k \end{array} \right\} \longrightarrow \left\{ \begin{array}{c} \text{category of finite sets with} \\ \text{continuous and transitive} \\ \text{left } \text{Gal}(k)\text{-action} \end{array} \right\}$$

gives an anti-equivalence.

# Grothendieck's formulation of Galois Theory

We start with a base field  $k$ , of which we fix separable and algebraic closures  $k_s \subseteq \bar{k}$ . Let  $\text{Gal}(k) := \text{Gal}(k_s|k)$ .

Let  $L$  be a finite separable extension of  $k$ ; *here we do not consider  $L$  as a subextension of  $k_s$* .  $L$  has only finitely many  $k$ -algebra homomorphisms into  $k_s$ , the number of these is equal to  $[L : k]$ .

So we may consider the finite set  $\text{Hom}_k(L, k_s)$  which is endowed by a natural left action of  $\text{Gal}(k)$  given by  $(g, \phi) \mapsto g \circ \phi$  for  $g \in \text{Gal}(k)$ ,  $\phi \in \text{Hom}_k(L, k_s)$ . This action is continuous.

## Main Theorem of Galois Theory, version I

Let  $k$  be a field with fixed separable closure  $k_s$ . The contravariant functor

$$\text{Hom}_k(-, k_s) : \left\{ \begin{array}{c} \text{category of finite} \\ \text{separable} \\ \text{extensions of } k \end{array} \right\} \longrightarrow \left\{ \begin{array}{c} \text{category of finite sets with} \\ \text{continuous and transitive} \\ \text{left } \text{Gal}(k)\text{-action} \end{array} \right\}$$

gives an anti-equivalence.

Here Galois extensions give rise to  $\text{Gal}(k)$ -sets isomorphic to some finite quotient of  $\text{Gal}(k)$ .

## Étale Algebras

A finite dimensional  $k$ -algebra  $A$  is étale (over  $k$ ) if it is isomorphic to a finite direct product of separable extensions of  $k$ .



# Grothendieck's formulation of Galois Theory

## Étale Algebras

A finite dimensional  $k$ -algebra  $A$  is étale (over  $k$ ) if it is isomorphic to a finite direct product of separable extensions of  $k$ .

## Main Theorem of Galois Theory - Grothendieck's version

Let  $k$  be a field.

# Grothendieck's formulation of Galois Theory

## Étale Algebras

A finite dimensional  $k$ -algebra  $A$  is étale (over  $k$ ) if it is isomorphic to a finite direct product of separable extensions of  $k$ .

## Main Theorem of Galois Theory - Grothendieck's version

Let  $k$  be a field. The contravariant functor

$$\mathrm{Hom}_k(-, k_s) : \left\{ \begin{array}{l} \text{category of finite} \\ \text{étale } k\text{-algebras} \end{array} \right\} \longrightarrow \left\{ \begin{array}{l} \text{category of finite sets with} \\ \text{continuous left } \mathrm{Gal}(k)\text{-action} \end{array} \right\}$$

gives an anti-equivalence.

# Grothendieck's formulation of Galois Theory

## Étale Algebras

A finite dimensional  $k$ -algebra  $A$  is étale (over  $k$ ) if it is isomorphic to a finite direct product of separable extensions of  $k$ .

## Main Theorem of Galois Theory - Grothendieck's version

Let  $k$  be a field. The contravariant functor

$$\mathrm{Hom}_k(-, k_s) : \left\{ \begin{array}{l} \text{category of finite} \\ \text{étale } k\text{-algebras} \end{array} \right\} \longrightarrow \left\{ \begin{array}{l} \text{category of finite sets with} \\ \text{continuous left } \mathrm{Gal}(k)\text{-action} \end{array} \right\}$$

gives an anti-equivalence.

Here separable field extensions give rise to sets with transitive  $\mathrm{Gal}(k)$ -action and Galois extensions to  $\mathrm{Gal}(k)$ -sets isomorphic to finite quotients of  $\mathrm{Gal}(k)$ .

# Topological Fundamental Group

× × ×

## Covers

We define the full subcategory  $\text{Cov}(X)$  of the category  $\text{Top}/X$  where  $p : Y \rightarrow X$  are subject to the condition: each point of  $X$  has an open neighbourhood  $V$  for which  $p^{-1}(V) \cong V \times I$ , where  $I$  is discrete.

## Covers

We define the full subcategory  $\text{Cov}(X)$  of the category  $\text{Top}/X$  where  $p : Y \rightarrow X$  are subject to the condition: each point of  $X$  has an open neighbourhood  $V$  for which  $p^{-1}(V) \cong V \times I$ , where  $I$  is discrete.

If  $X$  is connected, the fibres of  $p$  are all homeomorphic to the same discrete space  $I$ .

## Covers

We define the full subcategory  $\text{Cov}(X)$  of the category  $\text{Top}/X$  where  $p : Y \rightarrow X$  are subject to the condition: each point of  $X$  has an open neighbourhood  $V$  for which  $p^{-1}(V) \cong V \times I$ , where  $I$  is discrete.

If  $X$  is connected, the fibres of  $p$  are all homeomorphic to the same discrete space  $I$ .

Important examples arise from group actions on topological spaces. To obtain covers we need a technical restriction.

## Covers

We define the full subcategory  $\text{Cov}(X)$  of the category  $\text{Top}/X$  where  $p : Y \rightarrow X$  are subject to the condition: each point of  $X$  has an open neighbourhood  $V$  for which  $p^{-1}(V) \cong V \times I$ , where  $I$  is discrete.

If  $X$  is connected, the fibres of  $p$  are all homeomorphic to the same discrete space  $I$ .

Important examples arise from group actions on topological spaces. To obtain covers we need a technical restriction.

## Even (properly discontinuous) action

Let  $G$  be a group acting continuously from the left on a topological space  $Y$ . The action of  $G$  is even if each point  $y \in Y$  has some open neighbourhood  $U$  such that the open sets  $gU$  are pairwise disjoint for all  $g \in G$ .



## Covers

We define the full subcategory  $\text{Cov}(X)$  of the category  $\text{Top}/X$  where  $p : Y \rightarrow X$  are subject to the condition: each point of  $X$  has an open neighbourhood  $V$  for which  $p^{-1}(V) \cong V \times I$ , where  $I$  is discrete.

If  $X$  is connected, the fibres of  $p$  are all homeomorphic to the same discrete space  $I$ .

Important examples arise from group actions on topological spaces. To obtain covers we need a technical restriction.

## Even (properly discontinuous) action

Let  $G$  be a group acting continuously from the left on a topological space  $Y$ . The action of  $G$  is even if each point  $y \in Y$  has some open neighbourhood  $U$  such that the open sets  $gU$  are pairwise disjoint for all  $g \in G$ .

If  $G$  is a group acting evenly on a connected space  $Y$ , the projection  $p_G : Y \rightarrow G \backslash Y$  turns  $Y$  into a cover of  $G \backslash Y$ .

Henceforth we fix a base space  $X$  which will be assumed locally connected.

## Automorphism Group

Given a cover  $p : Y \rightarrow X$ , its automorphisms are to be automorphisms of  $Y$  as a space over  $X$ , i.e. topological automorphisms compatible with the projection  $p$ .

Henceforth we fix a base space  $X$  which will be assumed locally connected.

## Automorphism Group

Given a cover  $p : Y \rightarrow X$ , its automorphisms are to be automorphisms of  $Y$  as a space over  $X$ , i.e. topological automorphisms compatible with the projection  $p$ .

They form a group with respect to composition that we shall denote by  $\text{Aut}(Y|X)$ . By convention all automorphisms will be assumed to act from the left.

Henceforth we fix a base space  $X$  which will be assumed locally connected.

## Automorphism Group

Given a cover  $p : Y \rightarrow X$ , its automorphisms are to be automorphisms of  $Y$  as a space over  $X$ , i.e. topological automorphisms compatible with the projection  $p$ .

They form a group with respect to composition that we shall denote by  $\text{Aut}(Y|X)$ . By convention all automorphisms will be assumed to act from the left.

Note that for each point  $x \in X$  the group  $\text{Aut}(Y|X)$  maps the fibre  $p^{-1}(x)$  onto itself, so  $p^{-1}(x)$  is equipped with a natural action of  $\text{Aut}(Y|X)$ .

Henceforth we fix a base space  $X$  which will be assumed locally connected.

## Automorphism Group

Given a cover  $p : Y \rightarrow X$ , its automorphisms are to be automorphisms of  $Y$  as a space over  $X$ , i.e. topological automorphisms compatible with the projection  $p$ .

They form a group with respect to composition that we shall denote by  $\text{Aut}(Y|X)$ . By convention all automorphisms will be assumed to act from the left.

Note that for each point  $x \in X$  the group  $\text{Aut}(Y|X)$  maps the fibre  $p^{-1}(x)$  onto itself, so  $p^{-1}(x)$  is equipped with a natural action of  $\text{Aut}(Y|X)$ .

## Even Action

If  $p : Y \rightarrow X$  is a connected cover, the action of  $\text{Aut}(Y|X)$  on  $Y$  is even.

Henceforth we fix a base space  $X$  which will be assumed locally connected.

## Automorphism Group

Given a cover  $p : Y \rightarrow X$ , its automorphisms are to be automorphisms of  $Y$  as a space over  $X$ , i.e. topological automorphisms compatible with the projection  $p$ .

They form a group with respect to composition that we shall denote by  $\text{Aut}(Y|X)$ . By convention all automorphisms will be assumed to act from the left.

Note that for each point  $x \in X$  the group  $\text{Aut}(Y|X)$  maps the fibre  $p^{-1}(x)$  onto itself, so  $p^{-1}(x)$  is equipped with a natural action of  $\text{Aut}(Y|X)$ .

## Even Action

If  $p : Y \rightarrow X$  is a connected cover, the action of  $\text{Aut}(Y|X)$  on  $Y$  is even.

Conversely, if  $G$  is a group acting evenly on a connected space  $Y$ , the automorphism group of the cover  $p_G : Y \rightarrow G \backslash Y$  is precisely  $G$ .

## Galois Covers

Given a connected cover  $p : Y \rightarrow X$ , we may form the quotient of  $Y$  by the action of  $\text{Aut}(Y|X)$ .

## Galois Covers

Given a connected cover  $p : Y \rightarrow X$ , we may form the quotient of  $Y$  by the action of  $\text{Aut}(Y|X)$ .  $p$  factors as

$$\begin{array}{ccc} Y & \xrightarrow{\quad} & \text{Aut}(Y|X) \backslash Y \\ p \downarrow & \swarrow \bar{p} & \\ X & & \end{array}$$



## Galois Covers

Given a connected cover  $p : Y \rightarrow X$ , we may form the quotient of  $Y$  by the action of  $\text{Aut}(Y|X)$ .  $p$  factors as

$$\begin{array}{ccc} Y & \longrightarrow & \text{Aut}(Y|X) \backslash Y \\ p \downarrow & \swarrow \bar{p} & \\ X & & \end{array}$$

A connected cover  $p : Y \rightarrow X$  is said to be *Galois* if the induced map  $\bar{p}$  above is a homeomorphism.

## Galois Covers

Given a connected cover  $p : Y \rightarrow X$ , we may form the quotient of  $Y$  by the action of  $\text{Aut}(Y|X)$ .  $p$  factors as

$$\begin{array}{ccc} Y & \xrightarrow{\quad} & \text{Aut}(Y|X) \backslash Y \\ p \downarrow & \swarrow \bar{p} & \\ X & & \end{array}$$

A connected cover  $p : Y \rightarrow X$  is said to be *Galois* if the induced map  $\bar{p}$  above is a homeomorphism.

## Intermediate Covers

For a Galois cover  $p : Y \rightarrow X$ , a connected cover  $q : Z \rightarrow X$  is an intermediate cover if the following diagram commutes for some  $f : Y \rightarrow Z$

$$\begin{array}{ccc} Y & \xrightarrow{f} & Z \\ & \searrow p & \downarrow q \\ & & X \end{array}$$

## Galois Correspondence

Let  $p : Y \rightarrow X$  be a Galois cover with  $G = \text{Aut}(Y|X)$ . The maps

$$\begin{array}{ccc} & \text{\{Intermediate covers as before\}} & \\ & \uparrow \qquad \downarrow & \\ H \backslash Y \leftarrow H & & Z \mapsto \text{Aut}(Y|Z) \\ & \downarrow & \\ & \text{\{Subgroups of } G\} & \end{array}$$

yields a inclusion-preserving bijection.

## Galois Correspondence

Let  $p : Y \rightarrow X$  be a Galois cover with  $G = \text{Aut}(Y|X)$ . The maps

$$\begin{array}{ccc} & \{ \text{Intermediate covers as before} \} & \\ & \uparrow \quad \downarrow & \\ H \backslash Y \leftrightarrow H & & Z \mapsto \text{Aut}(Y|Z) \\ & \downarrow & \\ & \{ \text{Subgroups of } G \} & \end{array}$$

yields a inclusion-preserving bijection.  $f : Y \rightarrow Z$  is always Galois.

## Galois Correspondence

Let  $p : Y \rightarrow X$  be a Galois cover with  $G = \text{Aut}(Y|X)$ . The maps

$$\begin{array}{ccc} & \uparrow & \\ \{ \text{Intermediate covers as before} \} & & \\ H \backslash Y \leftarrow H & & Z \mapsto \text{Aut}(Y|Z) \\ & \downarrow & \\ & \{ \text{Subgroups of } G \} & \end{array}$$

yields a inclusion-preserving bijection.  $f : Y \rightarrow Z$  is always Galois.

The cover  $q : Z \rightarrow X$  is Galois if and only if  $H := \text{Aut}(Y|Z)$  is a normal subgroup of  $G$ ; in this case we have  $\text{Gal}(Z|X) \cong G/H$ .

## (Topological) Fundamental Group

For a topological group  $X$ , the *(topological) fundamental group of  $X$  with base point  $x$*   $\pi_1(X, x)$  is the group of homotopy classes of loops based at  $x \in X$ , where the group operation is given by concatenation of loops: for loops  $\gamma_1, \gamma_2 : [0, 1] \rightarrow X$ ,

$$(\gamma_1 \bullet \gamma_2)(x) = \begin{cases} \gamma_2(2x) & 0 \leq x \leq 1/2 \\ \gamma_1(2x - 1) & 1/2 \leq x \leq 1 \end{cases}$$

## (Topological) Fundamental Group

For a topological group  $X$ , the *(topological) fundamental group of  $X$  with base point  $x$*   $\pi_1(X, x)$  is the group of homotopy classes of loops based at  $x \in X$ , where the group operation is given by concatenation of loops: for loops  $\gamma_1, \gamma_2 : [0, 1] \rightarrow X$ ,

$$(\gamma_1 \bullet \gamma_2)(x) = \begin{cases} \gamma_2(2x) & 0 \leq x \leq 1/2 \\ \gamma_1(2x - 1) & 1/2 \leq x \leq 1 \end{cases}$$

If  $X$  is *path-connected*, i.e. any two points  $x$  and  $y$  may be joined by a path  $\rho$ , then  $\pi_1(X, x)$  is non-canonically isomorphic to  $\pi_1(X, y)$  via  $\gamma \mapsto \rho \bullet \gamma \bullet \rho^{-1}$ .

## (Topological) Fundamental Group

For a topological group  $X$ , the *(topological) fundamental group of  $X$  with base point  $x$*   $\pi_1(X, x)$  is the group of homotopy classes of loops based at  $x \in X$ , where the group operation is given by concatenation of loops: for loops  $\gamma_1, \gamma_2 : [0, 1] \rightarrow X$ ,

$$(\gamma_1 \bullet \gamma_2)(x) = \begin{cases} \gamma_2(2x) & 0 \leq x \leq 1/2 \\ \gamma_1(2x - 1) & 1/2 \leq x \leq 1 \end{cases}$$

If  $X$  is *path-connected*, i.e. any two points  $x$  and  $y$  may be joined by a path  $\rho$ , then  $\pi_1(X, x)$  is non-canonically isomorphic to  $\pi_1(X, y)$  via  $\gamma \mapsto \rho \bullet \gamma \bullet \rho^{-1}$ .

A path-connected space is always connected; it is *simply connected* if its fundamental group is trivial.



## (Topological) Fundamental Group

For a topological group  $X$ , the *(topological) fundamental group of  $X$  with base point  $x$*   $\pi_1(X, x)$  is the group of homotopy classes of loops based at  $x \in X$ , where the group operation is given by concatenation of loops: for loops  $\gamma_1, \gamma_2 : [0, 1] \rightarrow X$ ,

$$(\gamma_1 \bullet \gamma_2)(x) = \begin{cases} \gamma_2(2x) & 0 \leq x \leq 1/2 \\ \gamma_1(2x - 1) & 1/2 \leq x \leq 1 \end{cases}$$

If  $X$  is *path-connected*, i.e. any two points  $x$  and  $y$  may be joined by a path  $\rho$ , then  $\pi_1(X, x)$  is non-canonically isomorphic to  $\pi_1(X, y)$  via  $\gamma \mapsto \rho \bullet \gamma \bullet \rho^{-1}$ .

A path-connected space is always connected; it is *simply connected* if its fundamental group is trivial.

Given a cover  $p : Y \rightarrow X$ , the fibre  $p^{-1}(x)$  over a point  $x \in X$  carries a natural action by the group  $\pi_1(X, x)$ .

## Monodromy Action

Given  $y \in p^{-1}(x)$  and  $\alpha \in \pi_1(X, x)$  represented by a loop  $\gamma : [0, 1] \rightarrow X$  based at  $x$ ,

## Monodromy Action

Given  $y \in p^{-1}(x)$  and  $\alpha \in \pi_1(X, x)$  represented by a loop  $\gamma : [0, 1] \rightarrow X$  based at  $x$ , we define  $\alpha \cdot y := \tilde{\gamma}(1)$ , where  $\tilde{\gamma}$  is the unique lifting  $\tilde{\gamma}$  to  $Y$  with  $\tilde{\gamma}(0) = y$ .

## Monodromy Action

Given  $y \in p^{-1}(x)$  and  $\alpha \in \pi_1(X, x)$  represented by a loop  $\gamma : [0, 1] \rightarrow X$  based at  $x$ , we define  $\alpha \cdot y := \tilde{\gamma}(1)$ , where  $\tilde{\gamma}$  is the unique lifting  $\tilde{\gamma}$  to  $Y$  with  $\tilde{\gamma}(0) = y$ .

This defines a left action of  $\pi_1(X, x)$  on  $p^{-1}(x)$ , it is called the *monodromy action* on the fibre  $p^{-1}(x)$ .

## Monodromy Action

Given  $y \in p^{-1}(x)$  and  $\alpha \in \pi_1(X, x)$  represented by a loop  $\gamma : [0, 1] \rightarrow X$  based at  $x$ , we define  $\alpha \cdot y := \tilde{\gamma}(1)$ , where  $\tilde{\gamma}$  is the unique lifting  $\tilde{\gamma}$  to  $Y$  with  $\tilde{\gamma}(0) = y$ .

This defines a left action of  $\pi_1(X, x)$  on  $p^{-1}(x)$ , it is called the *monodromy action* on the fibre  $p^{-1}(x)$ .

## Fibre Functor

Fix a pointed space  $(X, x)$ . We define a functor

$$\text{Fib}_x : \text{Cov}(X) \longrightarrow \left\{ \begin{array}{l} \text{category of sets equipped} \\ \text{with a left } \pi_1(X, x)\text{-action} \end{array} \right\}, \quad (p : Y \rightarrow X) \longmapsto p^{-1}(x)$$

## Monodromy Action

Given  $y \in p^{-1}(x)$  and  $\alpha \in \pi_1(X, x)$  represented by a loop  $\gamma : [0, 1] \rightarrow X$  based at  $x$ , we define  $\alpha \cdot y := \tilde{\gamma}(1)$ , where  $\tilde{\gamma}$  is the unique lifting  $\tilde{\gamma}$  to  $Y$  with  $\tilde{\gamma}(0) = y$ .

This defines a left action of  $\pi_1(X, x)$  on  $p^{-1}(x)$ , it is called the *monodromy action* on the fibre  $p^{-1}(x)$ .

## Fibre Functor

Fix a pointed space  $(X, x)$ . We define a functor

$$\text{Fib}_x : \text{Cov}(X) \longrightarrow \left\{ \begin{array}{l} \text{category of sets equipped} \\ \text{with a left } \pi_1(X, x)\text{-action} \end{array} \right\}, \quad (p : Y \rightarrow X) \longmapsto p^{-1}(x)$$

If  $X$  is *connected and locally simply connected*,  $\text{Fib}_x$  induces an equivalence of categories.

## Monodromy Action

Given  $y \in p^{-1}(x)$  and  $\alpha \in \pi_1(X, x)$  represented by a loop  $\gamma : [0, 1] \rightarrow X$  based at  $x$ , we define  $\alpha \cdot y := \tilde{\gamma}(1)$ , where  $\tilde{\gamma}$  is the unique lifting  $\tilde{\gamma}$  to  $Y$  with  $\tilde{\gamma}(0) = y$ .

This defines a left action of  $\pi_1(X, x)$  on  $p^{-1}(x)$ , it is called the *monodromy action* on the fibre  $p^{-1}(x)$ .

## Fibre Functor

Fix a pointed space  $(X, x)$ . We define a functor

$$\text{Fib}_x : \text{Cov}(X) \longrightarrow \left\{ \begin{array}{l} \text{category of sets equipped} \\ \text{with a left } \pi_1(X, x)\text{-action} \end{array} \right\}, \quad (p : Y \rightarrow X) \longmapsto p^{-1}(x)$$

If  $X$  is *connected and locally simply connected*,  $\text{Fib}_x$  induces an equivalence of categories.

Connected covers correspond to  $\pi_1(X, x)$ -sets with transitive action and Galois covers to coset spaces of normal subgroups.

## Finite Covers

Call a cover  $Y \rightarrow X$  *finite* if it has finite fibres; for connected  $X$  these have the same cardinality, called the *degree* of the cover.



## Finite Covers

Call a cover  $Y \rightarrow X$  *finite* if it has finite fibres; for connected  $X$  these have the same cardinality, called the *degree* of the cover.

Let  $(X, x)$  be as before, then restricting  $\text{Fib}_x$  to the category of finite covers of  $X$  we get

$$\text{Fib}_x : \left\{ \begin{array}{c} \text{category of finite} \\ \text{covers of } X \end{array} \right\} \longrightarrow \left\{ \begin{array}{c} \text{category of sets equipped with a} \\ \text{continuous left } \pi_1(\widehat{X, x})\text{-action} \end{array} \right\}$$

is an equivalence of categories.

## Finite Covers

Call a cover  $Y \rightarrow X$  *finite* if it has finite fibres; for connected  $X$  these have the same cardinality, called the *degree* of the cover.

Let  $(X, x)$  be as before, then restricting  $\text{Fib}_x$  to the category of finite covers of  $X$  we get

$$\text{Fib}_x : \left\{ \begin{array}{c} \text{category of finite} \\ \text{covers of } X \end{array} \right\} \longrightarrow \left\{ \begin{array}{c} \text{category of sets equipped with a} \\ \text{continuous left } \pi_1(\widehat{X}, x)\text{-action} \end{array} \right\}$$

is an equivalence of categories.

Connected covers correspond to finite  $\pi_1(\widehat{X}, x)$ -sets with transitive action and Galois covers to coset spaces of open normal subgroups.

## Universal Cover

For a (connected and locally simply connected topological) pointed space  $(X, x)$  the functor  $\text{Fib}_x$  on  $\text{Cov}(X)$

## Universal Cover

For a (connected and locally simply connected topological) pointed space  $(X, x)$  the functor  $\text{Fib}_x$  on  $\text{Cov}(X)$  is representable by a cover  $\tilde{X}_x \rightarrow X$ , called the *universal cover*.

## Universal Cover

For a (connected and locally simply connected topological) pointed space  $(X, x)$  the functor  $\text{Fib}_x$  on  $\text{Cov}(X)$  is representable by a cover  $\tilde{X}_x \rightarrow X$ , called the *universal cover*.

Given an automorphism  $\phi \in \text{Aut}(\tilde{X}_x|X)$ , we have an induced bijection

$$- \circ \phi : \text{Hom}_X(\tilde{X}_x, Y) \rightarrow \text{Hom}_X(\tilde{X}_x, Y).$$

## Universal Cover

For a (connected and locally simply connected topological) pointed space  $(X, x)$  the functor  $\text{Fib}_x$  on  $\text{Cov}(X)$  is representable by a cover  $\tilde{X}_x \rightarrow X$ , called the *universal cover*.

Given an automorphism  $\phi \in \text{Aut}(\tilde{X}_x|X)$ , we have an induced bijection

$$- \circ \phi : \text{Hom}_X(\tilde{X}_x, Y) \rightarrow \text{Hom}_X(\tilde{X}_x, Y).$$

In this way we obtain a *right* action on  $\text{Hom}_X(\tilde{X}_x, Y) \cong \text{Fib}_x(Y)$  from the *left* action of  $\text{Aut}(\tilde{X}_x|X)$  on  $\tilde{X}_x$ .

## Universal Cover

For a (connected and locally simply connected topological) pointed space  $(X, x)$  the functor  $\text{Fib}_x$  on  $\text{Cov}(X)$  is representable by a cover  $\tilde{X}_x \rightarrow X$ , called the *universal cover*.

Given an automorphism  $\phi \in \text{Aut}(\tilde{X}_x|X)$ , we have an induced bijection

$$- \circ \phi : \text{Hom}_X(\tilde{X}_x, Y) \rightarrow \text{Hom}_X(\tilde{X}_x, Y).$$

In this way we obtain a *right* action on  $\text{Hom}_X(\tilde{X}_x, Y) \cong \text{Fib}_x(Y)$  from the *left* action of  $\text{Aut}(\tilde{X}_x|X)$  on  $\tilde{X}_x$ .

The cover  $\tilde{X}_x$  is a (simply) connected Galois cover of  $X$ ;

## Universal Cover

For a (connected and locally simply connected topological) pointed space  $(X, x)$  the functor  $\text{Fib}_x$  on  $\text{Cov}(X)$  is representable by a cover  $\tilde{X}_x \rightarrow X$ , called the *universal cover*.

Given an automorphism  $\phi \in \text{Aut}(\tilde{X}_x|X)$ , we have an induced bijection

$$- \circ \phi : \text{Hom}_X(\tilde{X}_x, Y) \rightarrow \text{Hom}_X(\tilde{X}_x, Y).$$

In this way we obtain a *right* action on  $\text{Hom}_X(\tilde{X}_x, Y) \cong \text{Fib}_x(Y)$  from the *left* action of  $\text{Aut}(\tilde{X}_x|X)$  on  $\tilde{X}_x$ .

The cover  $\tilde{X}_x$  is a (simply) connected Galois cover of  $X$ ; in fact, for  $X$  as above, a cover  $\tilde{X} \rightarrow X$  is universal if and only if it is simply connected.



## Universal Cover

For a (connected and locally simply connected topological) pointed space  $(X, x)$  the functor  $\text{Fib}_x$  on  $\text{Cov}(X)$  is representable by a cover  $\tilde{X}_x \rightarrow X$ , called the *universal cover*.

Given an automorphism  $\phi \in \text{Aut}(\tilde{X}_x|X)$ , we have an induced bijection

$$- \circ \phi : \text{Hom}_X(\tilde{X}_x, Y) \rightarrow \text{Hom}_X(\tilde{X}_x, Y).$$

In this way we obtain a *right* action on  $\text{Hom}_X(\tilde{X}_x, Y) \cong \text{Fib}_x(Y)$  from the *left* action of  $\text{Aut}(\tilde{X}_x|X)$  on  $\tilde{X}_x$ .

The cover  $\tilde{X}_x$  is a (simply) connected Galois cover of  $X$ ; in fact, for  $X$  as above, a cover  $\tilde{X} \rightarrow X$  is universal if and only if it is simply connected.

There is a natural isomorphism  $\text{Aut}(\tilde{X}_x|X) \cong \pi_1(X, x)$ .

## Universal Cover

For a (connected and locally simply connected topological) pointed space  $(X, x)$  the functor  $\text{Fib}_x$  on  $\text{Cov}(X)$  is representable by a cover  $\tilde{X}_x \rightarrow X$ , called the *universal cover*.

Given an automorphism  $\phi \in \text{Aut}(\tilde{X}_x|X)$ , we have an induced bijection

$$- \circ \phi : \text{Hom}_X(\tilde{X}_x, Y) \rightarrow \text{Hom}_X(\tilde{X}_x, Y).$$

In this way we obtain a *right* action on  $\text{Hom}_X(\tilde{X}_x, Y) \cong \text{Fib}_x(Y)$  from the *left* action of  $\text{Aut}(\tilde{X}_x|X)$  on  $\tilde{X}_x$ .

The cover  $\tilde{X}_x$  is a (simply) connected Galois cover of  $X$ ; in fact, for  $X$  as above, a cover  $\tilde{X} \rightarrow X$  is universal if and only if it is simply connected.

There is a natural isomorphism  $\text{Aut}(\tilde{X}_x|X) \cong \pi_1(X, x)$ . Moreover, for each cover  $Y \rightarrow X$  the left action of  $\text{Aut}(\tilde{X}_x|X)^{\text{op}}$  on  $\text{Fib}_x(Y)$  described above is exactly the monodromy action of  $\pi_1(X, x)$ .

## Universal Cover

For a (connected and locally simply connected topological) pointed space  $(X, x)$  the functor  $\text{Fib}_x$  on  $\text{Cov}(X)$  is representable by a cover  $\tilde{X}_x \rightarrow X$ , called the *universal cover*.

Given an automorphism  $\phi \in \text{Aut}(\tilde{X}_x|X)$ , we have an induced bijection

$$- \circ \phi : \text{Hom}_X(\tilde{X}_x, Y) \rightarrow \text{Hom}_X(\tilde{X}_x, Y).$$

In this way we obtain a *right* action on  $\text{Hom}_X(\tilde{X}_x, Y) \cong \text{Fib}_x(Y)$  from the *left* action of  $\text{Aut}(\tilde{X}_x|X)$  on  $\tilde{X}_x$ .

The cover  $\tilde{X}_x$  is a (simply) connected Galois cover of  $X$ ; in fact, for  $X$  as above, a cover  $\tilde{X} \rightarrow X$  is universal if and only if it is simply connected.

There is a natural isomorphism  $\text{Aut}(\tilde{X}_x|X) \cong \pi_1(X, x)$ . Moreover, for each cover  $Y \rightarrow X$  the left action of  $\text{Aut}(\tilde{X}_x|X)^{\text{op}}$  on  $\text{Fib}_x(Y)$  described above is exactly the monodromy action of  $\pi_1(X, x)$ .

Let  $X$  as above, let  $x$  and  $y$  be two base points and consider the universal covers  $\tilde{X}_x$  and  $\tilde{X}_y$  that represent  $\text{Fib}_x$  and  $\text{Fib}_y$ .

## Universal Cover

For a (connected and locally simply connected topological) pointed space  $(X, x)$  the functor  $\text{Fib}_x$  on  $\text{Cov}(X)$  is representable by a cover  $\tilde{X}_x \rightarrow X$ , called the *universal cover*.

Given an automorphism  $\phi \in \text{Aut}(\tilde{X}_x|X)$ , we have an induced bijection

$$- \circ \phi : \text{Hom}_X(\tilde{X}_x, Y) \rightarrow \text{Hom}_X(\tilde{X}_x, Y).$$

In this way we obtain a *right* action on  $\text{Hom}_X(\tilde{X}_x, Y) \cong \text{Fib}_x(Y)$  from the *left* action of  $\text{Aut}(\tilde{X}_x|X)$  on  $\tilde{X}_x$ .

The cover  $\tilde{X}_x$  is a (simply) connected Galois cover of  $X$ ; in fact, for  $X$  as above, a cover  $\tilde{X} \rightarrow X$  is universal if and only if it is simply connected.

There is a natural isomorphism  $\text{Aut}(\tilde{X}_x|X) \cong \pi_1(X, x)$ . Moreover, for each cover  $Y \rightarrow X$  the left action of  $\text{Aut}(\tilde{X}_x|X)^{\text{op}}$  on  $\text{Fib}_x(Y)$  described above is exactly the monodromy action of  $\pi_1(X, x)$ .

Let  $X$  as above, let  $x$  and  $y$  be two base points and consider the universal covers  $\tilde{X}_x$  and  $\tilde{X}_y$  that represent  $\text{Fib}_x$  and  $\text{Fib}_y$ . Then there's a bijection between homotopy classes of paths joining  $y$  to  $x$ , sometimes denoted as  $\pi_1(X; y, x)$ , and  $\text{Isom}_X(\tilde{X}_x, \tilde{X}_y)$ .

## Some Translations

For  $X$  and  $x$  (and  $y$ ) as before, our discussion tells us that we could have given an alternative definition of  $\pi_1(X, x)$  and  $\pi_1(X; y, x)$ .

## Some Translations

For  $X$  and  $x$  (and  $y$ ) as before, our discussion tells us that we could have given an alternative definition of  $\pi_1(\tilde{X}, x)$  and  $\pi_1(X; y, x)$ . Because, by Yoneda we obtain

$$\pi_1(X, x) \cong \text{Aut}(\tilde{X}_x | X)$$

## Some Translations

For  $X$  and  $x$  (and  $y$ ) as before, our discussion tells us that we could have given an alternative definition of  $\pi_1(\tilde{X}, x)$  and  $\pi_1(X; y, x)$ . Because, by Yoneda we obtain

$$\pi_1(X, x) \cong \text{Aut}(\tilde{X}_x | X) \cong \text{Aut}(\text{Hom}_X(\tilde{X}_x, -)) \cong \text{Aut}(\text{Fib}_x)$$

## Some Translations

For  $X$  and  $x$  (and  $y$ ) as before, our discussion tells us that we could have given an alternative definition of  $\pi_1(\tilde{X}, x)$  and  $\pi_1(X; y, x)$ . Because, by Yoneda we obtain

$$\pi_1(X, x) \cong \text{Aut}(\tilde{X}_x | X) \cong \text{Aut}(\text{Hom}_X(\tilde{X}_x, -)) \cong \text{Aut}(\text{Fib}_x)$$

$$\pi_1(X; y, x) \cong \text{Isom}_X(\tilde{X}_x, \tilde{X}_y)$$



## Some Translations

For  $X$  and  $x$  (and  $y$ ) as before, our discussion tells us that we could have given an alternative definition of  $\pi_1(\tilde{X}, x)$  and  $\pi_1(X; y, x)$ . Because, by Yoneda we obtain

$$\pi_1(X, x) \cong \text{Aut}(\tilde{X}_x | X) \cong \text{Aut}(\text{Hom}_X(\tilde{X}_x, -)) \cong \text{Aut}(\text{Fib}_x)$$

$$\pi_1(X; y, x) \cong \text{Isom}_X(\tilde{X}_x, \tilde{X}_y) \cong \text{Isom}(\text{Hom}_X(\tilde{X}_y, -), \text{Hom}_X(\tilde{X}_x, -)) \cong \text{Isom}(\text{Fib}_y, \text{Fib}_x)$$

## Some Translations

For  $X$  and  $x$  (and  $y$ ) as before, our discussion tells us that we could have given an alternative definition of  $\pi_1(X, x)$  and  $\pi_1(X; y, x)$ . Because, by Yoneda we obtain

$$\pi_1(X, x) \cong \text{Aut}(\tilde{X}_x | X) \cong \text{Aut}(\text{Hom}_X(\tilde{X}_x, -)) \cong \text{Aut}(\text{Fib}_x)$$

$$\pi_1(X; y, x) \cong \text{Isom}_X(\tilde{X}_x, \tilde{X}_y) \cong \text{Isom}(\text{Hom}_X(\tilde{X}_y, -), \text{Hom}_X(\tilde{X}_x, -)) \cong \text{Isom}(\text{Fib}_y, \text{Fib}_x)$$

As before, an element  $\rho \in \text{Isom}(\text{Fib}_y, \text{Fib}_x)$  induces an isomorphism

$$\text{Aut}(\text{Fib}_x) \rightarrow \text{Aut}(\text{Fib}_y) : \gamma \mapsto \rho^{-1} \circ \gamma \circ \rho.$$

## Some Translations

For  $X$  and  $x$  (and  $y$ ) as before, our discussion tells us that we could have given an alternative definition of  $\pi_1(X, x)$  and  $\pi_1(X; y, x)$ . Because, by Yoneda we obtain

$$\pi_1(X, x) \cong \text{Aut}(\tilde{X}_x | X) \cong \text{Aut}(\text{Hom}_X(\tilde{X}_x, -)) \cong \text{Aut}(\text{Fib}_x)$$

$$\pi_1(X; y, x) \cong \text{Isom}_X(\tilde{X}_x, \tilde{X}_y) \cong \text{Isom}(\text{Hom}_X(\tilde{X}_y, -), \text{Hom}_X(\tilde{X}_x, -)) \cong \text{Isom}(\text{Fib}_y, \text{Fib}_x)$$

As before, an element  $\rho \in \text{Isom}(\text{Fib}_y, \text{Fib}_x)$  induces an isomorphism

$$\text{Aut}(\text{Fib}_x) \rightarrow \text{Aut}(\text{Fib}_y) : \gamma \mapsto \rho^{-1} \circ \gamma \circ \rho.$$

Furthermore, the monodromy action is translated as follows: for any automorphism  $\phi \in \text{Aut}(\text{Fib}_x)$ , and a cover  $Y \rightarrow X$ , there's by definition a morphism  $\text{Fib}_x(Y) \rightarrow \text{Fib}_x(Y)$  induced by  $\phi$  which then gives a natural left action of  $\text{Aut}(\text{Fib}_x)$  on  $\text{Fib}_x(Y)$ .

# Étale Fundamental Group

× × ×

## Schemes

A scheme is a locally ringed space  $(X, \mathcal{O}_X)$  having an open covering  $\{U_i\}_{i \in I}$  such that for all  $i$  the locally ringed spaces  $(U_i, \mathcal{O}_X|_{U_i})$  are isomorphic to affine schemes  $(\operatorname{Spec} A_i, \mathcal{O}_{\operatorname{Spec} A_i})$ .

## Schemes

A scheme is a locally ringed space  $(X, \mathcal{O}_X)$  having an open covering  $\{U_i\}_{i \in I}$  such that for all  $i$  the locally ringed spaces  $(U_i, \mathcal{O}_X|_{U_i})$  are isomorphic to affine schemes  $(\operatorname{Spec} A_i, \mathcal{O}_{\operatorname{Spec} A_i})$ . In particular, for any  $p \in X$ , the stalk  $\mathcal{O}_{X,p}$  is a local ring with unique maximal ideal  $\mathfrak{m}_p$ . We call  $\kappa(p) := \mathcal{O}_{X,p}/\mathfrak{m}_p$  the *residue field* at  $p$ .

## Schemes

A scheme is a locally ringed space  $(X, \mathcal{O}_X)$  having an open covering  $\{U_i\}_{i \in I}$  such that for all  $i$  the locally ringed spaces  $(U_i, \mathcal{O}_X|_{U_i})$  are isomorphic to affine schemes  $(\operatorname{Spec} A_i, \mathcal{O}_{\operatorname{Spec} A_i})$ . In particular, for any  $p \in X$ , the stalk  $\mathcal{O}_{X,p}$  is a local ring with unique maximal ideal  $\mathfrak{m}_p$ . We call  $\kappa(p) := \mathcal{O}_{X,p} / \mathfrak{m}_p$  the *residue field* at  $p$ .

One can then identify  $p$  with  $\operatorname{Spec} \kappa(p)$ , i.e., the canonical morphism  $\operatorname{Spec} \kappa(p) \rightarrow X$  is an isomorphism onto  $p$ .

## Schemes

A scheme is a locally ringed space  $(X, \mathcal{O}_X)$  having an open covering  $\{U_i\}_{i \in I}$  such that for all  $i$  the locally ringed spaces  $(U_i, \mathcal{O}_X|_{U_i})$  are isomorphic to affine schemes  $(\operatorname{Spec} A_i, \mathcal{O}_{\operatorname{Spec} A_i})$ . In particular, for any  $p \in X$ , the stalk  $\mathcal{O}_{X,p}$  is a local ring with unique maximal ideal  $\mathfrak{m}_p$ . We call  $\kappa(p) := \mathcal{O}_{X,p}/\mathfrak{m}_p$  the *residue field* at  $p$ .

One can then identify  $p$  with  $\operatorname{Spec} \kappa(p)$ , i.e., the canonical morphism  $\operatorname{Spec} \kappa(p) \rightarrow X$  is an isomorphism onto  $p$ . In the affine case, this translates to saying for a prime  $\mathfrak{p}$  of a ring  $A$ , the canonical morphism  $\operatorname{Spec} A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} \rightarrow \operatorname{Spec} A$  has image  $\mathfrak{p}$ .



## Schemes

A scheme is a locally ringed space  $(X, \mathcal{O}_X)$  having an open covering  $\{U_i\}_{i \in I}$  such that for all  $i$  the locally ringed spaces  $(U_i, \mathcal{O}_X|_{U_i})$  are isomorphic to affine schemes  $(\text{Spec } A_i, \mathcal{O}_{\text{Spec } A_i})$ . In particular, for any  $p \in X$ , the stalk  $\mathcal{O}_{X,p}$  is a local ring with unique maximal ideal  $\mathfrak{m}_p$ . We call  $\kappa(p) := \mathcal{O}_{X,p}/\mathfrak{m}_p$  the *residue field* at  $p$ .

One can then identify  $p$  with  $\text{Spec } \kappa(p)$ , i.e., the canonical morphism  $\text{Spec } \kappa(p) \rightarrow X$  is an isomorphism onto  $p$ . In the affine case, this translates to saying for a prime  $\mathfrak{p}$  of a ring  $A$ , the canonical morphism  $\text{Spec } A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} \rightarrow \text{Spec } A$  has image  $\mathfrak{p}$ .

## Fibres

Given a morphism  $\phi : X \rightarrow S$  and a point  $p \in S$ , the fibre of  $\phi$  at  $p$  is the scheme  $X_p := X \times_S \text{Spec } \kappa(p)$ .

## Schemes

A scheme is a locally ringed space  $(X, \mathcal{O}_X)$  having an open covering  $\{U_i\}_{i \in I}$  such that for all  $i$  the locally ringed spaces  $(U_i, \mathcal{O}_X|_{U_i})$  are isomorphic to affine schemes  $(\text{Spec } A_i, \mathcal{O}_{\text{Spec } A_i})$ . In particular, for any  $p \in X$ , the stalk  $\mathcal{O}_{X,p}$  is a local ring with unique maximal ideal  $\mathfrak{m}_p$ . We call  $\kappa(p) := \mathcal{O}_{X,p}/\mathfrak{m}_p$  the *residue field* at  $p$ .

One can then identify  $p$  with  $\text{Spec } \kappa(p)$ , i.e., the canonical morphism  $\text{Spec } \kappa(p) \rightarrow X$  is an isomorphism onto  $p$ . In the affine case, this translates to saying for a prime  $\mathfrak{p}$  of a ring  $A$ , the canonical morphism  $\text{Spec } A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} \rightarrow \text{Spec } A$  has image  $\mathfrak{p}$ .

## Fibres

Given a morphism  $\phi : X \rightarrow S$  and a point  $p \in S$ , the fibre of  $\phi$  at  $p$  is the scheme  $X_p := X \times_S \text{Spec } \kappa(p)$ .

Recall that the underlying topological space of a fibre product of schemes is not the topological fibre product of the underlying spaces in general.

## Schemes

A scheme is a locally ringed space  $(X, \mathcal{O}_X)$  having an open covering  $\{U_i\}_{i \in I}$  such that for all  $i$  the locally ringed spaces  $(U_i, \mathcal{O}_X|_{U_i})$  are isomorphic to affine schemes  $(\text{Spec } A_i, \mathcal{O}_{\text{Spec } A_i})$ . In particular, for any  $p \in X$ , the stalk  $\mathcal{O}_{X,p}$  is a local ring with unique maximal ideal  $\mathfrak{m}_p$ . We call  $\kappa(p) := \mathcal{O}_{X,p}/\mathfrak{m}_p$  the *residue field* at  $p$ .

One can then identify  $p$  with  $\text{Spec } \kappa(p)$ , i.e., the canonical morphism  $\text{Spec } \kappa(p) \rightarrow X$  is an isomorphism onto  $p$ . In the affine case, this translates to saying for a prime  $\mathfrak{p}$  of a ring  $A$ , the canonical morphism  $\text{Spec } A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} \rightarrow \text{Spec } A$  has image  $\mathfrak{p}$ .

## Fibres

Given a morphism  $\phi : X \rightarrow S$  and a point  $p \in S$ , the fibre of  $\phi$  at  $p$  is the scheme  $X_p := X \times_S \text{Spec } \kappa(p)$ .

Recall that the underlying topological space of a fibre product of schemes is not the topological fibre product of the underlying spaces in general.

In the affine case, for a morphism  $\phi : \text{Spec } B \rightarrow \text{Spec } A$  and a prime  $\mathfrak{p} \in A$ , the fibre of  $\phi$  at  $\mathfrak{p}$  is  $\text{Spec } B \times_{\text{Spec } A} \text{Spec } A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} = \text{Spec}(B \otimes_A A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}) \cong \text{Spec } B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$ .

## Schemes

A scheme is a locally ringed space  $(X, \mathcal{O}_X)$  having an open covering  $\{U_i\}_{i \in I}$  such that for all  $i$  the locally ringed spaces  $(U_i, \mathcal{O}_X|_{U_i})$  are isomorphic to affine schemes  $(\text{Spec } A_i, \mathcal{O}_{\text{Spec } A_i})$ . In particular, for any  $p \in X$ , the stalk  $\mathcal{O}_{X,p}$  is a local ring with unique maximal ideal  $\mathfrak{m}_p$ . We call  $\kappa(p) := \mathcal{O}_{X,p}/\mathfrak{m}_p$  the *residue field* at  $p$ .

One can then identify  $p$  with  $\text{Spec } \kappa(p)$ , i.e., the canonical morphism  $\text{Spec } \kappa(p) \rightarrow X$  is an isomorphism onto  $p$ . In the affine case, this translates to saying for a prime  $\mathfrak{p}$  of a ring  $A$ , the canonical morphism  $\text{Spec } A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} \rightarrow \text{Spec } A$  has image  $\mathfrak{p}$ .

## Fibres

Given a morphism  $\phi : X \rightarrow S$  and a point  $p \in S$ , the fibre of  $\phi$  at  $p$  is the scheme  $X_p := X \times_S \text{Spec } \kappa(p)$ .

Recall that the underlying topological space of a fibre product of schemes is not the topological fibre product of the underlying spaces in general.

In the affine case, for a morphism  $\phi : \text{Spec } B \rightarrow \text{Spec } A$  and a prime  $\mathfrak{p} \in A$ , the fibre of  $\phi$  at  $\mathfrak{p}$  is  $\text{Spec } B \times_{\text{Spec } A} \text{Spec } A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} = \text{Spec}(B \otimes_A A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}) \cong \text{Spec } B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$ .

The underlying topological space of the fibre  $X_p$  is homeomorphic to the subspace  $\phi^{-1}(p)$  of the underlying space of  $X$ .

## Finite Étale Covers

A morphism  $\phi : X \rightarrow S$  is *flat* if, for every  $x \in X$ , the induced map of stalks  $\mathcal{O}_{S,\phi(x)} \rightarrow \mathcal{O}_{X,x}$  makes  $\mathcal{O}_{X,x}$  a flat  $\mathcal{O}_{S,\phi(x)}$ -module.

## Finite Étale Covers

A morphism  $\phi : X \rightarrow S$  is *flat* if, for every  $x \in X$ , the induced map of stalks  $\mathcal{O}_{S, \phi(x)} \rightarrow \mathcal{O}_{X, x}$  makes  $\mathcal{O}_{X, x}$  a flat  $\mathcal{O}_{S, \phi(x)}$ -module.

A morphism  $\phi : X \rightarrow S$  is called *finite* if it's affine, i.e. for any affine open  $\text{Spec } A$  of  $S$ ,  $\phi^{-1}(\text{Spec } A)$  is an affine open of  $X$ , say  $\text{Spec } B$ , and the corresponding map of rings  $\phi^\# : A \rightarrow B$  makes  $B$  a finitely generated module over  $A$ .

## Finite Étale Covers

A morphism  $\phi : X \rightarrow S$  is *flat* if, for every  $x \in X$ , the induced map of stalks  $\mathcal{O}_{S, \phi(x)} \rightarrow \mathcal{O}_{X, x}$  makes  $\mathcal{O}_{X, x}$  a flat  $\mathcal{O}_{S, \phi(x)}$ -module.

A morphism  $\phi : X \rightarrow S$  is called *finite* if it's affine, i.e. for any affine open  $\text{Spec } A$  of  $S$ ,  $\phi^{-1}(\text{Spec } A)$  is an affine open of  $X$ , say  $\text{Spec } B$ , and the corresponding map of rings  $\phi^\# : A \rightarrow B$  makes  $B$  a finitely generated module over  $A$ .

A finite and flat morphism  $\phi : X \rightarrow S$  is called a *finite étale morphism* if for any  $p \in S$ , the fibre  $X_p \cong \text{Spec } A$ , where  $A$  is a finite étale  $\kappa(p)$ -algebra.

## Finite Étale Covers

A morphism  $\phi : X \rightarrow S$  is *flat* if, for every  $x \in X$ , the induced map of stalks  $\mathcal{O}_{S, \phi(x)} \rightarrow \mathcal{O}_{X, x}$  makes  $\mathcal{O}_{X, x}$  a flat  $\mathcal{O}_{S, \phi(x)}$ -module.

A morphism  $\phi : X \rightarrow S$  is called *finite* if it's affine, i.e. for any affine open  $\text{Spec } A$  of  $S$ ,  $\phi^{-1}(\text{Spec } A)$  is an affine open of  $X$ , say  $\text{Spec } B$ , and the corresponding map of rings  $\phi^\# : A \rightarrow B$  makes  $B$  a finitely generated module over  $A$ .

A finite and flat morphism  $\phi : X \rightarrow S$  is called a *finite étale morphism* if for any  $p \in S$ , the fibre  $X_p \cong \text{Spec } A$ , where  $A$  is a finite étale  $\kappa(p)$ -algebra.

A surjective finite étale morphism is called a *finite étale cover*.



## Geometric Fibres

A *geometric point* of a scheme  $S$  is a morphism  $\bar{s} : \mathrm{Spec}(\Omega) \rightarrow S$ , where  $\Omega$  is an algebraically closed field.

## Geometric Fibres

A *geometric point* of a scheme  $S$  is a morphism  $\bar{s} : \operatorname{Spec}(\Omega) \rightarrow S$ , where  $\Omega$  is an algebraically closed field. The image of  $\bar{s}$  is a point  $s$  of  $S$  such that  $\Omega$  is an algebraic closure of  $\kappa(s)$ .

## Geometric Fibres

A *geometric point* of a scheme  $S$  is a morphism  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$ , where  $\Omega$  is an algebraically closed field. The image of  $\bar{s}$  is a point  $s$  of  $S$  such that  $\Omega$  is an algebraic closure of  $\kappa(s)$ .  $(S, \bar{s})$  is called a pointed scheme.

## Geometric Fibres

A *geometric point* of a scheme  $S$  is a morphism  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$ , where  $\Omega$  is an algebraically closed field. The image of  $\bar{s}$  is a point  $s$  of  $S$  such that  $\Omega$  is an algebraic closure of  $\kappa(s)$ .  $(S, \bar{s})$  is called a *pointed scheme*.

Given a morphism  $\phi : X \rightarrow S$  and a geometric point  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$  of  $S$ ,

## Geometric Fibres

A *geometric point* of a scheme  $S$  is a morphism  $\bar{s} : \operatorname{Spec}(\Omega) \rightarrow S$ , where  $\Omega$  is an algebraically closed field. The image of  $\bar{s}$  is a point  $s$  of  $S$  such that  $\Omega$  is an algebraic closure of  $\kappa(s)$ .  $(S, \bar{s})$  is called a *pointed scheme*.

Given a morphism  $\phi : X \rightarrow S$  and a geometric point  $\bar{s} : \operatorname{Spec}(\Omega) \rightarrow S$  of  $S$ , the *geometric fibre* of  $\phi$  over  $\bar{s}$  is defined to be  $X_{\bar{s}} := X \times_S \operatorname{Spec}(\Omega)$ .

## Geometric Fibres

A *geometric point* of a scheme  $S$  is a morphism  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$ , where  $\Omega$  is an algebraically closed field. The image of  $\bar{s}$  is a point  $s$  of  $S$  such that  $\Omega$  is an algebraic closure of  $\kappa(s)$ .  $(S, \bar{s})$  is called a *pointed scheme*.

Given a morphism  $\phi : X \rightarrow S$  and a geometric point  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$  of  $S$ , the *geometric fibre* of  $\phi$  over  $\bar{s}$  is defined to be  $X_{\bar{s}} := X \times_S \text{Spec}(\Omega)$ .

The fibres of  $\phi$  are spectra of finite étale algebras if and only if its geometric fibres are of the form  $\text{Spec}(\Omega \times \cdots \times \Omega)$ , i.e. they are finite disjoint unions of points defined over  $\Omega$ .

## Fibre Functor

For a scheme  $S$  denote by  $\mathbf{F\acute{E}t}_S$  the full subcategory of  $\mathbf{Sch}/S$  whose objects are finite étale covers of  $S$ .

## Fibre Functor

For a scheme  $S$  denote by  $\mathbf{F\acute{E}t}_S$  the full subcategory of  $\mathbf{Sch}/S$  whose objects are finite étale covers of  $S$ .

Fix a geometric point  $\bar{s} : \mathrm{Spec}(\Omega) \rightarrow S$ . For an object  $X \rightarrow S$  of  $\mathbf{F\acute{E}t}_S$  denote by  $\mathrm{Fib}_{\bar{s}}(X)$  the underlying set of the geometric fibre  $X_{\bar{s}}$ .



## Fibre Functor

For a scheme  $S$  denote by  $\mathbf{F\acute{E}t}_S$  the full subcategory of  $\mathbf{Sch}/S$  whose objects are finite étale covers of  $S$ .

Fix a geometric point  $\bar{s} : \mathrm{Spec}(\Omega) \rightarrow S$ . For an object  $X \rightarrow S$  of  $\mathbf{F\acute{E}t}_S$  denote by  $\mathrm{Fib}_{\bar{s}}(X)$  the underlying set of the geometric fibre  $X_{\bar{s}}$ .

Given a morphism  $X \rightarrow Y$  in  $\mathbf{F\acute{E}t}_S$ , there is an induced morphism of schemes  $X_{\bar{s}} \rightarrow Y_{\bar{s}}$ , whence a set-theoretic map  $\mathrm{Fib}_{\bar{s}}(X) \rightarrow \mathrm{Fib}_{\bar{s}}(Y)$ .

## Fibre Functor

For a scheme  $S$  denote by  $\mathbf{F\acute{E}t}_S$  the full subcategory of  $\mathbf{Sch}/S$  whose objects are finite étale covers of  $S$ .

Fix a geometric point  $\bar{s} : \mathrm{Spec}(\Omega) \rightarrow S$ . For an object  $X \rightarrow S$  of  $\mathbf{F\acute{E}t}_S$  denote by  $\mathrm{Fib}_{\bar{s}}(X)$  the underlying set of the geometric fibre  $X_{\bar{s}}$ .

Given a morphism  $X \rightarrow Y$  in  $\mathbf{F\acute{E}t}_S$ , there is an induced morphism of schemes  $X_{\bar{s}} \rightarrow Y_{\bar{s}}$ , whence a set-theoretic map  $\mathrm{Fib}_{\bar{s}}(X) \rightarrow \mathrm{Fib}_{\bar{s}}(Y)$ .

We have thus defined a set-valued functor

$$\mathrm{Fib}_{\bar{s}} : \mathbf{F\acute{E}t}_S \longrightarrow \mathbf{FinSet};$$

we call it the *fibre functor* at the geometric point  $\bar{s}$ .

# Étale Fundamental Group

## Fibre Functor

For a scheme  $S$  denote by  $\mathbf{F\acute{E}t}_S$  the full subcategory of  $\mathbf{Sch}/S$  whose objects are finite étale covers of  $S$ .

Fix a geometric point  $\bar{s} : \mathrm{Spec}(\Omega) \rightarrow S$ . For an object  $X \rightarrow S$  of  $\mathbf{F\acute{E}t}_S$  denote by  $\mathrm{Fib}_{\bar{s}}(X)$  the underlying set of the geometric fibre  $X_{\bar{s}}$ .

Given a morphism  $X \rightarrow Y$  in  $\mathbf{F\acute{E}t}_S$ , there is an induced morphism of schemes  $X_{\bar{s}} \rightarrow Y_{\bar{s}}$ , whence a set-theoretic map  $\mathrm{Fib}_{\bar{s}}(X) \rightarrow \mathrm{Fib}_{\bar{s}}(Y)$ .

We have thus defined a set-valued functor

$$\mathrm{Fib}_{\bar{s}} : \mathbf{F\acute{E}t}_S \longrightarrow \mathbf{FinSet};$$

we call it the *fibre functor* at the geometric point  $\bar{s}$ .

## Étale Fundamental Group

Given a pointed scheme  $(S, \bar{s})$ , we define the *étale fundamental group*  $\pi_1(S, \bar{s}) := \mathrm{Aut}(\mathrm{Fib}_{\bar{s}})$ .

## Fibre Functor Revisited

Note that there's a natural left action of  $\text{Aut}(\text{Fib}_{\bar{s}})$  on  $\text{Fib}_{\bar{s}}(X)$  for each finite étale cover  $X$  of  $S$ .

## Fibre Functor Revisited

Note that there's a natural left action of  $\text{Aut}(\text{Fib}_{\bar{s}})$  on  $\text{Fib}_{\bar{s}}(X)$  for each finite étale cover  $X$  of  $S$ . Therefore, we have

$$\text{Fib}_{\bar{s}} : \text{FÉt}_S \longrightarrow \left\{ \begin{array}{l} \text{category of finite} \\ \text{left } \pi_1(S, \bar{s})\text{-sets} \end{array} \right\}$$

## Fibre Functor Revisited

Note that there's a natural left action of  $\text{Aut}(\text{Fib}_{\bar{s}})$  on  $\text{Fib}_{\bar{s}}(X)$  for each finite étale cover  $X$  of  $S$ . Therefore, we have

$$\text{Fib}_{\bar{s}} : \text{FÉt}_S \longrightarrow \left\{ \begin{array}{l} \text{category of finite} \\ \text{left } \pi_1(S, \bar{s})\text{-sets} \end{array} \right\}$$

## Main Theorem (Grothendieck)

Let  $(S, \bar{s})$  be a connected pointed scheme.

## Fibre Functor Revisited

Note that there's a natural left action of  $\text{Aut}(\text{Fib}_{\bar{s}})$  on  $\text{Fib}_{\bar{s}}(X)$  for each finite étale cover  $X$  of  $S$ . Therefore, we have

$$\text{Fib}_{\bar{s}} : \text{FÉt}_S \longrightarrow \left\{ \begin{array}{l} \text{category of finite} \\ \text{left } \pi_1(S, \bar{s})\text{-sets} \end{array} \right\}$$

## Main Theorem (Grothendieck)

Let  $(S, \bar{s})$  be a connected pointed scheme.

- (1) The group  $\pi_1(S, \bar{s})$  is profinite, and its action on  $\text{Fib}_{\bar{s}}(X)$  is continuous for every  $X$  in  $\text{FÉt}_S$ .

## Fibre Functor Revisited

Note that there's a natural left action of  $\text{Aut}(\text{Fib}_{\bar{s}})$  on  $\text{Fib}_{\bar{s}}(X)$  for each finite étale cover  $X$  of  $S$ . Therefore, we have

$$\text{Fib}_{\bar{s}} : \mathbf{F\acute{E}t}_S \longrightarrow \left\{ \begin{array}{l} \text{category of finite} \\ \text{left } \pi_1(S, \bar{s})\text{-sets} \end{array} \right\}$$

## Main Theorem (Grothendieck)

Let  $(S, \bar{s})$  be a connected pointed scheme.

- (1) The group  $\pi_1(S, \bar{s})$  is profinite, and its action on  $\text{Fib}_{\bar{s}}(X)$  is continuous for every  $X$  in  $\mathbf{F\acute{E}t}_S$ .
- (2) The functor

$$\text{Fib}_{\bar{s}} : \mathbf{F\acute{E}t}_S \longrightarrow \left\{ \begin{array}{l} \text{category of finite} \\ \text{continuous left} \\ \pi_1(S, \bar{s})\text{-sets} \end{array} \right\}$$

induces an equivalence of categories.



## Fibre Functor Revisited

Note that there's a natural left action of  $\text{Aut}(\text{Fib}_{\bar{s}})$  on  $\text{Fib}_{\bar{s}}(X)$  for each finite étale cover  $X$  of  $S$ . Therefore, we have

$$\text{Fib}_{\bar{s}} : \mathbf{F\acute{E}t}_S \longrightarrow \left\{ \begin{array}{l} \text{category of finite} \\ \text{left } \pi_1(S, \bar{s})\text{-sets} \end{array} \right\}$$

## Main Theorem (Grothendieck)

Let  $(S, \bar{s})$  be a connected pointed scheme.

- (1) The group  $\pi_1(S, \bar{s})$  is profinite, and its action on  $\text{Fib}_{\bar{s}}(X)$  is continuous for every  $X$  in  $\mathbf{F\acute{E}t}_S$ .
- (2) The functor

$$\text{Fib}_{\bar{s}} : \mathbf{F\acute{E}t}_S \longrightarrow \left\{ \begin{array}{l} \text{category of finite} \\ \text{continuous left} \\ \pi_1(S, \bar{s})\text{-sets} \end{array} \right\}$$

induces an equivalence of categories. Here connected covers correspond to sets with transitive  $\pi_1(S, \bar{s})$ -action, and Galois covers to finite quotients of  $\pi_1(S, \bar{s})$ .

## Revisiting Galois Theory

Let  $S = \operatorname{Spec} k$  for a field  $k$ . Here a finite étale  $S$ -scheme  $X$  is the spectrum of a finite étale  $k$ -algebra.

## Revisiting Galois Theory

Let  $S = \operatorname{Spec} k$  for a field  $k$ . Here a finite étale  $S$ -scheme  $X$  is the spectrum of a finite étale  $k$ -algebra.

For a geometric point  $\bar{s}$  the fibre functor maps a connected cover  $X = \operatorname{Spec} L$  to the underlying set of  $\operatorname{Spec}(L \otimes_k \Omega)$ , which is a finite set indexed by the  $k$ -algebra homomorphisms  $L \rightarrow \Omega$ .

## Revisiting Galois Theory

Let  $S = \operatorname{Spec} k$  for a field  $k$ . Here a finite étale  $S$ -scheme  $X$  is the spectrum of a finite étale  $k$ -algebra.

For a geometric point  $\bar{s}$  the fibre functor maps a connected cover  $X = \operatorname{Spec} L$  to the underlying set of  $\operatorname{Spec}(L \otimes_k \Omega)$ , which is a finite set indexed by the  $k$ -algebra homomorphisms  $L \rightarrow \Omega$ .

The image of each such homomorphism lies in the separable closure  $k_s$  of  $k$  in  $\Omega$  via the embedding given by  $\bar{s}$ .

## Revisiting Galois Theory

Let  $S = \operatorname{Spec} k$  for a field  $k$ . Here a finite étale  $S$ -scheme  $X$  is the spectrum of a finite étale  $k$ -algebra.

For a geometric point  $\bar{s}$  the fibre functor maps a connected cover  $X = \operatorname{Spec} L$  to the underlying set of  $\operatorname{Spec}(L \otimes_k \Omega)$ , which is a finite set indexed by the  $k$ -algebra homomorphisms  $L \rightarrow \Omega$ .

The image of each such homomorphism lies in the separable closure  $k_s$  of  $k$  in  $\Omega$  via the embedding given by  $\bar{s}$ .

So finally we obtain that  $\operatorname{Fib}_{\bar{s}}(X) \cong \operatorname{Hom}_k(L, k_s)$  for all  $X$  of the form  $\operatorname{Spec} L$ , and therefore  $\pi_1(S, \bar{s}) \cong \operatorname{Gal}(k_s|k)$ .

## Revisiting Galois Theory

Let  $S = \operatorname{Spec} k$  for a field  $k$ . Here a finite étale  $S$ -scheme  $X$  is the spectrum of a finite étale  $k$ -algebra.

For a geometric point  $\bar{s}$  the fibre functor maps a connected cover  $X = \operatorname{Spec} L$  to the underlying set of  $\operatorname{Spec}(L \otimes_k \Omega)$ , which is a finite set indexed by the  $k$ -algebra homomorphisms  $L \rightarrow \Omega$ .

The image of each such homomorphism lies in the separable closure  $k_s$  of  $k$  in  $\Omega$  via the embedding given by  $\bar{s}$ .

So finally we obtain that  $\operatorname{Fib}_{\bar{s}}(X) \cong \operatorname{Hom}_k(L, k_s)$  for all  $X$  of the form  $\operatorname{Spec} L$ , and therefore  $\pi_1(S, \bar{s}) \cong \operatorname{Gal}(k_s|k)$ .

Thus in this case the theorem above is equivalent to Grothendieck's formulation of Galois Theory.

## Revisiting Galois Theory

Let  $S = \operatorname{Spec} k$  for a field  $k$ . Here a finite étale  $S$ -scheme  $X$  is the spectrum of a finite étale  $k$ -algebra.

For a geometric point  $\bar{s}$  the fibre functor maps a connected cover  $X = \operatorname{Spec} L$  to the underlying set of  $\operatorname{Spec}(L \otimes_k \Omega)$ , which is a finite set indexed by the  $k$ -algebra homomorphisms  $L \rightarrow \Omega$ .

The image of each such homomorphism lies in the separable closure  $k_s$  of  $k$  in  $\Omega$  via the embedding given by  $\bar{s}$ .

So finally we obtain that  $\operatorname{Fib}_{\bar{s}}(X) \cong \operatorname{Hom}_k(L, k_s)$  for all  $X$  of the form  $\operatorname{Spec} L$ , and therefore  $\pi_1(S, \bar{s}) \cong \operatorname{Gal}(k_s|k)$ .

Thus in this case the theorem above is equivalent to Grothendieck's formulation of Galois Theory.

Although in the above example the fibre functor is identified with the functor  $\operatorname{Hom}(\operatorname{Spec} k_s, -)$ , this does not mean that the fibre functor is representable,

## Revisiting Galois Theory

Let  $S = \operatorname{Spec} k$  for a field  $k$ . Here a finite étale  $S$ -scheme  $X$  is the spectrum of a finite étale  $k$ -algebra.

For a geometric point  $\bar{s}$  the fibre functor maps a connected cover  $X = \operatorname{Spec} L$  to the underlying set of  $\operatorname{Spec}(L \otimes_k \Omega)$ , which is a finite set indexed by the  $k$ -algebra homomorphisms  $L \rightarrow \Omega$ .

The image of each such homomorphism lies in the separable closure  $k_s$  of  $k$  in  $\Omega$  via the embedding given by  $\bar{s}$ .

So finally we obtain that  $\operatorname{Fib}_{\bar{s}}(X) \cong \operatorname{Hom}_k(L, k_s)$  for all  $X$  of the form  $\operatorname{Spec} L$ , and therefore  $\pi_1(S, \bar{s}) \cong \operatorname{Gal}(k_s|k)$ .

Thus in this case the theorem above is equivalent to Grothendieck's formulation of Galois Theory.

Although in the above example the fibre functor is identified with the functor  $\operatorname{Hom}(\operatorname{Spec} k_s, -)$ , this does not mean that the fibre functor is representable, for  $k_s$  is not a finite étale  $k$ -algebra.



## Revisiting Galois Theory

Let  $S = \operatorname{Spec} k$  for a field  $k$ . Here a finite étale  $S$ -scheme  $X$  is the spectrum of a finite étale  $k$ -algebra.

For a geometric point  $\bar{s}$  the fibre functor maps a connected cover  $X = \operatorname{Spec} L$  to the underlying set of  $\operatorname{Spec}(L \otimes_k \Omega)$ , which is a finite set indexed by the  $k$ -algebra homomorphisms  $L \rightarrow \Omega$ .

The image of each such homomorphism lies in the separable closure  $k_s$  of  $k$  in  $\Omega$  via the embedding given by  $\bar{s}$ .

So finally we obtain that  $\operatorname{Fib}_{\bar{s}}(X) \cong \operatorname{Hom}_k(L, k_s)$  for all  $X$  of the form  $\operatorname{Spec} L$ , and therefore  $\pi_1(S, \bar{s}) \cong \operatorname{Gal}(k_s|k)$ .

Thus in this case the theorem above is equivalent to Grothendieck's formulation of Galois Theory.

Although in the above example the fibre functor is identified with the functor  $\operatorname{Hom}(\operatorname{Spec} k_s, -)$ , this does not mean that the fibre functor is representable, for  $k_s$  is not a finite étale  $k$ -algebra. However, it is the union of its finite Galois subextensions which are.

## Pro-representable Functors

Let  $\mathcal{C}$  be a category, and  $F$  a set-valued functor on  $\mathcal{C}$ . We say that  $F$  is *pro-representable* if there exists an inverse system  $P = (P_\alpha, \phi_{\alpha\beta})$  of objects of  $\mathcal{C}$  indexed by a directed partially ordered set  $\Gamma$  and a natural isomorphism

$$\varinjlim \operatorname{Hom}(P_\alpha, -) \cong F.$$

## Pro-representable Functors

Let  $\mathcal{C}$  be a category, and  $F$  a set-valued functor on  $\mathcal{C}$ . We say that  $F$  is *pro-representable* if there exists an inverse system  $P = (P_\alpha, \phi_{\alpha\beta})$  of objects of  $\mathcal{C}$  indexed by a directed partially ordered set  $\Gamma$  and a natural isomorphism

$$\varinjlim \operatorname{Hom}(P_\alpha, -) \cong F.$$

Notice that in the above definition the inverse limit of the  $P_\alpha$  may not exist in  $\mathcal{C}$ , but the direct limit of the  $\operatorname{Hom}(P_\alpha, X)$  does in the category of sets.

## Pro-representable Functors

Let  $\mathcal{C}$  be a category, and  $F$  a set-valued functor on  $\mathcal{C}$ . We say that  $F$  is *pro-representable* if there exists an inverse system  $P = (P_\alpha, \phi_{\alpha\beta})$  of objects of  $\mathcal{C}$  indexed by a directed partially ordered set  $\Gamma$  and a natural isomorphism

$$\varinjlim \operatorname{Hom}(P_\alpha, -) \cong F.$$

Notice that in the above definition the inverse limit of the  $P_\alpha$  may not exist in  $\mathcal{C}$ , but the direct limit of the  $\operatorname{Hom}(P_\alpha, X)$  does in the category of sets.

## Pro-representability

$\operatorname{Fib}_{\bar{s}}$  is pro-representable.

## Paths

Let  $S$  be a connected scheme. Given two geometric points  $\bar{s} : \mathrm{Spec}(\Omega) \rightarrow S$  and  $\bar{s}' : \mathrm{Spec}(\Omega') \rightarrow S$ ,

## Paths

Let  $S$  be a connected scheme. Given two geometric points  $\bar{s} : \operatorname{Spec}(\Omega) \rightarrow S$  and  $\bar{s}' : \operatorname{Spec}(\Omega') \rightarrow S$ ,

- (1) there exists an isomorphism of fibre functors  $\operatorname{Fib}_{\bar{s}} \rightarrow \operatorname{Fib}_{\bar{s}'}$ .

## Paths

Let  $S$  be a connected scheme. Given two geometric points  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$  and  $\bar{s}' : \text{Spec}(\Omega') \rightarrow S$ ,

- (1) there exists an isomorphism of fibre functors  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$ .
- (2) there exists a continuous isomorphism of profinite groups  $\pi_1(S, \bar{s}) \rightarrow \pi_1(S, \bar{s}')$ .

## Paths

Let  $S$  be a connected scheme. Given two geometric points  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$  and  $\bar{s}' : \text{Spec}(\Omega') \rightarrow S$ ,

- (1) there exists an isomorphism of fibre functors  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$ .
- (2) there exists a continuous isomorphism of profinite groups  $\pi_1(S, \bar{s}) \rightarrow \pi_1(S, \bar{s}')$ .

An isomorphism  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$  of fibre functors is called a *path* (or *chemin*) from  $\bar{s}$  to  $\bar{s}'$ .



## Paths

Let  $S$  be a connected scheme. Given two geometric points  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$  and  $\bar{s}' : \text{Spec}(\Omega') \rightarrow S$ ,

- (1) there exists an isomorphism of fibre functors  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$ .
- (2) there exists a continuous isomorphism of profinite groups  $\pi_1(S, \bar{s}) \rightarrow \pi_1(S, \bar{s}')$ .

An isomorphism  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$  of fibre functors is called a *path* (or *chemin*) from  $\bar{s}$  to  $\bar{s}'$ .

## Homotopy Exact Sequence

Let  $X$  be a quasi-compact and geometrically integral scheme over a field  $k$ ; that is,  $X_{\bar{k}} := X \times_k \bar{k}$  is integral, for an algebraic closure  $\bar{k}$  of  $k$ .

## Paths

Let  $S$  be a connected scheme. Given two geometric points  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$  and  $\bar{s}' : \text{Spec}(\Omega') \rightarrow S$ ,

- (1) there exists an isomorphism of fibre functors  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$ .
- (2) there exists a continuous isomorphism of profinite groups  $\pi_1(S, \bar{s}) \rightarrow \pi_1(S, \bar{s}')$ .

An isomorphism  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$  of fibre functors is called a *path* (or *chemin*) from  $\bar{s}$  to  $\bar{s}'$ .

## Homotopy Exact Sequence

Let  $X$  be a quasi-compact and geometrically integral scheme over a field  $k$ ; that is,  $X_{\bar{k}} := X \times_k \bar{k}$  is integral, for an algebraic closure  $\bar{k}$  of  $k$ . Fix an algebraic closure  $\bar{k}$  of  $k$ , and let  $k_s|k$  be the corresponding separable closure.

## Paths

Let  $S$  be a connected scheme. Given two geometric points  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$  and  $\bar{s}' : \text{Spec}(\Omega') \rightarrow S$ ,

- (1) there exists an isomorphism of fibre functors  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$ .
- (2) there exists a continuous isomorphism of profinite groups  $\pi_1(S, \bar{s}) \rightarrow \pi_1(S, \bar{s}')$ .

An isomorphism  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$  of fibre functors is called a *path* (or *chemin*) from  $\bar{s}$  to  $\bar{s}'$ .

## Homotopy Exact Sequence

Let  $X$  be a quasi-compact and geometrically integral scheme over a field  $k$ ; that is,  $X_{\bar{k}} := X \times_k \bar{k}$  is integral, for an algebraic closure  $\bar{k}$  of  $k$ . Fix an algebraic closure  $\bar{k}$  of  $k$ , and let  $k_s|k$  be the corresponding separable closure. Let  $\bar{x}$  be a geometric point of  $X_{\bar{k}}$  with values in  $\bar{k}$ .

## Paths

Let  $S$  be a connected scheme. Given two geometric points  $\bar{s} : \text{Spec}(\Omega) \rightarrow S$  and  $\bar{s}' : \text{Spec}(\Omega') \rightarrow S$ ,

- (1) there exists an isomorphism of fibre functors  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$ .
- (2) there exists a continuous isomorphism of profinite groups  $\pi_1(S, \bar{s}) \rightarrow \pi_1(S, \bar{s}')$ .

An isomorphism  $\text{Fib}_{\bar{s}} \rightarrow \text{Fib}_{\bar{s}'}$  of fibre functors is called a *path* (or *chemin*) from  $\bar{s}$  to  $\bar{s}'$ .

## Homotopy Exact Sequence

Let  $X$  be a quasi-compact and geometrically integral scheme over a field  $k$ ; that is,  $X_{\bar{k}} := X \times_k \bar{k}$  is integral, for an algebraic closure  $\bar{k}$  of  $k$ . Fix an algebraic closure  $\bar{k}$  of  $k$ , and let  $k_s|k$  be the corresponding separable closure. Let  $\bar{x}$  be a geometric point of  $X_{\bar{k}}$  with values in  $\bar{k}$ . The sequence of profinite groups

$$1 \longrightarrow \pi_1(X_{\bar{k}}, \bar{x}) \longrightarrow \pi_1(X, \bar{x}) \longrightarrow \text{Gal}(k_s|k) \longrightarrow 1$$

induced by the maps  $X_{\bar{k}} \rightarrow X$  and  $X \rightarrow \text{Spec } k$  is exact.

## Under Base Change

Let  $k \subseteq K$  be an extension of algebraically closed fields, and let  $X$  be a proper integral scheme over  $k$ . Denote  $X_K := X \times_k K$ . The map  $\pi_1(X_K, \bar{x}_K) \rightarrow \pi_1(X, \bar{x})$  induced by the projection  $X_K \rightarrow X$  is an isomorphism for every geometric point  $\bar{x}$  of  $X$ .

## Under Base Change

Let  $k \subseteq K$  be an extension of algebraically closed fields, and let  $X$  be a proper integral scheme over  $k$ . Denote  $X_K := X \times_k K$ . The map  $\pi_1(X_K, \bar{x}_K) \rightarrow \pi_1(X, \bar{x})$  induced by the projection  $X_K \rightarrow X$  is an isomorphism for every geometric point  $\bar{x}$  of  $X$ .

## Comparison with Topological Fundamental Group

Let  $X$  be a connected scheme of finite type over  $\mathbb{C}$ . The analytification functor  $(Y \rightarrow X) \mapsto (Y^{\text{an}} \rightarrow X^{\text{an}})$  induces an equivalence of the category of finite étale covers of  $X$  with that of finite topological covers of  $X^{\text{an}}$ .

## Under Base Change

Let  $k \subseteq K$  be an extension of algebraically closed fields, and let  $X$  be a proper integral scheme over  $k$ . Denote  $X_K := X \times_k K$ . The map  $\pi_1(X_K, \bar{x}_K) \rightarrow \pi_1(X, \bar{x})$  induced by the projection  $X_K \rightarrow X$  is an isomorphism for every geometric point  $\bar{x}$  of  $X$ .

## Comparison with Topological Fundamental Group

Let  $X$  be a connected scheme of finite type over  $\mathbf{C}$ . The analytification functor  $(Y \rightarrow X) \mapsto (Y^{\text{an}} \rightarrow X^{\text{an}})$  induces an equivalence of the category of finite étale covers of  $X$  with that of finite topological covers of  $X^{\text{an}}$ . Consequently, for every  $\mathbf{C}$ -point  $\bar{x} : \text{Spec}(\mathbf{C}) \rightarrow X$  this functor induces an isomorphism

$$\pi_1^{\text{top}}(\widehat{X^{\text{an}}}, \bar{x}) \xrightarrow{\sim} \pi_1^{\text{ét}}(X, \bar{x})$$

# Tannakian Fundamental Group

× × ×



# Representations of Affine Group Schemes

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ .

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .  $A$  then carries additional structure; it is a (commutative) Hopf algebra.

# Representations of Affine Group Schemes

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .  $A$  then carries additional structure; it is a (commutative) Hopf algebra.

## Examples

The *additive group scheme* is the functor  $\mathbf{G}_a : A \mapsto (A, +)$ , the underlying additive group of  $A$ ;

# Representations of Affine Group Schemes

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .  $A$  then carries additional structure; it is a (commutative) Hopf algebra.

## Examples

The *additive group scheme* is the functor  $\mathbf{G}_a : A \mapsto (A, +)$ , the underlying additive group of  $A$ ; it's represented by the  $k$ -algebra  $k[x]$ .

# Representations of Affine Group Schemes

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .  $A$  then carries additional structure; it is a (commutative) Hopf algebra.

## Examples

The *additive group scheme* is the functor  $\mathbf{G}_a : A \mapsto (A, +)$ , the underlying additive group of  $A$ ; it's represented by the  $k$ -algebra  $k[x]$ .

The *multiplicative group scheme* is the functor  $\mathbf{G}_m : A \mapsto A^\times$ , the unit group of  $A$ ;

# Representations of Affine Group Schemes

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .  $A$  then carries additional structure; it is a (commutative) Hopf algebra.

## Examples

The *additive group scheme* is the functor  $\mathbf{G}_a : A \mapsto (A, +)$ , the underlying additive group of  $A$ ; it's represented by the  $k$ -algebra  $k[x]$ .

The *multiplicative group scheme* is the functor  $\mathbf{G}_m : A \mapsto A^\times$ , the unit group of  $A$ ; it's represented by the  $k$ -algebra  $k[x, x^{-1}]$ .

# Representations of Affine Group Schemes

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .  $A$  then carries additional structure; it is a (commutative) Hopf algebra.

## Examples

The *additive group scheme* is the functor  $\mathbf{G}_a : A \mapsto (A, +)$ , the underlying additive group of  $A$ ; it's represented by the  $k$ -algebra  $k[x]$ .

The *multiplicative group scheme* is the functor  $\mathbf{G}_m : A \mapsto A^\times$ , the unit group of  $A$ ; it's represented by the  $k$ -algebra  $k[x, x^{-1}]$ .

More generally, the *general linear group* is the functor  $\mathbf{GL}_n : A \mapsto \mathbf{GL}_n(A)$ , invertible  $n \times n$ -matrices with entries in  $A$ ;



# Representations of Affine Group Schemes

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .  $A$  then carries additional structure; it is a (commutative) Hopf algebra.

## Examples

The *additive group scheme* is the functor  $\mathbf{G}_a : A \mapsto (A, +)$ , the underlying additive group of  $A$ ; it's represented by the  $k$ -algebra  $k[x]$ .

The *multiplicative group scheme* is the functor  $\mathbf{G}_m : A \mapsto A^\times$ , the unit group of  $A$ ; it's represented by the  $k$ -algebra  $k[x, x^{-1}]$ .

More generally, the *general linear group* is the functor  $\mathbf{GL}_n : A \mapsto \mathbf{GL}_n(A)$ , invertible  $n \times n$ -matrices with entries in  $A$ ; it's represented by the  $k$ -algebra  $k[(x_{ij})_{i,j}, d^{-1}]$ , where  $d = \det(x_{ij}) \in k[(x_{ij})_{i,j}]$ .

# Representations of Affine Group Schemes

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .  $A$  then carries additional structure; it is a (commutative) Hopf algebra.

## Examples

The *additive group scheme* is the functor  $\mathbf{G}_a : A \mapsto (A, +)$ , the underlying additive group of  $A$ ; it's represented by the  $k$ -algebra  $k[x]$ .

The *multiplicative group scheme* is the functor  $\mathbf{G}_m : A \mapsto A^\times$ , the unit group of  $A$ ; it's represented by the  $k$ -algebra  $k[x, x^{-1}]$ .

More generally, the *general linear group* is the functor  $\mathbf{GL}_n : A \mapsto \mathbf{GL}_n(A)$ , invertible  $n \times n$ -matrices with entries in  $A$ ; it's represented by the  $k$ -algebra  $k[(x_{ij})_{i,j}, d^{-1}]$ , where  $d = \det(x_{ij}) \in k[(x_{ij})_{i,j}]$ .

The *Deligne torus*  $\mathbf{S} := \text{Res}_{\mathbf{C}/\mathbf{R}} \mathbf{G}_m : \text{Alg}_{\mathbf{R}} \rightarrow \text{Grp}$ ,  $A \mapsto (A \otimes_{\mathbf{R}} \mathbf{C})^\times$ ;

# Representations of Affine Group Schemes

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .  $A$  then carries additional structure; it is a (commutative) Hopf algebra.

## Examples

The *additive group scheme* is the functor  $\mathbf{G}_a : A \mapsto (A, +)$ , the underlying additive group of  $A$ ; it's represented by the  $k$ -algebra  $k[x]$ .

The *multiplicative group scheme* is the functor  $\mathbf{G}_m : A \mapsto A^\times$ , the unit group of  $A$ ; it's represented by the  $k$ -algebra  $k[x, x^{-1}]$ .

More generally, the *general linear group* is the functor  $\mathbf{GL}_n : A \mapsto \mathbf{GL}_n(A)$ , invertible  $n \times n$ -matrices with entries in  $A$ ; it's represented by the  $k$ -algebra  $k[(x_{ij})_{i,j}, d^{-1}]$ , where  $d = \det(x_{ij}) \in k[(x_{ij})_{i,j}]$ .

The *Deligne torus*  $\mathbf{S} := \text{Res}_{\mathbf{C}/\mathbf{R}} \mathbf{G}_m : \text{Alg}_{\mathbf{R}} \rightarrow \text{Grp}$ ,  $A \mapsto (A \otimes_{\mathbf{R}} \mathbf{C})^\times$ ; it's represented by the  $\mathbf{R}$ -algebra  $\mathbf{R}[a, b, x, y, (a^2 + b^2)^{-1}] / (x + b, y - a)$ .

# Representations of Affine Group Schemes

## Affine Group Schemes

Let  $k$  be a field. An *affine group scheme*  $G$  over  $k$  is a functor from  $\text{Alg}_k \rightarrow \text{Grp}$  that, when viewed as a set-valued functor, is representable by some  $k$ -algebra  $A$ . In particular, it corresponds to the affine scheme  $\text{Spec } A$ .  $A$  then carries additional structure; it is a (commutative) Hopf algebra.

## Examples

The *additive group scheme* is the functor  $\mathbf{G}_a : A \mapsto (A, +)$ , the underlying additive group of  $A$ ; it's represented by the  $k$ -algebra  $k[x]$ .

The *multiplicative group scheme* is the functor  $\mathbf{G}_m : A \mapsto A^\times$ , the unit group of  $A$ ; it's represented by the  $k$ -algebra  $k[x, x^{-1}]$ .

More generally, the *general linear group* is the functor  $\mathbf{GL}_n : A \mapsto \mathbf{GL}_n(A)$ , invertible  $n \times n$ -matrices with entries in  $A$ ; it's represented by the  $k$ -algebra  $k[(x_{ij})_{i,j}, d^{-1}]$ , where  $d = \det(x_{ij}) \in k[(x_{ij})_{i,j}]$ .

The *Deligne torus*  $\mathbf{S} := \text{Res}_{\mathbf{C}/\mathbf{R}} \mathbf{G}_m : \text{Alg}_{\mathbf{R}} \rightarrow \text{Grp}$ ,  $A \mapsto (A \otimes_{\mathbf{R}} \mathbf{C})^\times$ ; it's represented by the  $\mathbf{R}$ -algebra  $\mathbf{R}[a, b, x, y, (a^2 + b^2)^{-1}] / (x + b, y - a)$ . In particular,  $\mathbf{S}(\mathbf{R}) = \mathbf{C}^\times$  and  $\mathbf{S}(\mathbf{C}) = \mathbf{C}^\times \times \mathbf{C}^\times$ .

## Coalgebras & Comodules

A *coalgebra* over  $k$  is a  $k$ -vector space equipped with a comultiplication  $\Delta : A \rightarrow A \otimes_k A$  and a counit map  $\iota : A \rightarrow k$  subject to the coassociativity and counit axioms. Here,  $\Delta$  and  $\iota$  are only assumed to be maps of  $k$ -vector spaces.

## Coalgebras & Comodules

A *coalgebra* over  $k$  is a  $k$ -vector space equipped with a comultiplication  $\Delta : A \rightarrow A \otimes_k A$  and a counit map  $\iota : A \rightarrow k$  subject to the coassociativity and counit axioms. Here,  $\Delta$  and  $\iota$  are only assumed to be maps of  $k$ -vector spaces.

Let  $A$  be a coalgebra over a field  $k$ . A *right  $A$ -comodule* is a  $k$ -vector space  $M$  together with a  $k$ -linear map  $\rho : M \rightarrow M \otimes_k A$  such that certain diagrams commute.

# Representations of Affine Group Schemes

## Coalgebras & Comodules

A *coalgebra* over  $k$  is a  $k$ -vector space equipped with a comultiplication  $\Delta : A \rightarrow A \otimes_k A$  and a counit map  $\iota : A \rightarrow k$  subject to the coassociativity and counit axioms. Here,  $\Delta$  and  $\iota$  are only assumed to be maps of  $k$ -vector spaces.

Let  $A$  be a coalgebra over a field  $k$ . A *right  $A$ -comodule* is a  $k$ -vector space  $M$  together with a  $k$ -linear map  $\rho : M \rightarrow M \otimes_k A$  such that certain diagrams commute.

## Representations of Affine Group Schemes

In the case  $A$  is a commutative Hopf algebra, an  $A$ -comodule  $M$  gives rise to a representation of the corresponding affine group scheme  $G$ .

# Representations of Affine Group Schemes

## Coalgebras & Comodules

A *coalgebra* over  $k$  is a  $k$ -vector space equipped with a comultiplication  $\Delta : A \rightarrow A \otimes_k A$  and a counit map  $\iota : A \rightarrow k$  subject to the coassociativity and counit axioms. Here,  $\Delta$  and  $\iota$  are only assumed to be maps of  $k$ -vector spaces.

Let  $A$  be a coalgebra over a field  $k$ . A *right  $A$ -comodule* is a  $k$ -vector space  $M$  together with a  $k$ -linear map  $\rho : M \rightarrow M \otimes_k A$  such that certain diagrams commute.

## Representations of Affine Group Schemes

In the case  $A$  is a commutative Hopf algebra, an  $A$ -comodule  $M$  gives rise to a representation of the corresponding affine group scheme  $G$ .

Given a  $k$ -algebra  $R$ , an element of  $G(R)$  corresponds to a  $k$ -algebra morphism  $\lambda : A \rightarrow R$ . The composite

$$M \xrightarrow{\rho} M \otimes_k A \xrightarrow{\text{id} \otimes \lambda} M \otimes R$$

induces an  $R$ -linear map  $M \otimes_k R \rightarrow M \otimes_k R$  that depends on  $R$  in a functorial way.



# Representations of Affine Group Schemes

## Coalgebras & Comodules

A *coalgebra* over  $k$  is a  $k$ -vector space equipped with a comultiplication  $\Delta : A \rightarrow A \otimes_k A$  and a counit map  $\iota : A \rightarrow k$  subject to the coassociativity and counit axioms. Here,  $\Delta$  and  $\iota$  are only assumed to be maps of  $k$ -vector spaces.

Let  $A$  be a coalgebra over a field  $k$ . A *right  $A$ -comodule* is a  $k$ -vector space  $M$  together with a  $k$ -linear map  $\rho : M \rightarrow M \otimes_k A$  such that certain diagrams commute.

## Representations of Affine Group Schemes

In the case  $A$  is a commutative Hopf algebra, an  $A$ -comodule  $M$  gives rise to a representation of the corresponding affine group scheme  $G$ .

Given a  $k$ -algebra  $R$ , an element of  $G(R)$  corresponds to a  $k$ -algebra morphism  $\lambda : A \rightarrow R$ . The composite

$$M \xrightarrow{\rho} M \otimes_k A \xrightarrow{\text{id} \otimes \lambda} M \otimes R$$

induces an  $R$ -linear map  $M \otimes_k R \rightarrow M \otimes_k R$  that depends on  $R$  in a functorial way. By the comodule axioms, we thus obtain a functorial collection of left group actions  $G(R) \times (M \otimes_k R) \rightarrow M \otimes_k R$  such that for each  $g \in G(R)$  the map  $m \mapsto gm$  is  $R$ -linear.

# Representations of Affine Group Schemes

## Coalgebras & Comodules

A *coalgebra* over  $k$  is a  $k$ -vector space equipped with a comultiplication  $\Delta : A \rightarrow A \otimes_k A$  and a counit map  $\iota : A \rightarrow k$  subject to the coassociativity and counit axioms. Here,  $\Delta$  and  $\iota$  are only assumed to be maps of  $k$ -vector spaces.

Let  $A$  be a coalgebra over a field  $k$ . A *right  $A$ -comodule* is a  $k$ -vector space  $M$  together with a  $k$ -linear map  $\rho : M \rightarrow M \otimes_k A$  such that certain diagrams commute.

## Representations of Affine Group Schemes

In the case  $A$  is a commutative Hopf algebra, an  $A$ -comodule  $M$  gives rise to a representation of the corresponding affine group scheme  $G$ .

Given a  $k$ -algebra  $R$ , an element of  $G(R)$  corresponds to a  $k$ -algebra morphism  $\lambda : A \rightarrow R$ . The composite

$$M \xrightarrow{\rho} M \otimes_k A \xrightarrow{\text{id} \otimes \lambda} M \otimes R$$

induces an  $R$ -linear map  $M \otimes_k R \rightarrow M \otimes_k R$  that depends on  $R$  in a functorial way. By the comodule axioms, we thus obtain a functorial collection of left group actions  $G(R) \times (M \otimes_k R) \rightarrow M \otimes_k R$  such that for each  $g \in G(R)$  the map  $m \mapsto gm$  is  $R$ -linear. We call such a collection a left representation of the affine group scheme  $G$ .

# Representations of Affine Group Schemes

## Representations of Affine Group Schemes (contd.)

If moreover  $M$  is finite dimensional over  $k$  and we fix a  $k$ -basis of  $M$ , giving a representation of  $G$  becomes equivalent to giving a morphism of group schemes  $G \rightarrow \mathbf{GL}_n$ , i.e. a morphism of group-valued functors.

# Representations of Affine Group Schemes

## Representations of Affine Group Schemes (contd.)

If moreover  $M$  is finite dimensional over  $k$  and we fix a  $k$ -basis of  $M$ , giving a representation of  $G$  becomes equivalent to giving a morphism of group schemes  $G \rightarrow \mathbf{GL}_n$ , i.e. a morphism of group-valued functors.

The construction given gives a bijection between right comodules over the commutative Hopf algebra  $A$  and left representations of the corresponding affine group scheme  $G$ .

# Representations of Affine Group Schemes

## Representations of Affine Group Schemes (contd.)

If moreover  $M$  is finite dimensional over  $k$  and we fix a  $k$ -basis of  $M$ , giving a representation of  $G$  becomes equivalent to giving a morphism of group schemes  $G \rightarrow \mathbf{GL}_n$ , i.e. a morphism of group-valued functors.

The construction given gives a bijection between right comodules over the commutative Hopf algebra  $A$  and left representations of the corresponding affine group scheme  $G$ .

## Duals

Given a  $k$ -coalgebra  $A$ , the  $k$ -linear dual  $A^* := \operatorname{Hom}_k(A, k)$  is a  $k$ -algebra with unit that is not necessarily commutative.

# Representations of Affine Group Schemes

## Representations of Affine Group Schemes (contd.)

If moreover  $M$  is finite dimensional over  $k$  and we fix a  $k$ -basis of  $M$ , giving a representation of  $G$  becomes equivalent to giving a morphism of group schemes  $G \rightarrow \mathbf{GL}_n$ , i.e. a morphism of group-valued functors.

The construction given gives a bijection between right comodules over the commutative Hopf algebra  $A$  and left representations of the corresponding affine group scheme  $G$ .

## Duals

Given a  $k$ -coalgebra  $A$ , the  $k$ -linear dual  $A^* := \text{Hom}_k(A, k)$  is a  $k$ -algebra with unit that is not necessarily commutative.

The contravariant functor  $A \mapsto A^*$  induces an anti-isomorphism between the category of  $k$ -coalgebras finite dimensional over  $k$  and that of not necessarily commutative finite dimensional  $k$ -algebras with unit.

# Representations of Affine Group Schemes

## Representations of Affine Group Schemes (contd.)

If moreover  $M$  is finite dimensional over  $k$  and we fix a  $k$ -basis of  $M$ , giving a representation of  $G$  becomes equivalent to giving a morphism of group schemes  $G \rightarrow \mathbf{GL}_n$ , i.e. a morphism of group-valued functors.

The construction given gives a bijection between right comodules over the commutative Hopf algebra  $A$  and left representations of the corresponding affine group scheme  $G$ .

## Duals

Given a  $k$ -coalgebra  $A$ , the  $k$ -linear dual  $A^* := \text{Hom}_k(A, k)$  is a  $k$ -algebra with unit that is not necessarily commutative.

The contravariant functor  $A \mapsto A^*$  induces an anti-isomorphism between the category of  $k$ -coalgebras finite dimensional over  $k$  and that of not necessarily commutative finite dimensional  $k$ -algebras with unit.

Given a finite dimensional  $k$ -coalgebra  $A$ , the contravariant functor  $V \mapsto V^*$  induces an anti-isomorphism between the category of finitely generated right  $A$ -comodules and that of finitely generated left  $A^*$ -modules.

## [Prefixes] Tensor Categories

A *tensor category (with a unit)*, i.e. a *monoidal category*, is a category  $\mathcal{C}$  together with

- a functor  $- \otimes - : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ ;
- a natural isomorphism  $\alpha_{X,Y,Z} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$ ;
- an object  $\mathbb{1} \in \mathcal{C}$ ;
- a natural isomorphism  $\lambda_X : \mathbb{1} \otimes X \rightarrow X$ ;
- a natural isomorphism  $\rho_X : X \otimes \mathbb{1} \rightarrow X$ ;

such that certain diagrams commute.



# Tensor Categories

## [Prefixes] Tensor Categories

A *tensor category (with a unit)*, i.e. a *monoidal category*, is a category  $\mathcal{C}$  together with

- a functor  $- \otimes - : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ ;
- a natural isomorphism  $\alpha_{X,Y,Z} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$ ;
- an object  $\mathbb{1} \in \mathcal{C}$ ;
- a natural isomorphism  $\lambda_X : \mathbb{1} \otimes X \rightarrow X$ ;
- a natural isomorphism  $\rho_X : X \otimes \mathbb{1} \rightarrow X$ ;

such that certain diagrams commute.

Furthermore, if there's

- a natural isomorphism  $\tau_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ ;

such that a certain diagram commutes, then a tensor category is called an *abelian tensor category*, i.e. a *symmetric monoidal category*.

## [Prefixes] Tensor Categories

A *tensor category (with a unit)*, i.e. a *monoidal category*, is a category  $\mathcal{C}$  together with

- a functor  $- \otimes - : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ ;
- a natural isomorphism  $\alpha_{X,Y,Z} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$ ;
- an object  $\mathbb{1} \in \mathcal{C}$ ;
- a natural isomorphism  $\lambda_X : \mathbb{1} \otimes X \rightarrow X$ ;
- a natural isomorphism  $\rho_X : X \otimes \mathbb{1} \rightarrow X$ ;

such that certain diagrams commute.

Furthermore, if there's

- a natural isomorphism  $\tau_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ ;

such that a certain diagram commutes, then a tensor category is called an *abelian tensor category*, i.e. a *symmetric monoidal category*.

A *tensor functor* between two tensor categories  $\mathcal{C}$  and  $\mathcal{D}$  is a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  together with an isomorphism  $\phi_0 : \mathbb{1}_{\mathcal{D}} \rightarrow F(\mathbb{1}_{\mathcal{C}})$  and a natural isomorphism  $\phi_{X,Y} : F(X) \otimes F(Y) \rightarrow F(X \otimes Y)$  such that certain diagrams commute.

## [Prefixes] Tensor Categories

A *tensor category (with a unit)*, i.e. a *monoidal category*, is a category  $\mathcal{C}$  together with

- a functor  $- \otimes - : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ ;
- a natural isomorphism  $\alpha_{X,Y,Z} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$ ;
- an object  $\mathbb{1} \in \mathcal{C}$ ;
- a natural isomorphism  $\lambda_X : \mathbb{1} \otimes X \rightarrow X$ ;
- a natural isomorphism  $\rho_X : X \otimes \mathbb{1} \rightarrow X$ ;

such that certain diagrams commute.

Furthermore, if there's

- a natural isomorphism  $\tau_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ ;

such that a certain diagram commutes, then a tensor category is called an *abelian tensor category*, i.e. a *symmetric monoidal category*.

A *tensor functor* between two tensor categories  $\mathcal{C}$  and  $\mathcal{D}$  is a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  together with an isomorphism  $\phi_0 : \mathbb{1}_{\mathcal{D}} \rightarrow F(\mathbb{1}_{\mathcal{C}})$  and a natural isomorphism  $\phi_{X,Y} : F(X) \otimes F(Y) \rightarrow F(X \otimes Y)$  such that certain diagrams commute.

We call a tensor category *rigid* if for each object  $X$ , there's a *dual*  $X^*$ ,

## [Prefixes] Tensor Categories

A *tensor category (with a unit)*, i.e. a *monoidal category*, is a category  $\mathcal{C}$  together with

- a functor  $- \otimes - : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ ;
- a natural isomorphism  $\alpha_{X,Y,Z} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z)$ ;
- an object  $\mathbb{1} \in \mathcal{C}$ ;
- a natural isomorphism  $\lambda_X : \mathbb{1} \otimes X \rightarrow X$ ;
- a natural isomorphism  $\rho_X : X \otimes \mathbb{1} \rightarrow X$ ;

such that certain diagrams commute.

Furthermore, if there's

- a natural isomorphism  $\tau_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ ;

such that a certain diagram commutes, then a tensor category is called an *abelian tensor category*, i.e. a *symmetric monoidal category*.

A *tensor functor* between two tensor categories  $\mathcal{C}$  and  $\mathcal{D}$  is a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  together with an isomorphism  $\phi_0 : \mathbb{1}_{\mathcal{D}} \rightarrow F(\mathbb{1}_{\mathcal{C}})$  and a natural isomorphism  $\phi_{X,Y} : F(X) \otimes F(Y) \rightarrow F(X \otimes Y)$  such that certain diagrams commute.

We call a tensor category *rigid* if for each object  $X$ , there's a *dual*  $X^*$ , that is there exist morphisms  $\varepsilon_X : X \otimes X^* \rightarrow \mathbb{1}$  and  $\delta_X : \mathbb{1} \rightarrow X^* \otimes X$  so that certain diagrams commute.

## Revisiting $\text{Comod}_A$

Let  $A$  be a coalgebra over a field  $k$ , and  $\omega$  the forgetful functor from  $\text{Comod}_A$  to  $\text{Vecf}_k$ .

Assume that there is a tensor category structure on  $\text{Comod}_A$  for which  $\omega$  becomes a tensor functor, where  $\text{Vecf}_k$  carries its usual tensor structure.

## Revisiting $\text{Comod}_A$

Let  $A$  be a coalgebra over a field  $k$ , and  $\omega$  the forgetful functor from  $\text{Comod}_A$  to  $\text{Vecf}_k$ .

Assume that there is a tensor category structure on  $\text{Comod}_A$  for which  $\omega$  becomes a tensor functor, where  $\text{Vecf}_k$  carries its usual tensor structure.

- There is a canonical  $k$ -algebra structure with unit on  $A$  defined by means of coalgebra morphisms.

## Revisiting $\text{Comod}_A$

Let  $A$  be a coalgebra over a field  $k$ , and  $\omega$  the forgetful functor from  $\text{Comod}_A$  to  $\text{Vec}_k$ .

Assume that there is a tensor category structure on  $\text{Comod}_A$  for which  $\omega$  becomes a tensor functor, where  $\text{Vec}_k$  carries its usual tensor structure.

- There is a canonical  $k$ -algebra structure with unit on  $A$  defined by means of coalgebra morphisms.
- If moreover the tensor category structure on  $\text{Comod}_A$  is rigid, then  $A$  has the structure of a Hopf algebra.

## Revisiting $\text{Comodf}_A$

Let  $A$  be a coalgebra over a field  $k$ , and  $\omega$  the forgetful functor from  $\text{Comodf}_A$  to  $\text{Vecf}_k$ .

Assume that there is a tensor category structure on  $\text{Comodf}_A$  for which  $\omega$  becomes a tensor functor, where  $\text{Vecf}_k$  carries its usual tensor structure.

- There is a canonical  $k$ -algebra structure with unit on  $A$  defined by means of coalgebra morphisms.
- If moreover the tensor category structure on  $\text{Comodf}_A$  is rigid, then  $A$  has the structure of a Hopf algebra.
- Assume moreover the tensor category structure on  $\text{Comodf}_A$  is commutative, and  $\omega$  respects the commutativity constraints. Then  $A$  is a commutative Hopf algebra, and  $\text{Comodf}_A$  becomes equivalent to the category  $\text{Rep}_G$  of finite dimensional representations of the associated affine group scheme  $G$ .



## More on $\text{Rep}_G$

Observe that given a commutative  $k$ -algebra  $R$ , the forgetful functor  $\omega$  on  $\text{Rep}_G$  induces a tensor functor  $\omega \otimes R : V \mapsto V \otimes_k R$  with values in the tensor category of  $R$ -modules.

## More on $\text{Rep}_G$

Observe that given a commutative  $k$ -algebra  $R$ , the forgetful functor  $\omega$  on  $\text{Rep}_G$  induces a tensor functor  $\omega \otimes R : V \mapsto V \otimes_k R$  with values in the tensor category of  $R$ -modules.

We can thus define the group-valued functor  $\mathbf{Aut}^\otimes(\omega)$  on the category of  $k$ -algebras by sending  $R$  to the group of  $R$ -linear tensor functor isomorphisms  $\omega \otimes R \rightarrow \omega \otimes R$ .

## More on $\text{Rep}_G$

Observe that given a commutative  $k$ -algebra  $R$ , the forgetful functor  $\omega$  on  $\text{Rep}_G$  induces a tensor functor  $\omega \otimes R : V \mapsto V \otimes_k R$  with values in the tensor category of  $R$ -modules.

We can thus define the group-valued functor  $\mathbf{Aut}^\otimes(\omega)$  on the category of  $k$ -algebras by sending  $R$  to the group of  $R$ -linear tensor functor isomorphisms  $\omega \otimes R \rightarrow \omega \otimes R$ .

*There is a canonical isomorphism of group-valued functors  $G \rightarrow \mathbf{Aut}^\otimes(\omega)$ . Consequently, we have that  $\mathbf{Aut}^\otimes(\omega)$  is an affine group scheme.*

## More on $\text{Rep}_G$

Observe that given a commutative  $k$ -algebra  $R$ , the forgetful functor  $\omega$  on  $\text{Rep}_G$  induces a tensor functor  $\omega \otimes R : V \mapsto V \otimes_k R$  with values in the tensor category of  $R$ -modules.

We can thus define the group-valued functor  $\mathbf{Aut}^\otimes(\omega)$  on the category of  $k$ -algebras by sending  $R$  to the group of  $R$ -linear tensor functor isomorphisms  $\omega \otimes R \rightarrow \omega \otimes R$ .

*There is a canonical isomorphism of group-valued functors  $G \rightarrow \mathbf{Aut}^\otimes(\omega)$ . Consequently, we have that  $\mathbf{Aut}^\otimes(\omega)$  is an affine group scheme.*

This shows that one may recover an affine group scheme from the tensor category of its representations.

## More on $\text{Rep}_G$

Observe that given a commutative  $k$ -algebra  $R$ , the forgetful functor  $\omega$  on  $\text{Rep}_G$  induces a tensor functor  $\omega \otimes R : V \mapsto V \otimes_k R$  with values in the tensor category of  $R$ -modules.

We can thus define the group-valued functor  $\mathbf{Aut}^\otimes(\omega)$  on the category of  $k$ -algebras by sending  $R$  to the group of  $R$ -linear tensor functor isomorphisms  $\omega \otimes R \rightarrow \omega \otimes R$ .

*There is a canonical isomorphism of group-valued functors  $G \rightarrow \mathbf{Aut}^\otimes(\omega)$ . Consequently, we have that  $\mathbf{Aut}^\otimes(\omega)$  is an affine group scheme.*

This shows that one may recover an affine group scheme from the tensor category of its representations.

This is the algebraic analogue of the classical Tannaka–Krein theorem on topological groups which states that the continuous irreducible unitary representations of a compact group determine the group.

## More on $\text{Rep}_G$

Observe that given a commutative  $k$ -algebra  $R$ , the forgetful functor  $\omega$  on  $\text{Rep}_G$  induces a tensor functor  $\omega \otimes R : V \mapsto V \otimes_k R$  with values in the tensor category of  $R$ -modules.

We can thus define the group-valued functor  $\mathbf{Aut}^\otimes(\omega)$  on the category of  $k$ -algebras by sending  $R$  to the group of  $R$ -linear tensor functor isomorphisms  $\omega \otimes R \rightarrow \omega \otimes R$ .

*There is a canonical isomorphism of group-valued functors  $G \rightarrow \mathbf{Aut}^\otimes(\omega)$ . Consequently, we have that  $\mathbf{Aut}^\otimes(\omega)$  is an affine group scheme.*

This shows that one may recover an affine group scheme from the tensor category of its representations.

This is the algebraic analogue of the classical Tannaka–Krein theorem on topological groups which states that the continuous irreducible unitary representations of a compact group determine the group.

Now let  $G$  and  $H$  be affine group schemes. Given a group scheme homomorphism  $\phi : G \rightarrow H$ , every finite dimensional representation of  $H$  yields a representation of  $G$  via composition with  $\phi$ .

## More on $\text{Rep}_G$

Observe that given a commutative  $k$ -algebra  $R$ , the forgetful functor  $\omega$  on  $\text{Rep}_G$  induces a tensor functor  $\omega \otimes R : V \mapsto V \otimes_k R$  with values in the tensor category of  $R$ -modules.

We can thus define the group-valued functor  $\mathbf{Aut}^{\otimes}(\omega)$  on the category of  $k$ -algebras by sending  $R$  to the group of  $R$ -linear tensor functor isomorphisms  $\omega \otimes R \rightarrow \omega \otimes R$ .

*There is a canonical isomorphism of group-valued functors  $G \rightarrow \mathbf{Aut}^{\otimes}(\omega)$ . Consequently, we have that  $\mathbf{Aut}^{\otimes}(\omega)$  is an affine group scheme.*

This shows that one may recover an affine group scheme from the tensor category of its representations.

This is the algebraic analogue of the classical Tannaka–Krein theorem on topological groups which states that the continuous irreducible unitary representations of a compact group determine the group.

Now let  $G$  and  $H$  be affine group schemes. Given a group scheme homomorphism  $\phi : G \rightarrow H$ , every finite dimensional representation of  $H$  yields a representation of  $G$  via composition with  $\phi$ . In this way we obtain a tensor functor  $\phi^* : \text{Rep}_H \rightarrow \text{Rep}_G$  satisfying  $\omega_G \circ \phi^* = \omega_H$ .

## More on $\text{Rep}_G$

Observe that given a commutative  $k$ -algebra  $R$ , the forgetful functor  $\omega$  on  $\text{Rep}_G$  induces a tensor functor  $\omega \otimes R : V \mapsto V \otimes_k R$  with values in the tensor category of  $R$ -modules.

We can thus define the group-valued functor  $\mathbf{Aut}^{\otimes}(\omega)$  on the category of  $k$ -algebras by sending  $R$  to the group of  $R$ -linear tensor functor isomorphisms  $\omega \otimes R \rightarrow \omega \otimes R$ .

*There is a canonical isomorphism of group-valued functors  $G \rightarrow \mathbf{Aut}^{\otimes}(\omega)$ . Consequently, we have that  $\mathbf{Aut}^{\otimes}(\omega)$  is an affine group scheme.*

This shows that one may recover an affine group scheme from the tensor category of its representations.

This is the algebraic analogue of the classical Tannaka–Krein theorem on topological groups which states that the continuous irreducible unitary representations of a compact group determine the group.

Now let  $G$  and  $H$  be affine group schemes. Given a group scheme homomorphism  $\phi : G \rightarrow H$ , every finite dimensional representation of  $H$  yields a representation of  $G$  via composition with  $\phi$ . In this way we obtain a tensor functor  $\phi^* : \text{Rep}_H \rightarrow \text{Rep}_G$  satisfying  $\omega_G \circ \phi^* = \omega_H$ .

*The rule  $\phi \mapsto \phi^*$  induces a bijection between group scheme homomorphisms  $G \rightarrow H$  and tensor functors  $F : \text{Rep}_H \rightarrow \text{Rep}_G$  satisfying  $\omega_G \circ F = \omega_H$ .*



## Neutral Tannakian Categories

A *neutral Tannakian category* over a field  $k$  is a rigid  $k$ -linear abelian tensor category  $\mathcal{C}$  whose unit  $1$  satisfies  $\text{End}(1) \cong k$ ,

## Neutral Tannakian Categories

A *neutral Tannakian category* over a field  $k$  is a *rigid  $k$ -linear abelian tensor category*  $\mathcal{C}$  whose unit  $1$  satisfies  $\text{End}(1) \cong k$ , and is moreover equipped with an exact faithful tensor functor  $\omega : \mathcal{C} \rightarrow \text{Vecf}_k$ .

## Neutral Tannakian Categories

A *neutral Tannakian category* over a field  $k$  is a *rigid  $k$ -linear abelian tensor category*  $\mathcal{C}$  whose unit  $1$  satisfies  $\text{End}(1) \cong k$ , and is moreover equipped with an exact faithful tensor functor  $\omega : \mathcal{C} \rightarrow \text{Vecf}_k$ .

The functor  $\omega$  is called a (*neutral*) *fibre functor*.

## Neutral Tannakian Categories

A *neutral Tannakian category* over a field  $k$  is a *rigid  $k$ -linear abelian tensor category*  $\mathcal{C}$  whose unit  $1$  satisfies  $\text{End}(1) \cong k$ , and is moreover equipped with an exact faithful tensor functor  $\omega : \mathcal{C} \rightarrow \text{Vecf}_k$ .

The functor  $\omega$  is called a *(neutral) fibre functor*.

Given an affine group scheme  $G$  over  $k$ , the category  $\text{Rep}_G$  of finite dimensional representations of  $G$  with its usual tensor structure and the forgetful functor as fibre functor is a neutral Tannakian category.

## Neutral Tannakian Categories

A *neutral Tannakian category* over a field  $k$  is a *rigid  $k$ -linear abelian tensor category*  $\mathcal{C}$  whose unit  $\mathbb{1}$  satisfies  $\text{End}(\mathbb{1}) \cong k$ , and is moreover equipped with an exact faithful tensor functor  $\omega : \mathcal{C} \rightarrow \text{Vecf}_k$ .

The functor  $\omega$  is called a (*neutral*) *fibre functor*.

Given an affine group scheme  $G$  over  $k$ , the category  $\text{Rep}_G$  of finite dimensional representations of  $G$  with its usual tensor structure and the forgetful functor as fibre functor is a neutral Tannakian category.

## Tannakian Fundamental Group

Every neutral Tannakian category  $(\mathcal{C}, \omega)$  over  $k$  is equivalent to the category  $\text{Rep}_G$  of finite dimensional representations of an affine group scheme  $G$  over  $k$ .

# Tannakian Fundamental Group

## Neutral Tannakian Categories

A *neutral Tannakian category* over a field  $k$  is a *rigid  $k$ -linear abelian tensor category*  $\mathcal{C}$  whose unit  $1$  satisfies  $\text{End}(1) \cong k$ , and is moreover equipped with an exact faithful tensor functor  $\omega : \mathcal{C} \rightarrow \text{Vecf}_k$ .

The functor  $\omega$  is called a (*neutral*) *fibre functor*.

Given an affine group scheme  $G$  over  $k$ , the category  $\text{Rep}_G$  of finite dimensional representations of  $G$  with its usual tensor structure and the forgetful functor as fibre functor is a neutral Tannakian category.

## Tannakian Fundamental Group

Every neutral Tannakian category  $(\mathcal{C}, \omega)$  over  $k$  is equivalent to the category  $\text{Rep}_G$  of finite dimensional representations of an affine group scheme  $G$  over  $k$ .

The group scheme  $G$  as a functor is isomorphic to the functor  $\mathbf{G}(\mathcal{C}, \omega) = \mathbf{Aut}^{\otimes}(\omega)$  of tensor automorphisms of the fibre functor  $\omega$ .

# Tannakian Fundamental Group

## Neutral Tannakian Categories

A *neutral Tannakian category* over a field  $k$  is a *rigid  $k$ -linear abelian tensor category*  $\mathcal{C}$  whose unit  $1$  satisfies  $\text{End}(1) \cong k$ , and is moreover equipped with an exact faithful tensor functor  $\omega : \mathcal{C} \rightarrow \text{Vecf}_k$ .

The functor  $\omega$  is called a (*neutral*) *fibre functor*.

Given an affine group scheme  $G$  over  $k$ , the category  $\text{Rep}_G$  of finite dimensional representations of  $G$  with its usual tensor structure and the forgetful functor as fibre functor is a neutral Tannakian category.

## Tannakian Fundamental Group

Every neutral Tannakian category  $(\mathcal{C}, \omega)$  over  $k$  is equivalent to the category  $\text{Rep}_G$  of finite dimensional representations of an affine group scheme  $G$  over  $k$ .

The group scheme  $G$  as a functor is isomorphic to the functor  $\mathbf{G}(\mathcal{C}, \omega) = \mathbf{Aut}^{\otimes}(\omega)$  of tensor automorphisms of the fibre functor  $\omega$ . We call it the *Tannakian fundamental group* of  $(\mathcal{C}, \omega)$ .

# Examples

## Tannakian Fundamental Group (contd.)

Often,  $G(\mathcal{C}, \omega)$  is “too large” to handle directly.



## Tannakian Fundamental Group (contd.)

Often,  $\mathbf{G}(\mathcal{C}, \omega)$  is “too large” to handle directly. For  $X \in \mathcal{C}$ , let  $\langle X \rangle_{\otimes}$  be the Tannakian subcategory of  $\mathcal{C}$  generated by  $X$ . One puts  $\mathbf{G}(X) = \mathbf{G}(\langle X \rangle_{\otimes}, \omega)$ .

## Tannakian Fundamental Group (contd.)

Often,  $\mathbf{G}(\mathcal{C}, \omega)$  is “too large” to handle directly. For  $X \in \mathcal{C}$ , let  $\langle X \rangle_{\otimes}$  be the Tannakian subcategory of  $\mathcal{C}$  generated by  $X$ . One puts  $\mathbf{G}(X) = \mathbf{G}(\langle X \rangle_{\otimes}, \omega)$ . It turns out that  $\mathbf{G}(X)$  is finite-dimensional and  $\mathbf{G}(\mathcal{C}, \omega) = \varprojlim \mathbf{G}(X)$ .

# Examples

## Tannakian Fundamental Group (contd.)

Often,  $\mathbf{G}(\mathcal{C}, \omega)$  is “too large” to handle directly. For  $X \in \mathcal{C}$ , let  $\langle X \rangle_{\otimes}$  be the Tannakian subcategory of  $\mathcal{C}$  generated by  $X$ . One puts  $\mathbf{G}(X) = \mathbf{G}(\langle X \rangle_{\otimes}, \omega)$ . It turns out that  $\mathbf{G}(X)$  is finite-dimensional and  $\mathbf{G}(\mathcal{C}, \omega) = \varprojlim \mathbf{G}(X)$ .

## Familiar Examples

Let  $\mathcal{C} = \text{Vecf}_k$ , this is a neutral Tannakian category with the fibre functor being the identity.

# Examples

## Tannakian Fundamental Group (contd.)

Often,  $\mathbf{G}(\mathcal{C}, \omega)$  is “too large” to handle directly. For  $X \in \mathcal{C}$ , let  $\langle X \rangle_{\otimes}$  be the Tannakian subcategory of  $\mathcal{C}$  generated by  $X$ . One puts  $\mathbf{G}(X) = \mathbf{G}(\langle X \rangle_{\otimes}, \omega)$ . It turns out that  $\mathbf{G}(X)$  is finite-dimensional and  $\mathbf{G}(\mathcal{C}, \omega) = \varprojlim \mathbf{G}(X)$ .

## Familiar Examples

Let  $\mathcal{C} = \text{Vecf}_k$ , this is a neutral Tannakian category with the fibre functor being the identity. Then  $\mathbf{G}(\mathcal{C}, \omega) = \text{Spec } k$ .

# Examples

## Tannakian Fundamental Group (contd.)

Often,  $\mathbf{G}(\mathcal{C}, \omega)$  is “too large” to handle directly. For  $X \in \mathcal{C}$ , let  $\langle X \rangle_{\otimes}$  be the Tannakian subcategory of  $\mathcal{C}$  generated by  $X$ . One puts  $\mathbf{G}(X) = \mathbf{G}(\langle X \rangle_{\otimes}, \omega)$ . It turns out that  $\mathbf{G}(X)$  is finite-dimensional and  $\mathbf{G}(\mathcal{C}, \omega) = \varprojlim \mathbf{G}(X)$ .

## Familiar Examples

Let  $\mathcal{C} = \text{Vec}_k$ , this is a neutral Tannakian category with the fibre functor being the identity. Then  $\mathbf{G}(\mathcal{C}, \omega) = \text{Spec } k$ .

Let  $\mathcal{C} = \text{VecGr}_k$ , the category of graded  $k$ -vector spaces  $(V_n)_n$  such that  $\bigoplus_n V_n$  is finite dimensional. This is a neutral Tannakian category with the fibre functor being the forgetful functor.

# Examples

## Tannakian Fundamental Group (contd.)

Often,  $\mathbf{G}(\mathcal{C}, \omega)$  is “too large” to handle directly. For  $X \in \mathcal{C}$ , let  $\langle X \rangle_{\otimes}$  be the Tannakian subcategory of  $\mathcal{C}$  generated by  $X$ . One puts  $\mathbf{G}(X) = \mathbf{G}(\langle X \rangle_{\otimes}, \omega)$ . It turns out that  $\mathbf{G}(X)$  is finite-dimensional and  $\mathbf{G}(\mathcal{C}, \omega) = \varprojlim \mathbf{G}(X)$ .

## Familiar Examples

Let  $\mathcal{C} = \text{Vec}_k$ , this is a neutral Tannakian category with the fibre functor being the identity. Then  $\mathbf{G}(\mathcal{C}, \omega) = \text{Spec } k$ .

Let  $\mathcal{C} = \text{VecGr}_k$ , the category of graded  $k$ -vector spaces  $(V_n)_n$  such that  $\bigoplus_n V_n$  is finite dimensional. This is a neutral Tannakian category with the fibre functor being the forgetful functor. Then  $\mathbf{G}(\mathcal{C}, \omega) = \mathbf{G}_m$ .

# Examples

## Tannakian Fundamental Group (contd.)

Often,  $\mathbf{G}(\mathcal{C}, \omega)$  is “too large” to handle directly. For  $X \in \mathcal{C}$ , let  $\langle X \rangle_{\otimes}$  be the Tannakian subcategory of  $\mathcal{C}$  generated by  $X$ . One puts  $\mathbf{G}(X) = \mathbf{G}(\langle X \rangle_{\otimes}, \omega)$ . It turns out that  $\mathbf{G}(X)$  is finite-dimensional and  $\mathbf{G}(\mathcal{C}, \omega) = \varprojlim \mathbf{G}(X)$ .

## Familiar Examples

Let  $\mathcal{C} = \text{Vec}_k$ , this is a neutral Tannakian category with the fibre functor being the identity. Then  $\mathbf{G}(\mathcal{C}, \omega) = \text{Spec } k$ .

Let  $\mathcal{C} = \text{VecGr}_k$ , the category of graded  $k$ -vector spaces  $(V_n)_n$  such that  $\bigoplus_n V_n$  is finite dimensional. This is a neutral Tannakian category with the fibre functor being the forgetful functor. Then  $\mathbf{G}(\mathcal{C}, \omega) = \mathbf{G}_m$ .

Let  $\mathcal{C} = \text{Hod}_{\mathbf{R}}$ , the category of real Hodge structures.

# Examples

## Tannakian Fundamental Group (contd.)

Often,  $\mathbf{G}(\mathcal{C}, \omega)$  is “too large” to handle directly. For  $X \in \mathcal{C}$ , let  $\langle X \rangle_{\otimes}$  be the Tannakian subcategory of  $\mathcal{C}$  generated by  $X$ . One puts  $\mathbf{G}(X) = \mathbf{G}(\langle X \rangle_{\otimes}, \omega)$ . It turns out that  $\mathbf{G}(X)$  is finite-dimensional and  $\mathbf{G}(\mathcal{C}, \omega) = \varprojlim \mathbf{G}(X)$ .

## Familiar Examples

Let  $\mathcal{C} = \text{Vec}_k$ , this is a neutral Tannakian category with the fibre functor being the identity. Then  $\mathbf{G}(\mathcal{C}, \omega) = \text{Spec } k$ .

Let  $\mathcal{C} = \text{VecGr}_k$ , the category of graded  $k$ -vector spaces  $(V_n)_n$  such that  $\bigoplus_n V_n$  is finite dimensional. This is a neutral Tannakian category with the fibre functor being the forgetful functor. Then  $\mathbf{G}(\mathcal{C}, \omega) = \mathbf{G}_m$ .

Let  $\mathcal{C} = \text{Hod}_{\mathbf{R}}$ , the category of real Hodge structures. Recall that a real Hodge structure is a finite dimensional real vector space  $V$  such that

$$V \otimes_{\mathbf{R}} \mathbf{C} \cong \bigoplus_{p,q} V^{p,q}, \quad \text{with} \quad V^{q,p} = \overline{V^{p,q}}.$$



## Tannakian Fundamental Group (contd.)

Often,  $\mathbf{G}(\mathcal{C}, \omega)$  is “too large” to handle directly. For  $X \in \mathcal{C}$ , let  $\langle X \rangle_{\otimes}$  be the Tannakian subcategory of  $\mathcal{C}$  generated by  $X$ . One puts  $\mathbf{G}(X) = \mathbf{G}(\langle X \rangle_{\otimes}, \omega)$ . It turns out that  $\mathbf{G}(X)$  is finite-dimensional and  $\mathbf{G}(\mathcal{C}, \omega) = \varprojlim \mathbf{G}(X)$ .

## Familiar Examples

Let  $\mathcal{C} = \text{Vec}_k$ , this is a neutral Tannakian category with the fibre functor being the identity. Then  $\mathbf{G}(\mathcal{C}, \omega) = \text{Spec } k$ .

Let  $\mathcal{C} = \text{VecGr}_k$ , the category of graded  $k$ -vector spaces  $(V_n)_n$  such that  $\bigoplus_n V_n$  is finite dimensional. This is a neutral Tannakian category with the fibre functor being the forgetful functor. Then  $\mathbf{G}(\mathcal{C}, \omega) = \mathbf{G}_m$ .

Let  $\mathcal{C} = \text{Hod}_{\mathbf{R}}$ , the category of real Hodge structures. Recall that a real Hodge structure is a finite dimensional real vector space  $V$  such that

$$V \otimes_{\mathbf{R}} \mathbf{C} \cong \bigoplus_{p,q} V^{p,q}, \quad \text{with} \quad V^{q,p} = \overline{V^{p,q}}.$$

This is a neutral Tannakian category with the fibre functor being the forgetful functor.

# Examples

## Tannakian Fundamental Group (contd.)

Often,  $\mathbf{G}(\mathcal{C}, \omega)$  is “too large” to handle directly. For  $X \in \mathcal{C}$ , let  $\langle X \rangle_{\otimes}$  be the Tannakian subcategory of  $\mathcal{C}$  generated by  $X$ . One puts  $\mathbf{G}(X) = \mathbf{G}(\langle X \rangle_{\otimes}, \omega)$ . It turns out that  $\mathbf{G}(X)$  is finite-dimensional and  $\mathbf{G}(\mathcal{C}, \omega) = \varprojlim \mathbf{G}(X)$ .

## Familiar Examples

Let  $\mathcal{C} = \text{Vec}_k$ , this is a neutral Tannakian category with the fibre functor being the identity. Then  $\mathbf{G}(\mathcal{C}, \omega) = \text{Spec } k$ .

Let  $\mathcal{C} = \text{VecGr}_k$ , the category of graded  $k$ -vector spaces  $(V_n)_n$  such that  $\bigoplus_n V_n$  is finite dimensional. This is a neutral Tannakian category with the fibre functor being the forgetful functor. Then  $\mathbf{G}(\mathcal{C}, \omega) = \mathbf{G}_m$ .

Let  $\mathcal{C} = \text{Hod}_{\mathbf{R}}$ , the category of real Hodge structures. Recall that a real Hodge structure is a finite dimensional real vector space  $V$  such that

$$V \otimes_{\mathbf{R}} \mathbf{C} \cong \bigoplus_{p,q} V^{p,q}, \quad \text{with} \quad V^{q,p} = \overline{V^{p,q}}.$$

This is a neutral Tannakian category with the fibre functor being the forgetful functor. Then  $\mathbf{G}(\mathcal{C}, \omega) = \mathbf{S} = \text{Res}_{\mathbf{C}/\mathbf{R}} \mathbf{G}_m$ .

## $\mathbb{Q}_p$ -unipotent fundamental group

Let  $X/\overline{\mathbb{Q}}$  be a smooth variety and  $b \in X(\overline{\mathbb{Q}})$  a geometric point.

## $\mathbf{Q}_p$ -unipotent fundamental group

Let  $X/\overline{\mathbf{Q}}$  be a smooth variety and  $b \in X(\overline{\mathbf{Q}})$  a geometric point. We consider the category of unipotent  $\mathbf{Q}_p$ -local systems in the étale topology  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$ .

## $\mathbf{Q}_p$ -unipotent fundamental group

Let  $X/\overline{\mathbf{Q}}$  be a smooth variety and  $b \in X(\overline{\mathbf{Q}})$  a geometric point. We consider the category of unipotent  $\mathbf{Q}_p$ -local systems in the étale topology  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$ . The objects are locally constant sheaves of finite-dimensional  $\mathbf{Q}_p$ -vector spaces on  $X$  considered in the étale topology

## $\mathbf{Q}_p$ -unipotent fundamental group

Let  $X/\overline{\mathbf{Q}}$  be a smooth variety and  $b \in X(\overline{\mathbf{Q}})$  a geometric point. We consider the category of unipotent  $\mathbf{Q}_p$ -local systems in the étale topology  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$ . The objects are locally constant sheaves of finite-dimensional  $\mathbf{Q}_p$ -vector spaces on  $X$  considered in the étale topology that admit a filtration

$$L = L^0 \supset L^1 \supset \cdots \supset L^n \supset L^{n+1} = 0$$

such that each quotient  $L_i/L_{i+1}$  is isomorphic to a direct sum of the constant sheaf  $\mathbf{Q}_p$ .

## $\mathbf{Q}_p$ -unipotent fundamental group

Let  $X/\overline{\mathbf{Q}}$  be a smooth variety and  $b \in X(\overline{\mathbf{Q}})$  a geometric point. We consider the category of unipotent  $\mathbf{Q}_p$ -local systems in the étale topology  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$ . The objects are locally constant sheaves of finite-dimensional  $\mathbf{Q}_p$ -vector spaces on  $X$  considered in the étale topology that admit a filtration

$$L = L^0 \supset L^1 \supset \cdots \supset L^n \supset L^{n+1} = 0$$

such that each quotient  $L_i/L_{i+1}$  is isomorphic to a direct sum of the constant sheaf  $\mathbf{Q}_p$ .  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$  is a neutral Tannakian category with fibre functor

$$F_b^{\mathrm{un}} : \mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p) \rightarrow \mathrm{Vecf}_{\mathbf{Q}_p}, L \mapsto L_b$$

that associates to each sheaf its stalk at  $b$ .

## $\mathbf{Q}_p$ -unipotent fundamental group

Let  $X/\overline{\mathbf{Q}}$  be a smooth variety and  $b \in X(\overline{\mathbf{Q}})$  a geometric point. We consider the category of unipotent  $\mathbf{Q}_p$ -local systems in the étale topology  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$ . The objects are locally constant sheaves of finite-dimensional  $\mathbf{Q}_p$ -vector spaces on  $X$  considered in the étale topology that admit a filtration

$$L = L^0 \supset L^1 \supset \cdots \supset L^n \supset L^{n+1} = 0$$

such that each quotient  $L_i/L_{i+1}$  is isomorphic to a direct sum of the constant sheaf  $\mathbf{Q}_p$ .  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$  is a neutral Tannakian category with fibre functor

$$F_b^{\mathrm{un}} : \mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p) \rightarrow \mathrm{Vecf}_{\mathbf{Q}_p}, L \mapsto L_b$$

that associates to each sheaf its stalk at  $b$ .

The  $\mathbf{Q}_p$ -unipotent fundamental group is  $\pi_1^{\mathrm{un}, \mathbf{Q}_p}(X, b) := \mathbf{G}(\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p), F_b^{\mathrm{un}})$ .



## $\mathbf{Q}_p$ -unipotent fundamental group

Let  $X/\overline{\mathbf{Q}}$  be a smooth variety and  $b \in X(\overline{\mathbf{Q}})$  a geometric point. We consider the category of unipotent  $\mathbf{Q}_p$ -local systems in the étale topology  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$ . The objects are locally constant sheaves of finite-dimensional  $\mathbf{Q}_p$ -vector spaces on  $X$  considered in the étale topology that admit a filtration

$$L = L^0 \supset L^1 \supset \cdots \supset L^n \supset L^{n+1} = 0$$

such that each quotient  $L_i/L_{i+1}$  is isomorphic to a direct sum of the constant sheaf  $\mathbf{Q}_p$ .  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$  is a neutral Tannakian category with fibre functor

$$F_b^{\mathrm{un}} : \mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p) \rightarrow \mathrm{Vecf}_{\mathbf{Q}_p}, L \mapsto L_b$$

that associates to each sheaf its stalk at  $b$ .

The  $\mathbf{Q}_p$ -unipotent fundamental group is  $\pi_1^{\mathrm{un}, \mathbf{Q}_p}(X, b) := \mathbf{G}(\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p), F_b^{\mathrm{un}})$ .

It corresponds to the  $\mathbf{Q}_p$ -unipotent completion of the (geometric) étale fundamental group of  $X$ .

## $\mathbf{Q}_p$ -unipotent fundamental group

Let  $X/\overline{\mathbf{Q}}$  be a smooth variety and  $b \in X(\overline{\mathbf{Q}})$  a geometric point. We consider the category of unipotent  $\mathbf{Q}_p$ -local systems in the étale topology  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$ . The objects are locally constant sheaves of finite-dimensional  $\mathbf{Q}_p$ -vector spaces on  $X$  considered in the étale topology that admit a filtration

$$L = L^0 \supset L^1 \supset \dots \supset L^n \supset L^{n+1} = 0$$

such that each quotient  $L_i/L_{i+1}$  is isomorphic to a direct sum of the constant sheaf  $\mathbf{Q}_p$ .  $\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p)$  is a neutral Tannakian category with fibre functor

$$F_b^{\mathrm{un}} : \mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p) \rightarrow \mathrm{Vecf}_{\mathbf{Q}_p}, L \mapsto L_b$$

that associates to each sheaf its stalk at  $b$ .

The  $\mathbf{Q}_p$ -unipotent fundamental group is  $\pi_1^{\mathrm{un}, \mathbf{Q}_p}(X, b) := \mathbf{G}(\mathrm{Un}^{\mathrm{ét}}(X, \mathbf{Q}_p), F_b^{\mathrm{un}})$ .

It corresponds to the  $\mathbf{Q}_p$ -unipotent completion of the (geometric) étale fundamental group of  $X$ .

Here, we also have (torsor of) paths

$$\pi_1^{\mathrm{un}, \mathbf{Q}_p}(X; b, x) := \mathbf{Isom}^{\otimes}(F_b^{\mathrm{un}}, F_x^{\mathrm{un}})$$

# Examples

## de Rham fundamental group

Let  $X/K$  be a smooth variety where  $K$  is of characteristic 0.

## de Rham fundamental group

Let  $X/K$  be a smooth variety where  $K$  is of characteristic 0. We consider the category  $\mathrm{Un}(X)$  of unipotent (algebraic) vector bundles on  $X$ .

## de Rham fundamental group

Let  $X/K$  be a smooth variety where  $K$  is of characteristic 0. We consider the category  $\text{Un}(X)$  of unipotent (algebraic) vector bundles on  $X$ . The objects are vector bundles  $\mathcal{V}$  on  $X$  equipped with a integrable (flat) connection

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/K} \otimes \mathcal{V}$$

## de Rham fundamental group

Let  $X/K$  be a smooth variety where  $K$  is of characteristic 0. We consider the category  $\text{Un}(X)$  of unipotent (algebraic) vector bundles on  $X$ . The objects are vector bundles  $\mathcal{V}$  on  $X$  equipped with a integrable (flat) connection

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/K} \otimes \mathcal{V}$$

(that is, curvature  $\nabla_1 \circ \nabla = 0$ , where  $\nabla_1 : \Omega_{X/K} \otimes \mathcal{V} \rightarrow \Omega_{X/K}^2 \otimes \mathcal{V}$ )

## de Rham fundamental group

Let  $X/K$  be a smooth variety where  $K$  is of characteristic 0. We consider the category  $\text{Un}(X)$  of unipotent (algebraic) vector bundles on  $X$ . The objects are vector bundles  $\mathcal{V}$  on  $X$  equipped with a integrable (flat) connection

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/K} \otimes \mathcal{V}$$

(that is, curvature  $\nabla_1 \circ \nabla = 0$ , where  $\nabla_1 : \Omega_{X/K} \otimes \mathcal{V} \rightarrow \Omega_{X/K}^2 \otimes \mathcal{V}$ ) that admits a filtration

$$\mathcal{V} = \mathcal{V}_0 \supset \mathcal{V}_1 \supset \cdots \supset \mathcal{V}_n \supset \mathcal{V}_{n+1} = 0$$

by subbundles stable under  $\nabla$  such that each  $\mathcal{V}_i/\mathcal{V}_{i+1}$  is a trivial bundle with connection.

## de Rham fundamental group

Let  $X/K$  be a smooth variety where  $K$  is of characteristic 0. We consider the category  $\text{Un}(X)$  of unipotent (algebraic) vector bundles on  $X$ . The objects are vector bundles  $\mathcal{V}$  on  $X$  equipped with a integrable (flat) connection

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/K} \otimes \mathcal{V}$$

(that is, curvature  $\nabla_1 \circ \nabla = 0$ , where  $\nabla_1 : \Omega_{X/K} \otimes \mathcal{V} \rightarrow \Omega_{X/K}^2 \otimes \mathcal{V}$ ) that admits a filtration

$$\mathcal{V} = \mathcal{V}_0 \supset \mathcal{V}_1 \supset \cdots \supset \mathcal{V}_n \supset \mathcal{V}_{n+1} = 0$$

by subbundles stable under  $\nabla$  such that each  $\mathcal{V}_i/\mathcal{V}_{i+1}$  is a trivial bundle with connection. Morphisms are maps of sheaves preserving the connection.



## de Rham fundamental group

Let  $X/K$  be a smooth variety where  $K$  is of characteristic 0. We consider the category  $\mathrm{Un}(X)$  of unipotent (algebraic) vector bundles on  $X$ . The objects are vector bundles  $\mathcal{V}$  on  $X$  equipped with a integrable (flat) connection

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/K} \otimes \mathcal{V}$$

(that is, curvature  $\nabla_1 \circ \nabla = 0$ , where  $\nabla_1 : \Omega_{X/K} \otimes \mathcal{V} \rightarrow \Omega_{X/K}^2 \otimes \mathcal{V}$ ) that admits a filtration

$$\mathcal{V} = \mathcal{V}_0 \supset \mathcal{V}_1 \supset \cdots \supset \mathcal{V}_n \supset \mathcal{V}_{n+1} = 0$$

by subbundles stable under  $\nabla$  such that each  $\mathcal{V}_i/\mathcal{V}_{i+1}$  is a trivial bundle with connection. Morphisms are maps of sheaves preserving the connection.  $\mathrm{Un}(X)$  is a neutral Tannakian category with fibre functor

$$F_b^{\mathrm{dR}} : \mathrm{Un}(X) \rightarrow \mathrm{Vecf}_K, (\mathcal{V}, \nabla) \mapsto \mathcal{V}_b$$

that associates to each vector bundle its stalk at  $b$ .

## de Rham fundamental group

Let  $X/K$  be a smooth variety where  $K$  is of characteristic 0. We consider the category  $\mathrm{Un}(X)$  of unipotent (algebraic) vector bundles on  $X$ . The objects are vector bundles  $\mathcal{V}$  on  $X$  equipped with a integrable (flat) connection

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/K} \otimes \mathcal{V}$$

(that is, curvature  $\nabla_1 \circ \nabla = 0$ , where  $\nabla_1 : \Omega_{X/K} \otimes \mathcal{V} \rightarrow \Omega_{X/K}^2 \otimes \mathcal{V}$ ) that admits a filtration

$$\mathcal{V} = \mathcal{V}_0 \supset \mathcal{V}_1 \supset \cdots \supset \mathcal{V}_n \supset \mathcal{V}_{n+1} = 0$$

by subbundles stable under  $\nabla$  such that each  $\mathcal{V}_i/\mathcal{V}_{i+1}$  is a trivial bundle with connection. Morphisms are maps of sheaves preserving the connection.  $\mathrm{Un}(X)$  is a neutral Tannakian category with fibre functor

$$F_b^{\mathrm{dR}} : \mathrm{Un}(X) \rightarrow \mathrm{Vecf}_K, (\mathcal{V}, \nabla) \mapsto \mathcal{V}_b$$

that associates to each vector bundle its stalk at  $b$ .

The *de Rham fundamental group* is  $\pi_1^{\mathrm{dR}}(X, b) := \mathbf{G}(\mathrm{Un}(X), F_b^{\mathrm{dR}})$ .

## de Rham fundamental group

Let  $X/K$  be a smooth variety where  $K$  is of characteristic 0. We consider the category  $\text{Un}(X)$  of unipotent (algebraic) vector bundles on  $X$ . The objects are vector bundles  $\mathcal{V}$  on  $X$  equipped with a integrable (flat) connection

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/K} \otimes \mathcal{V}$$

(that is, curvature  $\nabla_1 \circ \nabla = 0$ , where  $\nabla_1 : \Omega_{X/K} \otimes \mathcal{V} \rightarrow \Omega_{X/K}^2 \otimes \mathcal{V}$ ) that admits a filtration

$$\mathcal{V} = \mathcal{V}_0 \supset \mathcal{V}_1 \supset \cdots \supset \mathcal{V}_n \supset \mathcal{V}_{n+1} = 0$$

by subbundles stable under  $\nabla$  such that each  $\mathcal{V}_i/\mathcal{V}_{i+1}$  is a trivial bundle with connection. Morphisms are maps of sheaves preserving the connection.  $\text{Un}(X)$  is a neutral Tannakian category with fibre functor

$$F_b^{\text{dR}} : \text{Un}(X) \rightarrow \text{Vecf}_K, (\mathcal{V}, \nabla) \mapsto \mathcal{V}_b$$

that associates to each vector bundle its stalk at  $b$ .

The *de Rham fundamental group* is  $\pi_1^{\text{dR}}(X, b) := \mathbf{G}(\text{Un}(X), F_b^{\text{dR}})$ .

Here, we also have the notion of (torsor of) paths

$$\pi_1^{\text{dR}}(X; b, x) := \mathbf{Isom}^{\otimes}(F_b^{\text{dR}}, F_x^{\text{dR}})$$

Fin.

- [1] Szamuely, Tamás (2009). Galois Groups and Fundamental Groups. *Cambridge University Press*.
- [2] Kim, M. (2010). Fundamental groups and Diophantine geometry. *Open Mathematics*, 8(4), 633-645.
- [3] Sujatha, R. (2008), Motives from a categorical point of view. *Autour des motifs*, Asian-French summer school on algebraic geometry and number theory, Vol. I, Panor. Synthèses 29, Soc. Math. France, Paris 2009, 27–48.