

M382D NOTES: DIFFERENTIAL TOPOLOGY

ARUN DEBRAY
JANUARY 25, 2016

These notes were taken in UT Austin's Math 382D (Differential Topology) class in Spring 2016, taught by Lorenzo Sadun. I live-T_EXed them using vim, and as such there may be typos; please send questions, comments, complaints, and corrections to a.debray@math.utexas.edu. Thanks to Thérèse Wu for finding a few typos.

CONTENTS

1. The Inverse and Implicit Function Theorems: 1/20/15	1
2. The Contraction Mapping Theorem: 1/22/15	3
3. Manifolds: 1/25/15	5

Lecture 1.

The Inverse and Implicit Function Theorems: 1/20/15

“The most important lesson of the start of this class is the proper pronunciation of my name [Sadun]: it rhymes with ‘balloon.’”

We’re basically going to march through the textbook (Guillemin and Pollack), with a little more in the beginning and a little more in the end; however, we’re going to be a bit more abstract, talking about manifolds more abstractly, rather than just embedding them in \mathbb{R}^n , though the theorems are mostly the same. At the beginning, we’ll discuss the analytic underpinnings to differential topology in more detail, and at the end, we’ll hopefully have time to discuss de Rham cohomology.

Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Its derivative is df ; what exactly is this? There are several possible answers.

- It’s the best linear approximation to f at a given point.
- It’s the matrix of partial derivatives.

What we need to do is make good, rigorous sense of this, more so than in multivariable calculus, and relate the two notions.

Definition. A function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is *differentiable* at an $a \in \mathbb{R}^n$ if there exists a linear map $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that

$$\lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - L(h)|}{|h|} = 0. \quad (1.1)$$

In this case, L is called the *differential* of f at a , written $df|_a$.

Note that $h \in \mathbb{R}^n$ and the numerator is in \mathbb{R}^m , so it’s quite important to have the magnitudes there, or else it would make no sense.

Another way to rewrite this is that $f(a+h) = f(a) + L(h) + o(\text{small})$, i.e. along with some small error (whatever that means). This makes sense of the first notion: L is a linear approximation to f near a . Now, let’s make sense of the second notion.

Theorem 1.1. If f is differentiable at a , then df is given by the matrix $\left(\frac{\partial f^i}{\partial x^j}\right)$.

Proof. The idea: if f is differentiable at a , then (1.1) holds for $h \rightarrow 0$ along any path!

So let’s take e_j be a unit vector and $h = te_j$ as $t \rightarrow 0$ in \mathbb{R} . Then, (1.1) reduces to

$$L(te_j) = \frac{f(a_1, a_2, \dots, a_j + t, a_{j+1}, \dots, a_n) - f(a)}{t},$$

and as $t \rightarrow 0$, this shows $L(\mathbf{e}_j)^i = \frac{\partial f^i}{\partial x^j}$. \square

In particular, if f is differentiable, then all partial derivatives exist. The converse is *false*: there exist functions whose partial derivatives exist at a point a , but are not differentiable. In fact, one can construct a function whose directional derivatives all exist, but is not differentiable! There will be an example on the first homework. The idea is that directional derivatives record linear paths, but differentiability requires all paths, and so making things fail along, say, a quadratic, will produce these strange counterexamples.

Nonetheless, if all partial derivatives exist, then we're almost there.

Theorem 1.2. *Suppose all partial derivatives of f exist at a and are continuous on a neighborhood of a , then f is differentiable at a .*

In calculus, one can formulate several “guiding” ideas, e.g. the whole change is the sum of the individual changes, the whole is the (possibly infinite) sum of the parts, and so forth. One particular one is: *one variable at a time*. This principle will guide the proof of this theorem.

Proof. The proof will be given for $m = 2$ and $n = 1$, but you can figure out the small details needed to generalize it; for larger n , just repeat the argument for each component.

We want to compute

$$\begin{aligned} f(a_1 + h_1, a_2 + h_2) - f(a_1, a_2) \\ = f(a_1 + h_1, a_2 + h_2) - f(a_1 + h_1, a_2) + f(a_1 + h_1, a_2) - f(a_1, a_2) \end{aligned}$$

Regrouping, this is two single-variable questions. In particular, we can apply the mean value theorem: there exist $c_1, c_2 \in \mathbb{R}$ such that

$$\begin{aligned} &= \frac{\partial f}{\partial x^2} \Big|_{(a_1+h_1, a_2+c_2)} h_2 + \frac{\partial f}{\partial x^1} \Big|_{(a_1+c_1, a_2)} h_1 \\ &= \left(\frac{\partial f}{\partial x^1} \Big|_{a_1+c_1, a_2} - \frac{\partial f}{\partial x^1} \Big|_a \right) h_1 + \left(\frac{\partial f}{\partial x^2} \Big|_{a_1+h_1, a_2+c_2} - \frac{\partial f}{\partial x^2} \Big|_a \right) h_2 + \left(\frac{\partial f}{\partial x^1} \Big|_a, \frac{\partial f}{\partial x^2} \Big|_a \right) \begin{pmatrix} h_1 \\ h_2 \end{pmatrix}, \end{aligned}$$

but since the partials are continuous, the left two terms go to 0, and since the last term is linear, it goes to 0 as $h \rightarrow 0$. \square

We'll often talk about *smooth* functions in this class, which are functions for which all higher-order derivatives exist and are continuous. Thus, they don't have the problems that one counterexample had.

Since we're going to be making linear approximations to maps, then we should discuss what happens when you perturb linear maps a little bit. Recall that if $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is linear, then its image $\text{Im}(L) \subset \mathbb{R}^m$ and its kernel $\ker(L) \subset \mathbb{R}^n$.

Suppose $n \leq m$; then, L is said to have *full rank* if $\text{rank } L = n$. This is an open condition: every full-rank linear function can be perturbed a little bit and stay linear. This will be very useful: if a (possibly nonlinear) function's differential has full rank, then one can say some interesting things about it.

If $n \geq m$, then full rank means rank m . This is once again stable (an open condition): such a linear map can be written $L = (A \mid B)$, where A is an invertible $m \times m$ matrix, and invertibility is an open condition (since it's given by the determinant, which is a continuous function).

To actually figure out whether a linear map has full rank, write down its matrix and row-reduce it, using Gaussian elimination. Then, you can read off a basis for the kernel, determining the free variables and the relations determining the other variables. In general, for a k -dimensional subspace of \mathbb{R}^n , you can pick k variables arbitrarily and these force the remaining $n - k$ variables. The point is: *the subspace is the graph of a function*.

Now, we can apply this to more general smooth functions.

Theorem 1.3. *Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is smooth and $df|_a$ has full rank.*

- (1) (Inverse function theorem) *If $n = m$, then there is a neighborhood U of a such that $f|_U$ is invertible, with a smooth inverse.*
- (2) (Implicit function theorem) *In general, there is a neighborhood U of a such that $U \cap f^{-1}(f(a))$ is the graph of some smooth function $g : \mathbb{R}^{n-m} \rightarrow \mathbb{R}^m$ (up to permutation of indices).*
- (3) (Immersion theorem) *If $n < m$, there's a neighborhood U of a such that $f(U)$ is the graph of a smooth $g : \mathbb{R}^n \rightarrow \mathbb{R}^{n-m}$.*

This time, the results are local rather than global, but once again, full rank means (local) invertibility when $m = n$, and more generally means that we can write all the points sent to $f(a)$ (analogous to a kernel) as the graph of a smooth function.

It's possible to sharpen these theorems slightly: instead of maximal rank, you can use that if $df|_a$ has block form with the square block invertible, then similar statements hold.

The content of these theorems, the way to think of them, is that in these cases, smooth functions locally behave like linear ones. But this is not too much of a surprise: differentiability means exactly that a function can be locally well approximated by a linear function. The point of the proof is that the higher-order terms also vanish.

For example, if $m = n = 1$, then full rank means the derivative is nonzero at a . In this case, it's increasing or decreasing in a neighborhood of a , and therefore invertible. On the other hand, if the derivative is 0, then bad things happen, because it's controlled by the higher-order derivatives, so one can have a noninvertible function (e.g. a constant) or an invertible function whose inverse isn't smooth (e.g. $y = x^3$ at $x = 0$).

This is not the last time in this class that maximal rank implies nice analytic results.

We're going to prove (2); then, as linear-algebraic corollaries, we'll recover the other two.

Lecture 2.

The Contraction Mapping Theorem: 1/22/15

Today, we're going to prove the generalized inverse function theorem, Theorem 1.3. We'll start with the case where $m = n$, which is also the simplest in the linear case (full rank means invertible, almost tautologically).

Theorem 2.1. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be smooth. if $df|_a$ is invertible, then*

- (1) f is invertible on a neighborhood of a ,
- (2) f^{-1} is smooth on a neighborhood of a , and
- (3) $d(f^{-1})|_{f(a)} = (df|_a)^{-1}$.

Proof of part (1). Without loss of generality, we can assume that $a = f(a) = 0$ by translating. We can also assume that $df|_a = I$, by precomposing with $df|_a^{-1}$:

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{f} & \mathbb{R}^n \\ \uparrow df|_a^{-1} & \nearrow & \\ \mathbb{R}^n & & \end{array}$$

If we prove the result for the diagonal arrow, then it is also true for f . Since the domain and codomain of f are different in this proof, we're going to call the former X and the latter Y , so $f : X \rightarrow Y$.

Now, since f is smooth, its derivative is continuous, so there's a neighborhood of a in X given by the x such that $\|df|_x - I\| < 1/2$.¹ And by shrinking this neighborhood, we can assume that it is a closed ball C .

On C , f is injective: if $x_1, x_2 \in C$, then since C is convex, then there's a line $\gamma(t) = x_1 + tv$ (where $v = x_2 - x_1$) joining x_1 to x_2 , and $\frac{df}{dt} = (df|_{\gamma(t)})v$. Therefore

$$\begin{aligned} f(x_2) - f(x_1) &= \left(\int_0^1 df|_{\gamma(t)} dt \right) v \\ &= \int_0^1 ((df|_{\gamma(t)} - I) + I) dt \\ &= x_2 - x_1 + \int_0^1 (df|_{\gamma(t)} - I)v dt. \end{aligned}$$

We can bound the integral below:

$$\left| \int_0^1 (df|_{\gamma(t)} - I)v dt \right| \leq \int_0^1 |(df|_{\gamma(t)} - I)v| dt \leq \int_0^1 \frac{1}{2}|v| dt = \frac{|v|}{2}.$$

¹There are many different norms on the space of $n \times n$ matrices, but since this is a finite-dimensional vector space, they are all equivalent. However, for this proof we're going to take the operator norm $\|A\| = \sup_{v \in \mathbb{R}^{n-1}} |Av|$.

Thus, since $x_2 - x_1 = v$, then their sum has magnitude at least $v/2$, so in particular it can't be zero. Thus, f is injective on C . The point is, since df is close to the identity on C , we get an error term that we can make small.

To construct an inverse, we need to make it surjective on a neighborhood of $f(a)$ in Y . The way to do this is called the contraction mapping principle, but we'll do it by hand for now and recover the general principle later.

To be precise, we'll iterate with a "poor-man's Newton's method:" if $y \in Y$, then given x_n , let $x_{n+1} = x_0 - (f(x_0) - y) = y + x_0 - f(x_0)$ (since we're using the derivative at the origin instead of at x , and this is just the identity). A fixed point of this iteration is a preimage of y .

Since

$$x_{n+1} - x_n = y + x_n - f(x_n) - (y + x_{n-1} - f(x_{n-1})) = (x_n - x_{n-1}) - (f(x_n) - f(x_{n-1})),$$

then $|x_{n+1} - x_n| < (1/2)|x_n - x_{n-1}|$, so in particular, this is a Cauchy sequence! Thus, it must converge, and to a value with magnitude no more than $2|y|$. Thus, if C has radius R , then for any y in the ball of radius $1/2$ from the origin (in Y), y has a preimage x , so f is surjective on this neighborhood. \square

Now, we can discuss the contraction mapping principle more generally.

Definition. Let X be a complete metric space and $T : X \rightarrow X$ be a continuous map such that $d(T(x), T(y)) \leq cd(x, y)$ for all $x, y \in X$ and some $c \in [0, 1)$. Then, T is called a *contraction mapping*.

Theorem 2.2 (Contraction mapping principle). *If X is a complete metric space and T a contraction mapping on X , then there's a unique fixed point x (i.e. $T(x) = x$).*

Proof. Uniqueness is pretty simple: if T has two fixed points x and x' such that $x \neq x'$, then $d(T(x), T(x')) \leq cd(x, x') = d(T(x), T(x'))$, and $c < 1$, so this is a contradiction, so $x = x'$.

Existence is basically the proof we just saw: pick an arbitrary $x_0 \in X$ and let $x_{n+1} = T(x_n)$. Then, $d(x_m, x_n) \leq c^{|n-m|}d(x_n, x_{n-1})$, so this sequence is Cauchy, and has a limit x . Then, since T is continuous, $T(x) = x$. \square

Now, back to the theorem.

Proof of Theorem 2.1, part (2). Once again, we assume $f(0) = 0$. By the fundamental theorem of calculus, on our neighborhood of 0,

$$y = f(x) = \int_0^1 df|_{tx}(x) dt.$$

Since we assumed that $df|_0 = I$, then in this neighborhood, this integral is small in x , so $y = x + o(|x|)$, and in particular $y = x + o(|y|)$, or rearranging, $x = f^{-1}(y) = y + o(|y|)$. Thus, f^{-1} is differentiable once at 0. Moreover, this is true in a neighborhood of $0 \in Y$, because we can repeat this argument for sufficiently close $y \in Y$.

To see why this is smooth, let $A(t)$ be an invertible function, so that $A(t)A(t)^{-1} = I$. Thus, $A'A^{-1} = A(A^{-1})' = 0$, so rearranging, $(A^{-1})' = -A^{-1}A'A^{-1}$. Thus, you can differentiate the inverse as many times as you can differentiate the original function, so f^{-1} is smooth, and (after some playing about with the chain rule) $d(f^{-1})|_a = (df|_a)^{-1}$. \square

We can use this to recover the rest of Theorem 1.3 as corollaries.

Proof of Theorem 1.3, part (2). First, for the implicit function theorem, let $n > m$ and $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be smooth with full rank, and choose a basis in which $df|_a = (A | B)$ in block form, where A is an invertible $m \times m$ matrix. The theorem statement is that we can write the first m coordinates as a function of the last $n - m$ coordinates: specifically, that there exists a neighborhood U of a such that $U = f^{-1}(f(a)) = U \cap \{g(y), y\}$ for some smooth $g : \mathbb{R}^{n-m} \rightarrow \mathbb{R}^m$.²

Now, the proof. Let $x \in \mathbb{R}^m$ and $y \in \mathbb{R}^{n-m}$, and let

$$F \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} f(x, y) \\ y \end{pmatrix}.$$

Hence,

$$dF|_a = \begin{pmatrix} A & B \\ 0 & I \end{pmatrix}.$$

This is invertible, since A is (its determinant is $\det A \neq 0$). Thus, we apply the inverse function theorem to F to conclude that a smooth F^{-1} exists, and so if π_1 denotes projection onto the first component, $x = \pi_1 \circ F^{-1}(0, y) = g(y)$. \square

²For example, if $n = 2$ and $m = 1$, consider $f(x) = |x|^2 - 1$, and $a = (\cos \theta, \sin \theta)$. Then, $f^{-1}(f(a))$ is the unit circle, so the implicit function is telling us that locally, the circle is a function of x_1 in terms of x_2 , or vice versa.

Lecture 3.

Manifolds: 1/25/15*“Erase any notes you have of the last eight minutes! But the first 40 minutes were okay.”*

Recall that we’ve been discussing Theorem 1.3, a collection of results called the inverse function theorem, the implicit function theorem, and the immersion theorem. These are local (not global) results, and generalize similar results for linear maps: not all matrices are square, but if a matrix has full rank, it can be written in two blocks, one of which is invertible. Using this with $df|_a$ as our matrix is the idea behind proving Theorem 1.3: the first n variables determine the remaining $(m - n)$ variables (for some values of m and n).

However, we don’t know which variables they are: you may have to permute x_1, \dots, x_n to get the last $m - n$ variables as smooth functions of the first ones. For example, for a circle, the tangent line is horizontal sometimes (so we can’t always parameterize in terms of x_2) and vertical at other times (so we can’t only use x_1).

Before we prove the immersion theorem (part (3) of Theorem 1.3), let’s recall what tools we use to talk about curves in the plane.

- (1) A common technique is using a *parameterized curve*, the image of a smooth $\gamma(t) : \mathbb{R} \rightarrow \mathbb{R}^2$ whose derivative is never zero (to avoid singularities). For example, $f(t) = t^2, t^3$ has a zero at the origin, but the curve one obtains is $y = \pm x^{3/2}$, which has a cusp at $(0, 0)$. This is the content of the immersion theorem.
- (2) Another way to describe curves is as level sets: $f(x, y) = c$, most famously the circle. This is the content of the implicit function theorem: this looks like a graph-like curve locally.
- (3) This brings us to the most simple method: graphs of functions, just like in calculus.

And the point of Theorem 1.3 is that these three approaches give you the same sets, *up to permutation of variables* (and that a curve is the graph of a function only locally). We have these three pictures of what higher-dimensional surfaces look like.

And that means that when we talk about manifolds, which are the analogue of higher-dimensional surfaces, we should keep these things in mind: a manifold may be defined abstractly, but we understand manifolds through these three visualizations.

Proof of Theorem 1.3, part (3). We’re going to prove the equivalent statement that if the first n rows of $df|_a$ are linearly independent, then the remaining $m - n$ variables are smooth functions in the first n .

Since $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, then let π_1 denote projection onto the first n coordinates, so we have a commutative diagram

$$\begin{array}{ccc} \mathbb{R}^n & \xrightarrow{f} & \mathbb{R}^m \\ & \searrow \pi_1 \circ f & \downarrow \pi_1 \\ & & \mathbb{R}^n. \end{array}$$

In block form, $df|_a = \begin{pmatrix} A \\ B \end{pmatrix}$, where A is invertible, and therefore $d(\pi_1 \circ f)|_a = A$. This is invertible, so $(\pi_1 \circ f)^{-1}$ has an inverse in a neighborhood of a , by the inverse function theorem. Thus, if π_2 denotes projection onto the last $m - n$ coordinates, then $g = \pi_2 \circ f \circ (\pi_1 \circ f)^{-1}$ writes the last $m - n$ coordinates in terms of the first n , as desired. \square

Now, we’re ready to talk about smooth manifolds.

Definition. A k -manifold X in \mathbb{R}^n is a set that locally looks like one of the descriptions (1), (2), or (3) for a smooth surface. That is, it satisfies one of the following descriptions.

- (1) For every $p \in X$, there’s a neighborhood U of p where one can write $N - k$ variables in smooth functions of the remaining k variables, i.e. there is a neighborhood $V \subset \mathbb{R}^k$ and a smooth $g : V \rightarrow \mathbb{R}^{N-k}$ such that $X \cap U = \{(x, g(x))\}$ (up to permutation).
- (2) X is locally the image of a smooth map, i.e. for every $p \in X$, there’s a neighborhood U of p and a smooth $f : \mathbb{R}^k \rightarrow \mathbb{R}^N$ with full rank such that the image of f in U is $X \cap U$. This is the “parameterized curve” analogue.
- (3) Locally, X is the level set of a smooth map $f : \mathbb{R}^N \rightarrow \mathbb{R}^{N-k}$ with full rank.

If k is understood from context (or not important), X will also be called a *manifold*.

The big theorem is that these three conditions are equivalent, and this follows directly from Theorem 1.3.

For example, suppose we have the graph of a smooth function $y = x^2$. How can we write this as the image of a smooth map? Well, $(x, y) = (t, t^2)$ has nonzero derivative, and we can do exactly the same thing (locally) for a manifold in general. And this is the level set $f(x, y) = 0$, where $f(x, y) = y - x^2$, and the same thing works (locally) for manifolds: for a general graph $\mathbf{y} = g(\mathbf{x})$, this is the level set of $f(\mathbf{x}, \mathbf{y}) = \mathbf{y} - g(\mathbf{x})$, whose derivative is $df = (I \mid -dg^T)$, which has full rank. Neat.

And perhaps most useful for now, something that's locally a graph is really easy to visualize: it's the bedrock on which one first defined curves and surfaces.

Now, that's a manifold in \mathbb{R}^n . As far as Guillemin and Pollack are concerned, that's the only kind of manifold there is, but we want to talk about abstract manifolds, but that means we'll need one more important property.

Suppose $X \subset \mathbb{R}^N$ is a manifold, and $p \in X$. We're going to look at a neighborhood of p as the image of a smooth $g_1 : \mathbb{R}^k \rightarrow \mathbb{R}^N$; this is the most common and most fundamental description of a manifold. However, this is not in general unique; suppose $g_2 : \mathbb{R}^k \rightarrow \mathbb{R}^N$ lands in a different neighborhood of p — though, by restricting to their intersection, we can assume we have two smooth maps (sometimes called *charts*) into the same neighborhood, and they both have inverses, so we have a well-defined function $g_2^{-1} \circ g_1 : \mathbb{R}^k \rightarrow \mathbb{R}^k$. Is it smooth?

Theorem 3.1. $g_2^{-1} \circ g_1$ is smooth.

The key assumption here is that dg_1 and dg_2 both have maximal rank.

Definition. The *tangent space* to X at p , denoted $T_p X$, is $\text{Im}(dg_1|_{g_1^{-1}(p)})$; it is a k -dimensional subspace of \mathbb{R}^N .

This is the set of velocity vectors of paths through p , which makes sense, because such a path must come from a path downstairs in \mathbb{R}^k , since g_1 is locally invertible.

Lemma 3.2. The tangent space is independent of choice of g_1 .

The idea is that any velocity vector must come from a path in both $\text{Im}(dg_1|_{g_1^{-1}(p)})$ and $\text{Im}(dg_2|_{g_2^{-1}(p)})$, so these two images are the same.

Then, we'll punt the proof of Theorem 3.1 to next lecture.