

SUMMER 2016 ALGEBRAIC GEOMETRY SEMINAR

ARUN DEBRAY

MAY 23, 2016

CONTENTS

1. Separability, Varieties and Rational Maps: 5/16/16	1
2. Proper Morphisms: 5/19/16	5
3. Dimension: 5/23/16	9

1. SEPARABILITY, VARIETIES AND RATIONAL MAPS: 5/16/16

Today's lecture was given by Tom Oldfield, on the first half of chapter 10.

This seminar has a website, located at

<https://www.ma.utexas.edu/users/toldfield/Seminars/Algebraicgeometryreading.html>.

The first half of Chapter 10 is about separated morphisms and varieties; it only took us 10 chapters! Vakil writes that he was very conflicted about leaving a proper treatment of algebraic varieties, a cornerstone of classical algebraic geometry, to so late in the notes. But from a modern perspective, our hands are tied: varieties are defined in terms of properties, which means building those properties out of other properties and out of the large amount of technology you need for modern algebraic geometry. With that technology out of the way, here we are.

One of these properties is separability. Let $\pi : X \rightarrow Y$ be a morphism of schemes; then, the **diagonal** is the induced morphism $\delta_\pi : X \rightarrow X \times_Y X$ defined by $x \mapsto (x, x)$; this maps into the fiber product because it fits into the diagram

$$\begin{array}{ccccc}
 X & & & & \\
 \delta_\pi \searrow & & 1_X \searrow & & \\
 & X \times_Y X & \xrightarrow{p_2} & X & \\
 1_X \searrow & \downarrow p_1 & & \downarrow \pi & \\
 & X & \xrightarrow{\pi} & Y &
 \end{array} \tag{1.1}$$

Here, p_1 and p_2 are the projections onto the first and second components, respectively, and 1_X is the identity map on X .

The diagonal has a few nice properties. Suppose $V \subset Y$ is open, and $U, U' \subset \pi^{-1}(V)$ are open subsets of X . Then, $U \times_V U' = p_1^{-1}(U) \cap p_2^{-1}(U')$: we constructed fiber products such that they send open embeddings to intersections. In particular, if $U \cong \text{Spec } A$, $U' \cong \text{Spec } A'$, and $V \cong \text{Spec } B$ are affine, $U \times_V U' \cong \text{Spec}(A \otimes_B A')$. Therefore $\delta_\pi^{-1}(U \times_V U') = \delta_\pi^{-1}(p_1^{-1}(U) \cap p_2^{-1}(U')) = U \cap U'$. That is, the diagonal turns intersections into fiber products.

This argument feels like it takes place in **Set**, but goes through word-for-word for schemes.

Definition 1.2. A morphism $\pi : X \rightarrow Y$ of schemes is a **locally closed embedding** if it factors as $\pi = \pi_1 \circ \pi_2$, where π_2 is a closed embedding and π_1 is an open embedding.

Proposition 1.3. For any $\pi : X \rightarrow Y$, δ_π is locally closed.

Proof. Let $\{V_i\}$ be an affine open cover of Y , so $V_i \cong \text{Spec } B_i$ for each B_i , and $\mathcal{U}_i = \{U_{ij}\}$ be an affine open cover of $\pi^{-1}(V_i)$ for each i . Then, $\{U_{ij} \times_{V_i} U_{ij'} : i, j, j'\}$ covers $X \times_Y X$. More interestingly, $\{U_{ij} \times_{V_i} U_{ij} : i, j\}$ covers $\text{Im}(\delta_\pi)$: this is because if $x \in U_{ij}$, then $\delta_\pi(x) \in p_1^{-1}(U_{ij})$ and in $p_2^{-1}(U_{ij})$, and $p_1^{-1}(U_{ij}) \cap p_2^{-1}(U_{ij}) = U_{ij} \times_{V_i} U_{ij}$.

Now, it suffices to show that $\delta_\pi : \delta_\pi^{-1}(U_{ij} \times_{V_i} U_{ij}) \rightarrow U_{ij} \times_{V_i} U_{ij}$ is closed, since the property of being a closed embedding is affine-local. Since each $U_{ij} \cong \text{Spec } A_{ij}$ is affine, then it suffices to understand what's happening ring-theoretically: the diagonal map corresponds to the ring morphism $A_{ij} \otimes_{V_i} A_{ij} \rightarrow A_{ij}$ sending $a \otimes a' \mapsto aa'$. This is clearly surjective, which is exactly the criterion for a morphism of schemes to be a closed embedding. \square

Corollary 1.4. *If X and Y are affine schemes, then δ_π is a closed embedding.*

Corollary 1.5. *If Δ denotes $\text{Im}(\delta_\pi)$, then for any open $V \subset Y$ and $U \subset \pi^{-1}(V)$, $\Delta \cap (U \times_V U') \cong U \cap U'$ is a homeomorphism of topological spaces.*

This follows because a locally closed embedding is homeomorphic onto its image.

These will all be super useful once we define separability, which we'll do now.

Definition 1.6. A morphism $\pi : X \rightarrow Y$ is **separated** if $\delta_\pi : X \rightarrow X \times_Y X$ is a closed embedding.

This is weird upon first glance: why do we look at the diagonal to understand things about a morphism? The answer is that the diagonal has nice category-theoretic properties, so we can prove some useful properties by doing a few diagram chases.

More geometrically, separability corresponds to the Hausdorff property in topological spaces, and there's a criterion for this in terms of the diagonal.

Proposition 1.7. *If T is a topological space, then T is Hausdorff iff the diagonal morphism $T \rightarrow T \times T$ is a closed embedding.*

Equivalently, the image $\Delta \subset T \times T$ is a closed subspace.

Remark. Since schemes are topological spaces, you might think this proves separated schemes are Hausdorff, but this is untrue: fiber products of schemes are generally not fiber products of underlying spaces, and therefore closed embeddings of schemes are not the same as closed embeddings of their underlying spaces.

Separability is a nice property, and is good to have. But like Hausdorffness, we generally won't need to use schemes that aren't separated.

Example 1.8.

- (1) By Corollary 1.4, all morphisms of affine schemes are separated.
- (2) If we can cover $X \times_Y X$ by the sets $U_{ij} \times_{V_i} U_{ij}$ (with these sets as in the proof of Proposition 1.3), then π is separated.
- (3) For a counterexample, let $X = \mathbb{A}_{(0,0)}^1$ be the “line with two origins” over a field k . This isn't a separated scheme: the diagonal is a “line with four origins,” and these cannot be separated topologically: every open set containing one contains all of them. So take one affine piece of X , which contains exactly one origin, and therefore its image ought to contain all four, but it doesn't, so $X \rightarrow \text{Spec } k$ isn't closed. This might feel a little imprecise, but one can make it fully rigorous.

We want separated morphisms to be nice: we'd like them to be preserved under base change and composition, and we'd like locally closed embeddings to be separated.

Proposition 1.9. *Locally closed embeddings are separated.*

This is the only example of a hands-on proof of a property; it's not hard, but the rest will be less abstract and easier. First, though, let's reframe it:

Proposition 1.10. *Any monomorphism of schemes is separated.¹*

Proof. By point (2) of Example 1.8, it suffices to prove that fiber products $U_{ij} \times_{V_i} U_{ij}$ cover $X \times_Y X$ for our affine covers. So let's look at the fiber diagram (1.1) again; it tells us that $\pi \circ p_1 = \pi \circ p_2$. But since π is a monomorphism, then $p_1 = p_2$, so for any $z \in X \times_Y Z$, $p_1(z) = p_2(z)$; call this point x_z . Then, if $x_z \in U_{ij}$, $z \in p^{-1}(U_{ij})$ and $z \in p_2^{-1}(U_{ij})$, and their intersection is the fiber product. \square

¹More is true in general; all you need is that $p_1 = p_2$ in the diagram (1.1), which is analogous to an injectivity condition on π . Hence, it suffices that π is injective as a map of sets, but this is a weird notion for schemes, so we generally phrase it in terms of monomorphisms.

Since locally closed embeddings are monomorphisms, Proposition 1.9 follows as a corollary.

At this point, we can define varieties, and Vakil does so, but can't do anything with them, so we'll come back to them in a little bit.

Proposition 1.11. *If A is a ring, $\mathbb{P}_A^n \rightarrow \operatorname{Spec} A$ is separated.*

The idea of the proof is to compute: we already know a cover of \mathbb{P}_A^n by $n + 1$ affine schemes, and can check that the induced map on rings is surjective.

The following proposition gives us an important geometric property of separability.

Proposition 1.12. *If A is a ring and $X \rightarrow \operatorname{Spec} A$ is separated, then for any affine open subsets $U, V \subset X$, $U \cap V$ is also affine.*

Proof. The diagonal is a closed embedding, so $\delta : U \times V \rightarrow U \times_A V$ is also a closed embedding. Therefore $U \times V$ is isomorphic to a closed subscheme of an affine scheme, and therefore is affine. \square

It's surprising how useful these arguments with the diagonal are: we got a useful and nontrivial result in one line! In general, you can prove a weirdly large amount of things by factoring them through the diagonal. In fact, let's use it to define another property.

Definition 1.13. A morphism $\pi : X \rightarrow Y$ is **quasiseparated** if δ_π is quasicompact.

This isn't the same as the other definition we were given, that for all affine $V \subset Y$ and $U, U' \subset \pi^{-1}(V)$, $U \cap U'$ is quasicompact. But it turns out to be equivalent.

Proposition 1.14. *$\pi : X \rightarrow Y$ is quasiseparated in the sense of Definition 1.13 iff it's quasiseparated in the sense we defined previously.*

The proof is a diagram chase involving the “magic diagram” for fiber products. This states that if $X_1, X_2 \rightarrow Y \rightarrow Z$ are maps in some category and the relevant fiber products exist, the diagram

$$\begin{array}{ccc} X_1 \times_Y X_2 & \longrightarrow & X_1 \times_Z X_2 \\ \downarrow & & \downarrow \\ Y & \xrightarrow{\delta} & Y \times_Z Y \end{array}$$

is a fiber diagram; the proof is a diagram chase following from the associativity of products, or checking the universal property. This diagram is also very ubiquitous for proofs like these.

Proposition 1.15. *Separability and quasiseparability are preserved under base change.*

Proof. Suppose $\pi : X \rightarrow Y$ is separated and $\varphi : S \rightarrow Y$ is another map of schemes, so there's an induced morphism $\pi' : Z = X \times_Y S \rightarrow S$ fitting into the diagram

$$\begin{array}{ccc} Z & \xrightarrow{\pi'} & S \\ \downarrow p_1 & & \downarrow \varphi \\ X & \xrightarrow{\pi} & Y. \end{array}$$

The magic diagram for this is the fiber diagram

$$\begin{array}{ccc} Z & \xrightarrow{\delta_{\pi'}} & Z \times_S Z \\ \downarrow & & \downarrow \\ X & \xrightarrow{\delta_\pi} & X \times_Y X. \end{array}$$

If π is separated, δ_π is closed, and therefore $\delta_{\pi'}$ is closed (since closed embeddings are preserved under base change), so π' is separated. The same argument works with π quasiseparated and δ_π quasicompact. \square

There are a few related properties that we won't prove, but whose proofs are very similar to the previous one.

Proposition 1.16. *Separability and quasiseparability are*

- (1) *local on the target*,
- (2) *closed under composition*, and
- (3) *closed under taking products*: if $\pi : X \rightarrow Y$ and $\pi' : X' \rightarrow Y'$ are separated morphisms of schemes over a scheme S , then $\pi \times \pi' : X \times_S X' \rightarrow Y \times_S Y'$ is separated; if π and π' are merely quasiseparated, so is $\pi \times \pi'$.

Each of these is a diagram chase with the right diagram, and not a particularly hard one; the last one follows as a general categorical consequence of the others.

Now, though, we can define varieties.

Definition 1.17. Let k be a field. A **k -variety** is a k -scheme $X \rightarrow \operatorname{Spec} k$ that is reduced, separated, and of finite type. A **subvariety** of a given variety X is a reduced, locally closed subscheme.

Reducedness is a property of X , but the others are properties of the structure morphism $X \rightarrow \operatorname{Spec} k$. Notice that the affine line with doubled origin is reduced and of finite type, so separability is important for avoiding pathologies.

It's nontrivial that a subvariety $Y \subset X$ is itself a variety. X is finite type over $\operatorname{Spec} k$, so it's covered by finitely many affine opens that are schemes of finitely generated k -algebras, which are Noetherian, so X is Noetherian. Hence, $Y \hookrightarrow X$ is a finite-type morphism into a Noetherian scheme, so Y is finite type; but we do need separability to be preserved under composition, which we just saw how to prove.

We did not require varieties to be irreducible; irreducibility doesn't behave as well as we would like, unless k is particularly nice.

Proposition 1.18. *The product of irreducible varieties over an algebraically closed field k is an irreducible k -variety.*

This follows from the nontrivial fact that if A and B are k -algebras that are integral domains, then $A \otimes_k B$ is an integral domain.

The last important thing we'll discuss today is a big meta-theorem about classes of morphisms.

Theorem 1.19 (Cancellation theorem). *Consider a commutative diagram*

$$\begin{array}{ccc} X & \xrightarrow{\pi} & Y \\ & \searrow \tau & \swarrow \rho \\ & Z, & \end{array}$$

i.e. $\tau = \rho \circ \pi$, and let P be a property of morphisms preserved under base change and composition. If τ has P and δ_ρ has P , then π also has P .

The name is because we're "cancelling" ρ out of the composition.

The proof uses the notion of the graph of a morphism.

Definition 1.20. Let X and Y be schemes over a scheme S , and $\pi : X \rightarrow Y$ be a map of S -schemes. Then, the **graph** of π is the morphism $\Gamma_\pi : X \rightarrow X \times_S Y$ defined by $\Gamma_\pi; (1_X, \pi)$.

That is, this sends a point to its image on the graph. We use this because any morphism factors through its graph. Then, since δ_ρ has P , so must Γ_π , which is useful. It seems weirdly abstract and pointless, but the idea is that the nice properties of the diagonal, including locally closed embeddings, can be canceled off. In fact, if Y is separated, we can cancel off properties of closed embeddings, and if Y is quasiseparated, we can cancel off properties of quasicompact morphisms.

Rational Maps. Let's talk about rational maps, which are rational maps defined almost everywhere, and up to almost everywhere agreement. Rational maps are usually only defined on reduced varieties, since it's nearly impossible to get a hold on them otherwise; they're inherently geometric, and geometry tends to involve varieties.

Definition 1.21. A **rational map** $\pi : X \dashrightarrow Y$ is an equivalence class of morphisms $f : U \rightarrow Y$, where $U \subset X$ is a dense open subset; (f, U) and (f', U') are considered equivalent if there's a dense open set $V \subset U \cap U'$ if $f|_V = f'|_V$. One says π is **dominant** if its image is dense, or equivalently, for all nonempty opens $V \subseteq Y$, $\pi^{-1}(V) \neq \emptyset$.

Notice that dominance is well-defined, as it's independent of choice of representative.

Proposition 1.22. *Let X and Y be irreducible schemes, then $\pi : X \dashrightarrow Y$ is dominant iff the generic point of X maps to the generic point of Y .*

Proof. In the reverse direction, the generic point η_Y of Y is contained in every open subset of Y , so the preimage contains the generic point η_X of X , and in particular is nonempty.

In the other direction, suppose $\pi(\eta_X) \neq \eta_Y$; let $U = Y \setminus \overline{\pi(\eta_X)}$, which is an open subset. Thus, $\eta_X \notin \pi^{-1}(U)$, which is an open set. Since η_X is dense, it meets every nonempty open, so $\pi^{-1}(U)$ is empty, and therefore π isn't dominant. \square

This is a pretty useful characterization of dominance. But why do we care about dominance? Because of composition.

Remark. Let $\pi : X \dashrightarrow Y$ and $\rho : Y \dashrightarrow Z$ be rational maps. If π is dominant and X is irreducible, it's possible to make sense of $\rho \circ \pi : X \dashrightarrow Z$ as a rational map, which is dominant iff ρ is.

This is nontrivial: if π isn't dominant, one might discover that the domain of ρ doesn't intersect the image of π ; if they do, however, π^{-1} of the domain of definition of ρ is a nonempty open of X ; since X is irreducible, it must be dense.

Definition 1.23. A rational map $\pi : X \dashrightarrow Y$ is **birational** if it's dominant and there exists a dominant $\psi : Y \dashrightarrow X$ such that as rational maps, $\pi \circ \psi \sim 1_X$ and $\psi \circ \pi \sim 1_Y$. In this case, one says π and ψ are **birational(ly equivalent)**.

Proposition 1.24. *Let X and Y be reduced schemes; then, X and Y are birational iff there exist dense open subschemes $U \subset X$ and $V \subset Y$ such that $U \cong V$.*

The idea is that we can let U and V be the domains of definition for our rational maps.

The notion of rationality is very specific to algebraic geometry; in the differentiable category, it's complete nonsense. Since any manifold can be triangulated, any two manifolds of the same dimension are birationally equivalent: remove the edges of the triangles, and you get a dense open set; clearly, any two triangles are birational. However, there exist algebraic varieties of the same dimension that aren't birationally equivalent.

Definition 1.25. A variety X over k is **rational** if it's birational to \mathbb{A}_k^n for some n .

For example, \mathbb{P}_k^n is rational. Rationality loses some information, but what it keeps is interesting.

Finally, let's see what dominance means in terms of ring morphisms.

Definition 1.26. Let $\varphi : \text{Spec } A \rightarrow \text{Spec } B$ be a morphism of affine schemes and $\varphi^\# : B \rightarrow A$ be the induced map on global sections. Then, φ is dominant (i.e. as a rational map) iff $\ker(\varphi^\#) \subset \mathfrak{N}(A)$.

Here, $\mathfrak{N}(A)$ denotes its nilradical, the intersection of all prime ideals of A (equivalently, the ideal of nilpotent elements). That is, if A and B are reduced, dominance is equivalent to injectivity! Interestingly, this also corresponds to an inclusion of function fields, i.e. a field extension! We've reduced a geometric problem to a problem about algebra. Often, we can go in the other direction, e.g. for varieties. In this setting, birationality means isomorphism on the function fields.

2. PROPER MORPHISMS: 5/19/16

These are Arun's lecture notes on rational maps to separated schemes and proper morphisms, corresponding to sections 10.2 and 10.3 in Vakil's notes. I'm planning on talking about the following topics:

- Rational maps to separated schemes, including the reduced-to-separated theorem and some corollaries.
- The definition of proper morphisms, and that they form a nice class of morphisms. Projective A -schemes are proper over A .

Throughout this lecture, S is a scheme, which will often be the base scheme.

Rational Maps to Separated Schemes. If X and Y are spaces and $\pi, \pi' : X \rightrightarrows Y$ are continuous, it's sometimes useful to talk about the locus where they agree, $\{x \in X : \pi(x) = \pi'(x)\}$. Categorically, this is the equalizer $\text{Eq}(\pi, \pi') \hookrightarrow X$, which is characterized by the property that if $\varphi : W \rightarrow X$ is a continuous map such that $\pi \circ \varphi = \pi' \circ \varphi$, then it factors through $\text{Eq}(\pi, \pi')$, i.e. there's a unique $h : W \rightarrow \text{Eq}(\pi, \pi')$ such that the following diagram commutes.

$$\begin{array}{ccc} W & & \\ \downarrow & \searrow \varphi & \\ \text{Eq}(\pi, \pi') & \xrightarrow{\quad} & X \xrightleftharpoons[\pi']{\pi} Y. \end{array}$$

$\exists! \downarrow h$

So if we can do this for schemes, we'll have a subscheme where two morphisms agree, rather than just a set. The universal property for the equalizer is the same as for the fiber product

$$\begin{array}{ccc} \text{Eq}(\pi, \pi') & \longrightarrow & Y \\ \downarrow i & \lrcorner & \downarrow \delta \\ X & \xrightarrow{(\pi, \pi')} & Y \times Y, \end{array} \quad (2.1)$$

where δ is the diagonal morphism. We know fiber products of schemes exist, so equalizers do too.

Lemma 2.2 (Vakil ex. 10.2.A). *If $\pi, \pi' : X \rightrightarrows Y$ are two morphisms of schemes over S , then $i : \text{Eq}(\pi, \pi') \hookrightarrow X$ is a locally closed subscheme of X . If Y is separated over S , $\text{Eq}(\pi, \pi')$ is a closed subscheme.*

Proof. Since we're over S , the product in (2.1) should be replaced with $Y \times_S Y$, the product in \mathbf{Sch}_S . Since δ is a locally closed embedding, and this is a property preserved under base change, then i is too. If $Y \rightarrow S$ is separated, then δ is a closed embedding, and this is also preserved by pullbacks. \square

Remark. The locus where two maps agree does not need to be reduced, e.g. if $\pi, \pi' : \mathbb{A}_{\mathbb{C}}^1 \rightarrow \mathbb{A}_{\mathbb{C}}^1$ are defined by $\pi(x) = 0$ and $\pi'(x) = x^2$, then they agree “to first order” at 0, and $\text{Eq}(\pi, \pi') = \text{Spec } \mathbb{C}[x]/(x^2)$.

The central result about these is the reduced-to-separated theorem.

Theorem 2.3 (Reduced-to-separated theorem (Vakil Thm. 10.2.2)). *Let $\pi, \pi' : X \rightrightarrows Y$ be two morphisms of S -schemes. If X is reduced, Y is separated over S , and π and π' agree on a dense open subset, then $\pi = \pi'$.*

This is equality in the sense of morphisms of schemes, which is stronger than pointwise equality.

Proof. By Lemma 2.2, $\text{Eq}(\pi, \pi') \hookrightarrow X$ is a closed subscheme, but it contains a dense open set. Since X is reduced, its only closed subscheme containing a dense open set is itself. \square

Corollary 2.4. *If X is reduced, Y is separated, and $\pi : X \dashrightarrow Y$ is a rational map, then there is a maximal $U \subset X$ such that $\pi|_U : U \rightarrow Y$ is an honest morphism. In particular, this is true for rational functions on reduced schemes.*

This U is called the **domain of definition** of π ; its complement is sometimes called the **locus of indeterminacy**.

Proof. We can choose U to be the union of all domains of representatives of π . If $f_1 : V_1 \rightarrow Y$ and $f_2 : V_2 \rightarrow Y$ are two morphisms representing π , then f_1 and f_2 agree on a dense open subset of $V_1 \cap V_2$, so by the reduced-to-separated theorem agree on all of $V_1 \cap V_2$. Thus, we can glue representing morphisms on their intersection and therefore define π on all of U . \square

Next, we need to digress slightly to understand the image of a locally closed embedding. This is from section 8.3 of the notes.

If $\pi : X \rightarrow Y$ is a morphism of schemes, it's in particular a continuous function, so its image $\pi(X) \subset Y$ is a subspace. This will be referred to as the **set-theoretic image**. As usual, the topological version of a thing tends to be less well-behaved than the scheme-theoretic one, so we'll define an image of π that's a subscheme of Y . Schemes are locally cut out by equations, so it seems reasonable to say that a closed subscheme $i : Z \hookrightarrow Y$ **contains the image** of π if functions in \mathcal{O}_Y that vanish on Z also vanish when pulled back to X . That is, the composition $\mathcal{I}_{Z/Y} \rightarrow \mathcal{O}_Y \rightarrow \pi_* \mathcal{O}_X$ is zero, where $\mathcal{I}_{Z/Y} = \ker(i^\# : \mathcal{O}_Y \rightarrow i_* \mathcal{O}_Z)$ is the sheaf of ideals associated to the closed embedding of Z into Y .

Definition 2.5. The **scheme-theoretic image** $\text{Im}(\pi)$ of π is the intersection of all closed subschemes containing the image of π .² If π is a locally closed embedding, $\text{Im}(\pi)$ is also called the **scheme-theoretic closure** of π .

That is, $\text{Im}(\pi)$ is the smallest closed subscheme of Y such that locally vanishing on $\text{Im}(\pi)$ implies locally vanishing when pulled back to X .

Theorem 2.6 (Vakil cor. 8.3.5). *Let $\pi : X \rightarrow Y$ be a morphism of schemes. If X is reduced or Y is quasicompact, the closure of the set-theoretic image of π is the underlying set of $\text{Im}(\pi)$.*

We lack the time to prove this, but it follows from the defining properties of closed embeddings.

Just like we defined the graph of a morphism of S -schemes $\pi : X \rightarrow Y$ to be $\Gamma_\pi = (\text{id}, \pi) : X \rightarrow X \times_S Y$, we can define the graph of a rational map in nice situations.

Definition 2.7. Let $\pi : X \dashrightarrow Y$ be a rational map over S , where X is reduced and Y is separated over S . For any representative morphism $f : U \rightarrow Y$ of π , the **graph of the rational map** π , denoted Γ_π , is the scheme-theoretic closure of the map $\Gamma_f \hookrightarrow U \times_S Y \hookrightarrow X \times_S Y$. (The first map is a closed embedding, and the second is an open embedding.)

The following diagram might make this definition clearer.

$$\begin{array}{ccc} \Gamma_\pi & \xrightarrow{\text{cl.}} & X \times_S Y \\ \uparrow & \swarrow & \searrow \\ X & \xrightarrow{\pi} & Y \end{array}$$

A priori this definition depends on the choice of representative, but fortunately, this isn't actually the case.

Proposition 2.8 (Vakil ex. 10.2.E). *The graph of a rational map π is independent of choice of representative.*

Proof. Let $\xi' : U \rightarrow Y$ and $\xi : V \rightarrow Y$ be two representatives of π . Without loss of generality, we can assume V is the maximal domain of definition for π , so $U \subset V$ and $\xi' = \xi|_U$. Thus, we have a bunch of embeddings fitting into the diagram

$$\begin{array}{ccccc} \Gamma_{\xi'} & \xrightarrow{\text{cl.}} & U \times Y & \xrightarrow{\text{op.}} & X \times Y \\ \downarrow & & \downarrow \text{op.} & \nearrow \text{op.} & \\ \Gamma_\xi & \xrightarrow{\text{cl.}} & V \times Y & & \end{array}$$

Thus, $\Gamma_{\xi'}$ factors as a subset of a closed subset of $V \times Y$, so its scheme-theoretic closure, which is just the closure of its underlying set by Theorem 2.6, must factor through this. In particular, the graph of π as defined with respect to ξ' embeds into $V \times Y$. Thus, we can assume $V = X$, since everything takes place inside V . In this case, Γ_π as defined by ξ is just Γ_ξ , and $\Gamma_\xi \cong X$ by projection onto the first factor. This projection restricts to an isomorphism $\Gamma_{\xi'} \cong U$, and carries the embedding $\Gamma_\xi \hookrightarrow \Gamma_{\xi'}$ to the embedding $U \hookrightarrow X$. Finally, to form the graph of π with respect to ξ' , we take the closure, and since U is a dense open subset, we get X , or all of Γ_ξ . \square

Finally, we discuss one application to effective Cartier divisors. (This is actually an excuse to introduce effective Cartier divisors, since they show up again and again.)

Definition 2.9. A closed embedding $\pi : X \hookrightarrow Y$ is an **effective Cartier divisor** if $\mathcal{I}_{X/Y}$ is locally generated by a single non-zerodivisor. That is, there's an affine open cover \mathcal{U} of Y such that for each $U_i = \text{Spec } A_i \in \mathcal{U}$, there's a $t_i \in A_i$ that is not a zerodivisor and such that $\mathcal{I}_{X/Y}(U_i) = A_i/(t_i)$.

Proposition 2.10 (Vakil ex. 10.2.G). *Let X be a reduced S -scheme and Y be a separated S -scheme. If $i : D \hookrightarrow X$ is an effective Cartier divisor, there is at most one way to extend an S -morphism $\pi : X \setminus D \rightarrow Y$ to all of X .*

²There's something to prove here, that containing the image of π is well-behaved under intersections.

Proof. This is true if we know it on an affine cover, so without loss of generality assume $X = \operatorname{Spec} A$ is affine and $D = V(t)$ for some $t \in A$ that isn't a zerodivisor. If $D(t) = X \setminus D$ is dense in X , then we're done by Theorem 2.3. Since X is reduced, then by Theorem 2.6 this is equivalent to the scheme-theoretic closure of $D(t)$ being all of X . Given a closed subscheme $Z \hookrightarrow X$, we want to understand when functions vanishing on Z pull back to the zero function on $D(t)$. The map $\Gamma(X, \mathcal{O}_X) \rightarrow \Gamma(D(t), \mathcal{O}_X)$ is also $A \rightarrow A_t$; since t isn't a zerodivisor, this is injective, so a function pulls back to 0 on $D(t)$ iff it vanishes on all of X . Hence, $\operatorname{Im}(D(t) \hookrightarrow X) = X$ as desired. \square

Proper Morphisms. The next topological notion we introduce to algebraic geometry is that of a proper map. Recall that a continuous map of topological spaces is proper if the preimage of any compact set is compact. Compactness doesn't really behave the same way in algebraic geometry, so we'll have to define properness in a different way, which will satisfy similar properties.

Proper maps are closed maps, meaning the image of a closed set is closed. This would be a reasonable starting point, except that closed maps are not preserved by fiber products. It turns out the right way to fix this is just to pick the ones that behave well.

Definition 2.11. A morphism $\pi : X \rightarrow Y$ of schemes is **universally closed** if for all morphisms $Z \rightarrow Y$, the pullback $Z \times_Y X \rightarrow Z$ is a closed map.

That is, it remains closed under arbitrary base change.

Lemma 2.12. *Universal closure is a “nice” property of schemes, i.e. local on the target, closed under composition, and preserved by base change.*

Proof. Clearly, universal closure is closed under composition, and by definition, it's preserved by fiber products. Being a closed map is local on the target, and therefore so is universal closure. \square

We use universal closure to define the property we really care about.

Definition 2.13. A morphism $\pi : X \rightarrow Y$ is **proper** if it's separated, finite type, and universally closed. If A is a ring, an A -scheme X is said to be **proper over A** if the structure morphism $X \rightarrow \operatorname{Spec} A$ is proper.

Example 2.14. Closed embeddings are our first example of proper morphisms: they're affine, and therefore separated. Closed embeddings are closed maps, and since the pullback of a closed embedding is a closed embedding, a closed embedding is universally closed. Finally, closed morphisms are finite type (which boils down the fact that if $B \twoheadrightarrow A$ is a surjective ring map, A is a finitely generated B -algebra).³

This agrees with our intuition for topological spaces, which is good.

Proposition 2.15 (Vakil prop. 10.3.4).

- (1) *Properness is a “nice” property of schemes (in the sense of Lemma 2.12).*
- (2) *Properness is closed under products: if $\pi : X \rightarrow Y$ and $\pi' : X' \rightarrow Y'$ are proper morphisms of S -schemes, then $\pi \times \pi' : X \times_S X' \rightarrow Y \times_S Y'$ is proper.*
- (3) *Given a commutative diagram*

$$\begin{array}{ccc} X & \xrightarrow{\pi} & Y \\ & \searrow \tau & \swarrow \rho \\ & Z, & \end{array}$$

if τ is proper and ρ is separated, then π is proper.

For example, by (3), any morphism from a proper k -scheme to a separated k -scheme is proper (let $Z = \operatorname{Spec} k$).

Proof. Everything in this proposition comes nearly for free. We already knew finite type and separability to be nice properties of schemes, and by Lemma 2.12, so is universal closure; since properness is having all three at once, it too must be a nice property. (2) is a formal consequence of (1), which is proven for any nice class

³The same line of reasoning shows that finite morphisms are proper, which is a generalization: they're affine, hence separated, and closed maps; since they're preserved under base change, they must also be universally closed. Finally, finite morphisms are finite type.

of morphisms in Vakil's ex. 9.4.F. Finally, since closed embeddings are proper, the cancellation theorem from last lecture applies to prove (3). \square

According to Vakil, the next example is the most important example of proper morphisms.

Theorem 2.16 (Vakil thm. 10.3.5). *If A is a ring and X is a projective A -scheme, $X \rightarrow \operatorname{Spec} A$ is proper.*

Proof. Since X is projective, the structure morphism factors as $X \hookrightarrow \mathbb{P}_A^n \rightarrow A$, a closed embedding followed by the structure map for \mathbb{P}_A^n . Since closed embeddings are proper (Example 2.14), it suffices to show $\mathbb{P}_A^n \rightarrow \operatorname{Spec} A$ is proper, because proper morphisms are closed under composition. Projective schemes are finite type, and we proved last time that $\mathbb{P}_A^n \rightarrow \operatorname{Spec} A$ is separated, so it remains to check universal closure.

If $\varphi : X \rightarrow \operatorname{Spec} A$ is an arbitrary morphism, we would like for the map $\mathbb{P}_A^n \times_A X \rightarrow X$ to be closed. Since $\mathbb{P}_A^n = \mathbb{P}_{\mathbb{Z}}^n \times_A \operatorname{Spec} \mathbb{Z}$, then we have the following commutative diagram, in which both squares are pullback squares:

$$\begin{array}{ccc} \mathbb{P}_A^n \times_A X & \xrightarrow{\quad} & X \\ \downarrow & \lrcorner & \downarrow \varphi \\ \mathbb{P}_A^n & \xrightarrow{\quad} & \operatorname{Spec} A \\ \downarrow & \lrcorner & \downarrow \\ \mathbb{P}_{\mathbb{Z}}^n & \xrightarrow{\quad} & \operatorname{Spec} \mathbb{Z} \end{array}$$

By checking the universal property, we see that the outer rectangle is a pullback square too: in other words, $\mathbb{P}_A^n \times_A X = \mathbb{P}_X^n$, so it suffices to show that the structure map $\mathbb{P}_X^n \rightarrow X$ is closed for arbitrary X . Being a closed map is a local condition, so we can check on an affine cover of \mathbb{P}_X^n ; pulling back by $\operatorname{Spec} B \hookrightarrow X$ gives us $\mathbb{P}_B^n \rightarrow \operatorname{Spec} B$, so it suffices to know that the structure map is closed for all rings B . This is precisely the fundamental theorem of elimination theory (Thm. 7.4.7 in Vakil's notes), so we're done. \square

Perhaps surprisingly, the converse is almost true: it's difficult to come up with examples of schemes that are proper, but not projective.

The last thing we'll prove about proper schemes is another analogue of compactness. Recall that if M is a compact, connected complex manifold, all holomorphic functions on M are constant. We'll be able to prove a scheme-theoretic analogue of this.

Proposition 2.17 (Vakil 10.3.7). *Let k be an algebraically closed field and X be a connected, reduced, proper k -scheme. Then $\Gamma(X, \mathcal{O}_X) \cong k$.*

Proof. First, we can naturally identify $\Gamma(X, \mathcal{O}_X)$ with the ring of k -scheme maps $X \rightarrow \mathbb{A}_k^1$: using the $(\Gamma, \operatorname{Spec})$ adjunction, $\operatorname{Hom}_{\operatorname{Sch}_k}(X, \mathbb{A}_k^1) = \operatorname{Hom}_{\operatorname{Alg}_k}(k[t], \Gamma(X, \mathcal{O}_X)) = \Gamma(X, \mathcal{O}_X)$, so functions on X are actually a ring of functions, which is nice.

Let $f \in \Gamma(X, \mathcal{O}_X)$, so f corresponds to a morphism $\pi : X \rightarrow \mathbb{A}_k^1$. If $i : \mathbb{A}_k^1 \hookrightarrow \mathbb{P}_k^1$ is the usual open embedding, let $\pi' = i \circ \pi$. Since X is proper and \mathbb{P}_k^1 is separated over k , then π' must be proper, by Proposition 2.15, part 3 (let $Z = \operatorname{Spec} k$). Thus, π' is closed, so the set-theoretic image of π' is a closed, connected subset of \mathbb{P}_k^1 . Since \mathbb{P}_k^1 has the cofinite topology, then $\operatorname{Im}(\pi)$ must be a single closed point p or all of \mathbb{P}_k^1 , but if the latter, it can't factor through i . Since π' factors through \mathbb{A}_k^1 , p is a closed point in \mathbb{A}_k^1 , hence identified with an element of k .

The underlying set of the scheme-theoretic image of π is the closure of the set-theoretic image, so it's just p again; since X is reduced, so is its scheme-theoretic image. Thus, $\pi : X \rightarrow \mathbb{A}_k^1$ is a constant map of schemes $x \mapsto p$, and tracing through the adjunction, this corresponds to the constant function $f = p \in \Gamma(X, \mathcal{O}_X)$. \square

3. DIMENSION: 5/23/16

Today's lecture was given by Gill Grindstaff, on the first half of Chapter 11. This section relies on a lot of commutative algebra, which can make it difficult.

There are equivalent topological and algebraic formulations of the definition of the dimension of a scheme. We'd like this to agree with the intuitive notions of dimension: \mathbb{A}^n should be n -dimensional, for example.

To motivate these definitions, recall that the dimension of a vector space V is the cardinality of some (and therefore any) basis for V . However, it's equivalent to say that the dimension of V is the supremum of lengths

of nested chains of subspaces $0 \subsetneq W_1 \subsetneq W_2 \subsetneq \cdots \subsetneq W_n = V$ (so we don't count 0). The scheme-theoretic definition will resemble this.

Definition 3.1.

- The **Krull dimension of a topological space** X is the supremum of lengths of chains $X_1 \subsetneq X_2 \subsetneq \cdots \subsetneq X_n = X$ in which each X_i is a closed, irreducible subset of X .
- The **Krull dimension of a ring** A is the supremum of lengths of chains $0 \subsetneq \mathfrak{p}_1 \subsetneq \mathfrak{p}_2 \subsetneq \cdots \subsetneq \mathfrak{p}_n$ of prime ideals in A .

Fact. If A is a ring, then $\dim \operatorname{Spec} A = \dim A$. This is because $\mathfrak{p}_i \mapsto V(\mathfrak{p}_i)$ defines an inclusion-reversing bijection between the poset of prime ideals of A and the poset of closed irreducible subspaces of $\operatorname{Spec} A$, and in particular sends chains of nested subsets to chains of nested subsets (in the other direction).

Example 3.2.

- (1) Every prime ideal of \mathbb{Z} is of the form $\mathfrak{p} = (p)$ for a prime number p , or is the zero ideal. Hence, the longest chain we can make is $(p) \supset (0)$, so $\dim \mathbb{Z} = 1$.
- (2) Similarly, for any field k , in $k[t]$ the longest chains we can make are $(f(t)) \supset (0)$ for f irreducible, so $\dim \mathbb{A}_k^1 = 1$.
- (3) In $k[x]/(x^2)$, (0) is the only prime ideal, so $\dim k[x]/(x^2) = 0$.

Dimension is *not* local, unlike the dimension of manifolds: consider the space $Z \subset \mathbb{A}^3$ consisting of the union of the xy -plane and the z -axis, which is not irreducible. Then, the dimension of the xy -plane is 2 but the dimension of the z -axis is 1. We do know that $\dim(Z)$ is the maximum of the dimensions of its irreducible subsets, however.

Definition 3.3. A scheme X is **equidimensional** if each of its irreducible components has the same dimension.

An equidimensional scheme of dimension 1 is called a **curve**; an equidimensional scheme of dimension 2 is called a **surface**; and so forth.

In order to get a handle on dimension, we'll need to do some commutative algebra.

Theorem 3.4. Let $\pi : \operatorname{Spec} A \rightarrow \operatorname{Spec} B$ be induced from an integral extension $A \rightarrow B$ of rings. Then, $\dim \operatorname{Spec} A = \dim \operatorname{Spec} B$.

This follows from an algebraic result.

Theorem 3.5 (Going-up theorem). Let $A \hookrightarrow B$ be an integral extension, $\mathfrak{p}_1 \subset \mathfrak{p}_2$ be prime ideals of A , and \mathfrak{q}_1 be a prime ideal of B such that $\mathfrak{q}_1 \cap A = \mathfrak{p}_1$.⁴ Then, there is a prime ideal $\mathfrak{q}_2 \subset B$ such that $\mathfrak{q}_1 \subset \mathfrak{q}_2$ and $\mathfrak{q}_2 \cap A = \mathfrak{p}_2$.

This can be inductively extended to chains of prime ideals, which proves one half of Theorem 3.4.

The proof of Theorem 3.5 depends on the following lemma.

Lemma 3.6. Let $A \hookrightarrow B$ be an integral extension. If B is a field, then A is a field.

The following exercise is another nice property.

Exercise 3.7. Let $\nu : \tilde{X} \rightarrow X$ be the normalization. Then, $\dim \tilde{X} = \dim X$.

The **normalization** of X replaces rings with their integral closures on an affine cover; after checking that this behaves well, it defines a nice scheme that X embeds into as an open dense subset. The key to the proof is that the dimension of a ring is the same as the dimension of its integral closure, which follows from Theorem 3.4.

The next thing we'd like to define is codimension, but there are some weird pathologies: recalling Z , the union of the xy -plane and the z -axis, what's the codimension of the z -axis in Z ? Should it be 0, since there's nothing above it? Or is it 1, since Z is 2-dimensional and the z -axis is 1-dimensional? There's no good answer, and as a result one only defines codimension inside irreducible schemes.

Definition 3.8. Let X be an *irreducible* topological space and $Y \subset X$. Then, the **codimension** $\operatorname{codim}_X Y$ is the supremum of lengths of chains of irreducible closed subsets $\bar{Y} \subsetneq Z_1 \subsetneq Z_2 \subsetneq \cdots \subsetneq Z_n = X$.

⁴One often says that \mathfrak{q}_1 **lies over** \mathfrak{p}_1 .

Since Y might not be closed, we must start with \overline{Y} . This is satisfactory: a closed point has codimension 2 inside \mathbb{A}^2 .

There's also an algebraic analogue of codimension.

Definition 3.9. The **codimension of a prime ideal** \mathfrak{p} in a ring A , written $\text{codim}_R \mathfrak{p}$, is the supremum of lengths of *decreasing* chains of prime ideals $\mathfrak{p} \supsetneq \mathfrak{q}_1 \supsetneq \mathfrak{q}_2 \supsetneq \cdots \supsetneq \mathfrak{q}_n \supsetneq (0)$.

In particular, this implies that $\text{codim}_R \mathfrak{p} = \dim R_{\mathfrak{p}}$.

Here are some useful results about dimension.

Proposition 3.10. *Let R be a UFD and $\mathfrak{p} \subset R$ be a codimension-one prime ideal. Then, \mathfrak{p} is principal.*

Theorem 3.11. *Let A be a finitely generated k -algebra that's an integral domain. Then, $\dim \text{Spec } A = \text{tr. deg. } K(A)/k$.*

Here, $K(A)$ denotes the field of fractions of A , and $\text{tr. deg. } K/k$ is the **transcendence degree** of a field extension K/k ; the idea is: how many transcendental elements do you need to adjoin to get K ? This intuition turns out to be correct.

Lemma 3.12 (Noether normalization). *Let S be a finitely generated k -algebra that's an integral domain, such that $\text{tr. deg. } K(A)/k = n$. Then, there exist $x_1, \dots, x_n \in A$, algebraically independent⁵ over k , such that A is a finite extension of $k[x_1, \dots, x_n]$.*

These are very useful because the transcendence degree is much easier to understand than all prime ideals of a ring.

Just as we have a going-up theorem, there's also one in the pther dimension.

Theorem 3.13 (Going-down theorem). *Let $\phi : B \hookrightarrow A$ be a finite extension of rings (i.e. A is finitely-generated as a B -module), where B is an integrally closed domain and A is an integral domain. Suppose $\mathfrak{q}_1 \subset \mathfrak{q}_2$ are prime ideals of B and $\mathfrak{p}_2 \subset A$ is a prime ideal such that $\phi^{-1}(\mathfrak{p}_2) = \mathfrak{q}_2$ (so it lies over \mathfrak{q}_2). Then, there exists a prime $\mathfrak{p}_1 \subset \mathfrak{p}_2$ such that \mathfrak{p}_1 lies over \mathfrak{q}_1 .*

This requires more assumptions and is harder to prove.

At this point, we talked about a few exercises.

Lemma 3.14. *Let X be a topological space and $U \subset X$ be open. Then, there's a bijection between the irreducible closed subsets of U and the irreducible closed subsets of X meeting U .*

This resembles a theorem from commutative algebra establishing a bijection between prime ideals of B/I and prime ideals of B containing I (where $I \subset B$ is an ideal).

The proof of Lemma 3.14 sets up the bijection by sending an irreducible closed subset $F \subset U$ to $\overline{F} \subset X$, and sends an irreducible $F' \subset X$ to $F' \cap U \subset U$.

Exercise 3.15 (Vakil ex. 11.1.B). Show that a scheme has dimension n iff it can be covered by affine open subsets of dimension at most n , where equality is achieved for some affine scheme in the cover.

⁵Just as linear independence means not satisfying any nonzero linear relation, **algebraic independence** means not satisfying any nonzero polynomial relation.