

FURUTA’S 10/8 THEOREM

ARUN DEBRAY
JANUARY 23, 2019

These notes were taken in a learning seminar on Furuta’s 10/8 theorem in Spring 2019. I live-TeXed them using vim, and as such there may be typos; please send questions, comments, complaints, and corrections to a.debray@math.utexas.edu.

CONTENTS

1. Introduction to Seiberg-Witten theory: 1/23/19	1
References	2

1. INTRODUCTION TO SEIBERG-WITTEN THEORY: 1/23/19

Riccardo gave the first, introductory talk.

In 1982, Matsumoto conjectured that if M is a closed spin manifold, $b_2(M) \geq (11/8)|\sigma(M)|$. Here $b_2(M)$ is the second Betti number and $\sigma(M)$ is the signature. Equality holds for the K3 surface, so this is the best one can do.

In this seminar we’ll study a theorem of Furuta which makes major progress on this conjecture.

Theorem 1.1 (10/8 theorem [Fur01]). *If the intersection form of M is indefinite, $b_2(M) \geq (10/8)|\sigma(M)| + 2$.*

If the intersection form is definite, work of Donaldson [Don83] says that, up to a change of orientation, the intersection form is diagonalizable, so that case is dealt with.

Furuta’s proof uses both Seiberg-Witten theory and equivariant homotopy theory. It can be pushed a little bit farther, but not enough to prove the $11/8^{\text{th}}$ conjecture, as shown recently by Hopkins-Lin-Shi-Xu [HLSX18].

Today we’ll discuss some background for the proof.

Definition 1.2. Let $V \rightarrow M$ be a rank- n real oriented vector bundle. A *spin structure* on V is data $\mathfrak{s} = (P_{\text{Spin}}(V), \tau)$, where $P_{\text{Spin}}(V) \rightarrow M$ is a principal Spin_n -bundle and τ is an isomorphism

$$\tau: P_{\text{Spin}}(V) \times_{\text{Spin}_n} \mathbb{R}^n \xrightarrow{\cong} V.$$

A spin structure on a manifold M is a spin structure on TM .

Remark 1.3. There are other equivalent definitions of spin structures – for example, just as an orientation is a trivialization of V over the 1-skeleton of M , a spin structure is equivalent to a trivialization over the 2-skeleton. ◀

Here’s a cool theorem about spin manifolds.

Theorem 1.4 (Rokhlin [Roh52]). *If M is a spin manifold, $\sigma(M) \equiv 0 \pmod{16}$.*

The signature makes sense when $4 \mid \dim M$. Smoothness is crucial here; there are topological spin 4-manifolds, whatever that means, that do not satisfy this theorem. Freedman’s E_8 manifold is an example.

Suppose M is a spin 4-manifold. The representation theory of Spin_4 , in particular the fact that the spin representation S splits as $S^+ \oplus S^-$, leads to two quaternionic line bundles $\mathbb{S}^+, \mathbb{S}^- \rightarrow M$ with Hermitian metrics. Physics cares about these bundles, and will lead to powerful theorems in manifold topology.

These bundles have more structure: in particular, they are Clifford bundles.

Definition 1.5. Let $S \rightarrow M$ be a real vector bundle with a Euclidean metric $\langle \cdot, \cdot \rangle$. A *Clifford bundle* structure is data of, for each $x \in M$, the data of a Clifford algebra action $\text{Cl}(T_x M)$ on S_x that varies smoothly in x , such that the Clifford action is skew-adjoint, meaning

$$\langle v \cdot s_1, s_2 \rangle = -\langle s_1, v \cdot s_2 \rangle.$$

We also require the existence of a connection which is compatible with the Levi-Civita connection on TM .

Given the data of a Clifford bundle, there's an operator called the *Dirac operator* D , which is the following composition:

$$(1.6) \quad C^\infty(S) \xrightarrow{\nabla^{C\ell}} C^\infty(T^*M \otimes S) \xrightarrow{\langle \cdot, \cdot \rangle} C^\infty(TM \otimes S) \xrightarrow{\text{Clifford action}} C^\infty(S).$$

This operator is denoted \not{D} , a convention due to Feynman. It is a first-order, elliptic differential operator; ellipticity means that its analysis is nice.

Thus we can consider the *Seiberg-Witten equations* on a spin 4-manifold. Let $(a, \varphi) \in \Omega_M^1(i\mathbb{R}) \times \Gamma(\mathbb{S}^+)$; then the equations are

$$(1.7a) \quad \not{D}\varphi + \rho(a)(\varphi) = 0$$

$$(1.7b) \quad \rho(d^+a) - \varphi \otimes \varphi^* + \frac{1}{2}|\varphi|^2 \text{id} = 0$$

$$(1.7c) \quad d^*a = 0.$$

On a non-spin manifold, the equations are a little more complicated.

REFERENCES

- [Don83] S. K. Donaldson. An application of gauge theory to four-dimensional topology. *J. Differential Geom.*, 18(2):279–315, 1983. https://projecteuclid.org/download/pdf_1/euclid.jdg/1214437665. 1
- [Fur01] M. Furuta. Monopole equation and the $\frac{11}{8}$ -conjecture. *Math. Res. Lett.*, 8(3):279–291, 2001. http://intlpress.com/site/pub/files/_fulltext/journals/mrl/2001/0008/0003/MRL-2001-0008-0003-a005.pdf. 1
- [HLSX18] Michael J. Hopkins, Jianfeng Lin, XiaoLin Danny Shi, and Zhouli Xu. Intersection forms of spin 4-manifolds and the Pin(2)-equivariant Mahowald invariant. 2018. <http://arxiv.org/abs/1812.04052>. 1
- [Roh52] V. A. Rohlin. New results in the theory of four-dimensional manifolds. *Doklady Akad. Nauk SSSR (N.S.)*, 84:221–224, 1952. 1