


Riemann Surfaces



UT Austin, Spring 2016

M392C NOTES: RIEMANN SURFACES

ARUN DEBRAY
JANUARY 20, 2016

These notes were taken in UT Austin's Math 392C (Riemann Surfaces) class in Spring 2016, taught by Tim Perutz. I live-TeXed them using `vim`, and as such there may be typos; please send questions, comments, complaints, and corrections to a.debray@math.utexas.edu. The image on the front cover is M.C. Escher's *Circle Limit III* (1959), sourced from <http://www.wikiart.org/en/m-c-escher/circle-limit-iii>.

CONTENTS

1. Review of Complex Analysis: 1/20/16

2

Lecture 1.

Review of Complex Analysis: 1/20/16

Riemann surfaces is a subject that combines the topology of structures with complex analysis: a Riemann surface is a surface endowed with a notion of holomorphic function. This turns out to be an extremely rich idea; it's closely connected to complex analysis but also to algebraic geometry. For example, the data of a compact Riemann surface along with a projective embedding specifies a proper algebraic curve over \mathbb{C} , in the domain of algebraic geometry.¹ In fact, the algebraic geometry course that's currently ongoing is very relevant to this one.

The theory of Riemann surfaces ties into many other domains, some of them quite applied: number theory (via modular forms), symplectic topology (pseudo-holomorphic forms), integrable systems, group theory, and so on: so a very broad range of mathematics graduate students should find it interesting.

Moreover, by comparison with algebraic geometry or the theory of complex manifolds, there's very low overhead; we will quickly be able to write down some quite nontrivial examples and prove some deep theorems: by the middle of the semester, hopefully we will prove the analytic Riemann-Roch theorem, the fundamental theorem on compact Riemann surfaces, and use it to prove a classification theorem, called the uniformization theorem.

The course textbook is S.K. Donaldson's *Riemann Surfaces*, and the course website is at <http://www.ma.utexas.edu/users/perutz/RiemannSurfaces.html>; it currently has notes for this week's material, a rapid review of complex function theory. We will assume a small amount of complex analysis (on the level of Cauchy's theorem; much less than the complex analysis prelim) and topology (specifically, the relationship between the fundamental group and covering spaces). Some experience with calculus on manifolds will be helpful. Some real analysis will be helpful, and midway through the semester there will be a few Hilbert spaces. Thus, though this is a topics course, the demands on your knowledge will more resemble a prelim course.

Let's warm up by (quickly) reviewing basic complex analysis; the notes on the course website will delve into more detail. For the rest of this lecture, G denotes an open set in \mathbb{C} .

The following definition is fundamental.

Definition. A function $f : G \rightarrow \mathbb{C}$ is *holomorphic* if for all $z \in G$, the complex derivative

$$f'(z) = \lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h}$$

exists. The set of holomorphic functions $G \rightarrow \mathbb{C}$ is denoted $\mathcal{O}(G)$, after the Italian *funzione olomorfa*.

¹This sentence is packed with jargon you're not assumed to know yet.

Note that even though it makes sense for the limit to be infinite, this is not allowed.

First, let's establish a few basic properties.

- If $H \subset G$ is open and $f \in \mathcal{O}(G)$, then $f|_H \in \mathcal{O}(H)$.
- The sum, product, quotient, and chain rules hold for holomorphic functions, so $\mathcal{O}(G)$ is a commutative ring (with multiplication given pointwise) and in fact a commutative \mathbb{C} -algebra.²

In other words, holomorphic functions define a *sheaf* of \mathbb{C} -algebras on G .

By a rephrasing of the definition, then if f is holomorphic on G , then it has a *derivative* f' on G , i.e. for all $z \in G$, one can write $f(z+h) = f(z) + f'(z)h + \varepsilon_z(h)$, where $\varepsilon_z(h) \in o(h)$ (that is, $\varepsilon_z(h)/h \rightarrow 0$ as $h \rightarrow 0$). Thus, a holomorphic function is differentiable in the real sense, as a function $G \rightarrow \mathbb{R}^2$. This means that there's an \mathbb{R} -linear map $D_z f : \mathbb{C} \rightarrow \mathbb{C}$ such that $f(z+h) = f(z) + (D_z f)(h) + o(h)$: here, $D_z f(h) = f'(z)h$.

However, we actually know that $D_z f$ is \mathbb{C} -linear. This is known as the *Cauchy-Riemann condition*. Since it's *a priori* \mathbb{R} -linear, saying that it's \mathbb{C} -linear is equivalent to it commuting with multiplication by i . $D_z f$ is represented by the Jacobian matrix

$$D_z f = \begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{bmatrix}.$$

A short calculation shows that this commutes with i iff the following equations, called the *Cauchy-Riemann equations*, hold:

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}. \quad (1.1)$$

The content of this is exactly that $D_z f$ is complex linear.

Conversely, suppose $f : G \rightarrow \mathbb{C}$ is differentiable in the real sense. Then, if it satisfies (1.1), then $D_z f$ is complex linear. But a complex linear map $\mathbb{C} \rightarrow \mathbb{C}$ must be multiplication by a complex number $f'(z)$, so f is holomorphic, with derivative f' .

Power Series. The notation $D(c, R)$ means the open disc centered at c with radius R , i.e. all points $z \in \mathbb{C}$ such that $|z - c| < R$.

Definition. Let $A(z) = \sum_{n=0}^{\infty} a_n(z-c)^n$ be a \mathbb{C} -valued power series centered at a $c \in \mathbb{C}$. Then, its *radius of convergence* is $R = \sup\{|z-c| : A(z) \text{ converges}\}$, which may be 0, a positive real number, or ∞ .

Theorem 1.1. Suppose $A(z) = \sum_{n \geq 0} a_n(z-c)^n$ has radius of convergence R . Then:

- (1) $R^{-1} = \limsup |a_n|^{1/n}$;
- (2) $A(z)$ converges absolutely on $D(c, R)$ to a function $f(z)$;
- (3) the convergence is uniform on smaller discs $D(c, r)$ for $r < R$;
- (4) the series $B(z) = \sum_{n \geq 1} n a_n(z-c)^{n-1}$ has the same radius of convergence R , so converges on $D(c, R)$ to a function $g(z)$; and
- (5) $f \in \mathcal{O}(D(c, R))$ and $f' = g$.

These aren't extremely hard to prove: the first few rely on various series convergence tests from calculus, though the last one takes some more effort.

Paths and Cauchy's Theorem. By a *path* we mean a continuous and piecewise C^1 map $[a, b] \rightarrow \mathbb{C}$ for some real numbers $a < b$. That is, it breaks up into a finite number of chunks on which it has a continuous derivative. A *loop* is a path γ such that $\gamma(a) = \gamma(b)$.

If γ is a C^1 path in G (so its image is in G) and $f : G \rightarrow \mathbb{C}$ is continuous, we define the *integral*

$$\int_{\gamma} f = \int_a^b f(\gamma(t)) \gamma'(t) dt.$$

This is a complex-valued function, because the rightmost integral has real and imaginary parts. This makes sense as a Riemann integral, because these real and imaginary parts are continuous. This is additive on the join of paths, so we can extend the definition to piecewise C^1 paths. Moreover, integrals behave the expected way under reparameterization, and so on.

²A \mathbb{C} -algebra is a commutative ring A with an injective map $\mathbb{C} \hookrightarrow A$, which in this case is the constant functions.

Theorem 1.2 (Fundamental theorem of calculus). *If $F \in \mathcal{O}(G)$ and $\gamma : [a, b] \rightarrow G$ is a path, then*

$$\int_{\gamma} F'(z) \, dz = F(\gamma(b)) - F(\gamma(a)).$$

This is easy to deduce from the standard fundamental theorem of calculus. In particular, if γ is a loop, then the integral of a holomorphic function is 0.

Now, an extremely important theorem.

Definition. A *star-domain* is an open set $G \subset \mathbb{C}$ with a $z^* \in G$ such that for all $z \in G$, the line segment $[z^*, z]$ joining z^* and z is contained in G .

For example, any convex set is a star-domain.

Theorem 1.3 (Cauchy). *If G is a star-domain, γ is a loop in G , and $f \in \mathcal{O}(G)$, then $\int_{\gamma} f = 0$. Indeed, $f = F'$, where*

$$F(z) = \int_{[z^*, z]} f.$$

The proof is in the notes, but the point is that you can check that this definition of F produces a holomorphic function whose derivative is f ; then, you get the result. The idea is to compare $F(z+h)$ and $F(z)$ should be comparable, which depends on an explicit calculation of an integral of a holomorphic function around a triangle, which is not hard.

Cauchy didn't prove Cauchy's theorem this way; instead, he proved Green's theorem, using the Cauchy-Riemann equations. This is short and satisfying, but requires assuming that all holomorphic functions are C^1 . This is true (which is great), but the standard (and easiest) way to show this is... Cauchy's theorem.