

M392C NOTES: RATIONAL HOMOTOPY THEORY

ARUN DEBRAY
SEPTEMBER 17, 2015

These notes were taken in UT Austin's Math 392C (rational homotopy theory) class in Fall 2015, taught by Jonathan Campbell. I live- \TeX ed them using vim, and as such there may be typos; please send questions, comments, complaints, and corrections to a.debray@math.utexas.edu.

CONTENTS

1.	Postnikov Towers and Principal Fibrations: 8/27/15	1
2.	Serre Theory: 9/1/15	4
3.	Rational Homotopy Groups of Spheres: 9/3/15	7
4.	Commutative Differential Graded \mathbb{Q} -algebras and Model Categories: 9/8/15	9
5.	Homotopy in Model Categories: 9/15/15	13
6.	Today is the Cofibrantly Generated Model Categories Day: 9/17/15	16

Lecture 1.

Postnikov Towers and Principal Fibrations: 8/27/15

First, we'll outline some aspects of the course.

$X \rightarrow Y$ is a rational equivalence if $\pi_*(X) \otimes \mathbb{Q} \xrightarrow{\cong} \pi_*(Y) \otimes \mathbb{Q}$. The goal is to define a category $\text{Ho}(\text{Top}_{\mathbb{Q}})$ where, more or less, the isomorphisms are rational equivalences. The point is that this is a purely algebraic category, equivalent to a category of differential graded algebras, $\text{Ho}(\text{CDGA}_{\mathbb{Q}})$.

The first half of the course will deal with something called Sullivan's method: we'll get our hands on rational equivalence, and produce the rationalization functor $X \mapsto X_{\mathbb{Q}}$. We're developing it as it "could have been done," with some computations to show that things get a lot easier over \mathbb{Q} (e.g. homology of Eilenberg-MacLane spaces is the same as for spheres).

Then, we'll have to talk about model categories, which is a good way of producing homotopy categories or homotopy theories for more than just topological categories. Intuitively, a model category is a category in which one can do homotopy theory. Using this, we'll talk about the homotopy theory of commutative, differential algebras over \mathbb{Q} .

This isn't how it was originally done by Sullivan et al., and so we'll also discuss the classical construction. We'll also produce functors from simplicial sets to differential graded \mathbb{Q} -algebras and topological spaces, with adjoints and so on. One of these, turning a simplicial set into a differential graded \mathbb{Q} -algebra, will resemble the functor Ω^* of differential forms, but is more combinatorial.

This will enable us to prove equivalence, with all sorts of cool consequences: Whitehead products appear in the differential graded algebras category; automorphisms of CDGAs correspond to automorphisms of $\text{Top}_{\mathbb{Q}}$, which relate to automorphisms of topological spaces nicely, and so on.

The rest of the course will discuss Quillen's model, which relates differential graded Lie algebras to rational spaces. That might not mean anything right now, and we'll have to learn a little more machinery for it. Thus, this course will cover some classical and some modern algebraic topology, making the useful notion of model categories nice and concrete.

Here are some good references for this subject.

- Griffiths-Morgan, *Rational Homotopy Theory and Differential Forms*; it's all right, and geometric (they use simplicial complexes, rather than simplicial sets, and therefore don't get as nice of a result). The

second edition came out a year ago, but is similar to the first edition. The beginning has a beautiful exposition of algebraic topology in general.

- There's a GTM by Felix, Halperin, and Thomas, called *Rational Homotopy Theory*. It's pretty beefy, and the one gripe the professor has is that it doesn't use model categories at all, making things opaque. But there's definitely a bootlegged copy...
- Katherine Hess, who is a great writer, has a survey paper, about 20 pages, called *Rational Homotopy Theory*.

Those are the only expository works, but there are also some papers.

- Sullivan, "Infinitesimal Computations in Algebraic Topology." Sullivan is crazy, and the paper is very hard to read. Hopefully after the course everything is easier to read.
- Quillen, "Rational Homotopy Theory." This paper also isn't that easy to read.

There are a few other sources; things will be well cited in this class.

Now for some math.

Definition. Let X be a connected topological space. A *Postnikov tower* is a sequence

$$\begin{array}{c} X_n \\ \downarrow \\ \vdots \\ \downarrow \\ X_2 \\ \downarrow \\ X \longrightarrow X_1 \end{array}$$

such that

- (1) there are maps $X \rightarrow X_i$,
- (2) $\pi_i(X) \cong \pi_i(X_n)$ for $i \leq n$, and
- (3) $\pi_i(X_n) = 0$ for $i > n$.

As a consequence of the three properties, the homotopy fiber $X_n \rightarrow X_{n-1}$ is a $K(\pi_n(X), n)$, i.e. an Eilenberg-MacLane space. In some sense, this is a "co-cellular" way of building a space out of Eilenberg-MacLane spaces.

Theorem 1.1. *Postnikov towers exist.*

The proof is easy: just attach cells to X to kill homotopy above a given degree. But that's not so useful of a characterization. We want to know: what information in stage n determines stuff in stage $(n+1)$?

To produce spaces with certain fibers, classifying maps are useful. Suppose X_{n+1} arises as a (homotopy) pullback: if \star denotes a contractible space, this would look like

$$\begin{array}{ccc} X_{n+1} & \longrightarrow & \star \\ \downarrow & & \downarrow \\ X_n & \longrightarrow & K(\pi_{n+1}(X), n+2). \end{array}$$

It would be nice if fibrations with fiber $K(G, n)$ were classified by maps $X \rightarrow K(G, n+1)$, because then we could work with the cohomology group $H^{n+2}(X, \pi_{n+1}(X))$. It's not very easy to compute stuff in this cohomology group, however.

In any case, not every fibration is even classified in this manner!

Definition. A fibration $K(\pi, n) \rightarrow E \rightarrow B$ is *principal* if it arises as a pullback of a path fibration as follows.

$$\begin{array}{ccc} K(\pi, n) & \xlongequal{\quad} & K(\pi, n) \\ \downarrow & & \downarrow \\ E & \longrightarrow & P \\ \downarrow & & \downarrow \\ B & \longrightarrow & K(\pi, n+1) \end{array}$$

There's an equivalent, less useful, formulation in the lecture notes. The reason we like our formulation is the following theorem.

Theorem 1.2. *A connected CW complex X with $\pi_1(X)$ acting trivially on $\pi_n(X)$ has a Postnikov tower composed of principal fibrations.*

As a consequence, X_{n+1} is determined from X_n by a map $k_n : X_n \rightarrow K(\pi_{n+1}(X), n+2)$; this determines a class $[k_n] \in H^{n+2}(X_n, \pi_{n+1}(X))$, called a *k-invariant*. This is why we care about Postnikov towers: they are built up nicely in stages, using cohomology classes that, in nice cases, we can compute. And so in rational homotopy theory, where the homotopy groups are nicer, the *k-invariants* are nicer.

We'll use spectral sequences in this class; an introduction to them can be found in the professor's lecture notes.

Another takeaway from these results is that Eilenberg-MacLane spaces are pretty fundamental building blocks. Though they have nice homotopy, their cohomology groups are generally pretty nasty, leading to computations called Steenrod operations. But rationally, there's a nice result.

Theorem 1.3.

$$H^*(K(\mathbb{Z}, n); \mathbb{Q}) = \begin{cases} \mathbb{Q}[x], & n \text{ even} \\ \Lambda_{\mathbb{Q}}(x), & n \text{ odd}, \end{cases},$$

where the generators x have degree n .

These are the simplest differential graded \mathbb{Q} -algebras, and suggest that all spaces' rational homotopy will be built out of them (which is true).

Proof. As a base case, $H^*(K(\mathbb{Z}, 1); \mathbb{Q}) = H^*(S^1; \mathbb{Q}) = \Lambda_{\mathbb{Q}}(x)$, which is fine. More generally, we'll use the fibration

$$K(\mathbb{Z}, n-1) \longrightarrow \star \longrightarrow K(\mathbb{Z}, n).$$

By induction, if n is odd, then $H^*(K(\mathbb{Z}, n-1); \mathbb{Q}) = \mathbb{Q}[x]$, with $\deg x = n-1$, since $n-1$ is even. Let's use the Serre spectral sequence, for which

$$E_2^{p,q} = H^p(K(\mathbb{Z}, n-1); \mathbb{Q}) \otimes H^q(K(\mathbb{Z}, n-1); \mathbb{Q}).$$

For example, when $n = 3$, we have

$$\begin{array}{cccccccc} \mathbb{Q}y^3 & & & & & & & \\ 0 & & & & & & & \\ \mathbb{Q}y^2 & & & & & & & \\ 0 & & & & & & & \\ \mathbb{Q}y & 0 & 0 & \mathbb{Q}x \otimes \mathbb{Q}y & & & & \\ 0 & 0 & 0 & 0 & & & & \\ 0 & 0 & 0 & \mathbb{Q}x & H^4 & H^5 & \dots & \end{array}$$

Here, degree increases from 0 to the right and going upwards. This also uses the Hurevitch theorem. Then, we remark that d_2 , from $(0, 2)$ to $(3, 0)$, has to be an isomorphism, because the E_∞ -page is 0.¹ But since the Serre spectral sequence is linear, we also have isomorphisms from $(4, 0)$ to $(2, 2)$ to $(0, 4)$; specifically, $d_3: \mathbb{Q}y^2 \rightarrow \mathbb{Q}x \otimes \mathbb{Q}y$, so $H^6(K(\mathbb{Z}, 3); \mathbb{Q}) = 0$. And this means that $x^2 = 0$. Then, we continue by induction to show that all higher $H^q(K(\mathbb{Z}, 3); \mathbb{Q})$ are zero. Thus, $H^*(K(\mathbb{Z}, 3); \mathbb{Q}) = \Lambda_{\mathbb{Q}}(x)$, and the case for general n is similar. \square

¹Though the first few lectures will use spectral sequences, they won't be very important after that, so don't drop the course if this is the only thing making you uncomfortable.

Exercise 1. Handle the case where n is even, which is somewhat similar.

Again, this is suggestive: Eilenberg-MacLane spaces build topological spaces up, and they have differential graded algebras for their rational cohomology groups.

Next lecture, we'll discuss Serre theory, the tricks that Serre used to compute the rational homotopy groups of the spheres. These are strong clues that, rationally, things are much nicer.

After that, we'll discuss rational equivalence, and then CDGAs and their homotopy theory, necessitating a discussion of model categories. The course will get less computational at this point.

Lecture 2.

Serre Theory: 9/1/15

"Whistle guy has really got me off my game!"

Last lecture may have gone a little fast, so we'll reboot and say some things that we didn't mean to assume. Then, we'll start Serre theory.

Definition. If G is a group, an *Eilenberg-MacLane space* $K(G, n)$ is a space whose homotopy groups are

$$\pi_i(K(G, n)) = \begin{cases} G, & i = n \\ 0, & i \neq n. \end{cases}$$

For example, S^1 is a $K(\mathbb{Z}, 1)$, $\mathbb{C}P^\infty$ is a $K(\mathbb{Z}, 2)$, and $\mathbb{R}P^\infty$ is a $K(\mathbb{Z}/2, 1)$. Also, $K(G, n-1) = \Omega K(G, n)$ (i.e. the space of loops). One can think of building spaces out of these, because they're the simplest spaces from the perspective of homotopy theory.

Another of the reasons they're important is the following theorem, which is hard to prove.

Theorem 2.1. $[X, K(G, n)] = \tilde{H}^n(X; G)$.

In particular, the cohomology functor is representable.

This is why, last time, a map $k : X \rightarrow K(\pi_{n+1}(X), n+2)$ corresponded to $H^{n+2}(X; \pi_{n+1}(X))$: k -invariants arise from the representability of cohomology.

Postnikov towers are a way of using Eilenberg-MacLane spaces to build a space up, one homotopy group at a time.

Theorem 2.2. *Eilenberg-MacLane spaces exist for all n and all G , where G is abelian if $n > 1$.*

Recall by the Eckmann-Hilton argument that $\pi_n(X)$ is always abelian when $n > 1$.

Proof idea. When $n = 1$, one can produce a fiber bundle where the total bundle is trivial and the fiber is G , with the discrete topology, thus producing a sequence $G \rightarrow EG \rightarrow BG$, and one can show that $\pi_1(BG) = G$ and $\pi_i(BG) = 0$ for $i \geq 2$. This is done in chapter 1 of Hatcher.

If $n > 1$, take a resolution for G :

$$\mathbb{Z}[r_\beta] \longrightarrow \mathbb{Z}[g_\alpha] \longrightarrow G,$$

where the g_α are generators and r_β are relations. Then, consider a bouquet of spheres $\bigvee_\alpha S_\alpha^n$; each relation gives a map $S^n \rightarrow \bigvee_\alpha S_\alpha^n$ using the degree of the relation, so glue cells onto this bouquet via the relations, forming pushouts of the form

$$\begin{array}{ccc} \bigvee_\beta S^n & \longrightarrow & \bigvee_\alpha S_\alpha^n \\ \downarrow & & \uparrow \\ \bigvee_\beta D^n & \longrightarrow & X \end{array}$$

Then, the n -skeleton $X^{(n)}$ is given by the generators, and so we have an exact sequence

$$\pi_{n+1}(X, X^{(n)}) \longrightarrow \pi_n(X^{(n)}) \longrightarrow \pi_n(X) \longrightarrow \pi_n(X, X^{(n)}).$$

Then, $\pi_{n+1}(X, X^{(n)}) = \mathbb{Z}[r_\beta]$, $\pi_n(X^{(n)}) = \mathbb{Z}[g_\alpha]$, and $\pi_n(X, X^{(n)}) = 0$, so $\pi_n(X) = G$. There are only n -cells, so there's no lower homotopy, and we can kill all higher homotopy in a standardized manner. \square

Anyways, last time we talked about how $H^*(K(\mathbb{Z}, n), \mathbb{Q})$ is $\mathbb{Q}[x]$ when n is even and $\Lambda_{\mathbb{Q}}(x)$ when n is odd; this suggests that since the constituents of spaces have simple rational homotopy, then maybe rational homotopy provides some insights.

We proved this with the Serre spectral sequence, and we'll use it a few more times to get nice short exact sequences, so let's carefully state what's going on.

Theorem 2.3 (Homological Serre sequence). *Let $F \rightarrow E \rightarrow B$ be a fibration with $\pi_1(B)$ acting trivially on $H^*(F, G)$. Then, there is a spectral sequence $\{E_{p,q}^r, d_r\}$ such that:*

- (1) $d_r : E_{p,q}^r \rightarrow E_{p-r, q-r+1}^r$, i.e. d_r is of degree $(-r, r-1)$.
- (2) $E_{n, n-p}^\infty = F_n^p / F_n^{p-1}$, where F_n^\bullet is some filtration of $H_n(E; G)$, i.e. $H_n(E; G) = F_n^n \supset F_n^{n-1} \supset \dots \supset F_n^0 = 0$.
- (3) $E_{p,q}^2 = H_p(B; H_q(F; G))$.

Remark. If G is a field, then Künneth gives us the nicer result that $E_{p,q}^2 = H_p(B) \otimes H_q(E)$.

Theorem 2.4 (Cohomological Serre sequence). *With the same setup as Theorem 2.3, there exists a spectral sequence $(E_r^{p,q}, d_r)$ such that:*

- (1) $d_r : E_r^{p,q} \rightarrow E_r^{p+q, q-r+1}$.
- (2) $E_\infty^{p, n-p} = F_p^n / F_{p-1}^n$, where F^\bullet is some filtration of $H^n(E; G)$.
- (3) $E_2^{p,q} \cong H^p(B; H^q(F; G))$.

The cohomological Serre sequence is multiplicative: there's a product that ultimately comes from the cup product. This lends some rigidity to the cohomological theory that is often very useful.

Example 2.5. Suppose we're in a case of the homological Serre spectral sequence such that $E_{p,q}^2$ for $0 < q < n$. It turns out that in situations like this, you can leverage your knowledge of the sequence to obtain useful exact sequences.

In this case, the first differential that does anything interesting is $d_{n+1} : E_{n+1,0}^2 \rightarrow E_{0,n}^2$ (previous differentials all map to zero). This means that $E_{n+1,0}^3 = E_{n+1,0}^2 / \text{Im}(d_{n+1})$, and nothing else hits this, so this is also $E_{n+1,0}^\infty$. Thus, we have a sequence

$$E_{n+1,0}^2 \longrightarrow E_{0,n}^2 \longrightarrow E_{0,n}^\infty \longrightarrow 0. \quad (2.1)$$

Furthermore, there is a filtration

$$H_n(E) = F_n^n \supset F_n^{n-1} \supset \dots \supset F_n^0,$$

with $E_{p, n-p}^\infty = F_p^p / F_p^{p-1}$. In particular, $E_{0,n}^\infty = F_n^0$, $F_{1,n}^\infty = 0$, and $E_{n,0}^\infty = F_n^n / F_n^{n-1} = H_n(X) / E_{0,n}^\infty$. That is, we have a sequence

$$0 \longrightarrow E_{0,n}^\infty \longrightarrow H_n(X) \longrightarrow E_{n,0}^\infty \longrightarrow 0,$$

which we can join to (2.1) to produce

$$E_{n+1,0}^2 \longrightarrow E_{0,n}^2 \longrightarrow E_{0,n}^\infty \longrightarrow H_n(E) \longrightarrow E_{n,0}^\infty \longrightarrow 0.$$

This may seem a little contrived, but it happens, for example, when a fiber in a fibration is n -connected.

Serre Theory. This was sometimes called Serre's thesis. It will use some very abstract computations to determine the rational homotopy groups of the spheres.

Definition. Let \mathcal{C} be one of the three classes (the *Serre classes*): FG of finitely generated abelian groups, \mathcal{T}_P , the torsion abelian groups with orders drawn from a set P , and \mathcal{F}_P , the finite groups in \mathcal{T}_P .

Lemma 2.6. *The classes \mathcal{C} are closed under extension: that is, if $A, C \in \mathcal{C}$ and $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is short exact, then $B \in \mathcal{C}$. Moreover, for any $A, B \in \mathcal{C}$, $A \otimes B \in \mathcal{C}$ and $\text{Tor}(A, B) \in \mathcal{C}$.*

The point is that the disgusting machinery of the Serre spectral sequence mostly leaves a Serre class intact, which will be useful for proving the Hurewicz theorem mod \mathcal{C} . First, though, we'll need more lemmas.

Lemma 2.7. *Let $F \rightarrow E \rightarrow B$ be a fibration satisfying the hypotheses of the homological Serre spectral sequence, and assume F , E , and B are all path-connected. If any two of $H_*(B)$, $H_*(E)$, and $H_*(F)$ are in \mathcal{C} , then so is the third.*

Proof. We'll show that if $H_*(B)$ and $H_*(F)$ are in \mathcal{C} , then $H_*(E)$ is. Recall that

$$\begin{aligned} E_{p,q}^2 &= H_p(B; H_q(F)) \\ &= H_p(B; \mathbb{Z}) \otimes H_q(F; \mathbb{Z}) \oplus \text{Tor}(H_{p-1}(B), H_q(F)), \end{aligned}$$

by the universal coefficient theorem. The first two terms are in \mathcal{C} by assumption, and Lemma 2.6 implies the last one is. Thus, $E_{p,q}^2 \in \mathcal{C}$, so all subsequent pages must be too (since homology is just kernels and images, which don't pop us out of \mathcal{C}). Since $H_n(X)$ has successive filtrations whose quotients are $E_{p,q}^\infty$, which are all in \mathcal{C} , then $H_n(E) \in \mathcal{C}$. \square

Lemma 2.8. *If $\pi \in \mathcal{C}$, then $H_k(K(\pi, n), \mathbb{Z}) \in \mathcal{C}$ for all k .*

At this point, there aren't many guesses for tools we can use: the Serre sequence is basically the only tool we have for homotopy groups.

Proof. Recall that we have a fibration $K(\pi, n-1) \rightarrow * \rightarrow K(\pi, n)$; we'll apply the Serre sequence. By induction, it's sufficient to consider the case $n = 1$.

For right now, we'll consider $\mathcal{C} = FG$. By Künneth, it's sufficient to show for $K(\mathbb{Z}, 1)$ and $K(\mathbb{Z}/m, 1)$, which are S^1 and lens spaces, for which this is true. \square

Lemma 2.9. *Let X be simply connected. Then, $\pi_n(X) \in \mathcal{C}$ implies $H_n(X) \in \mathcal{C}$.*

Remark. The converse also holds (i.e. if we know this for all n , then X must be simply connected), but we won't show that yet.

Proof of Lemma 2.9. We'll use a Postnikov tower.

$$\begin{array}{ccc} K(\pi_n(X), n) & \longrightarrow & X_n \\ & & \downarrow \\ & & X_{n-1} \end{array} \tag{2.2}$$

From Lemma 2.8, we know that $H_k(K(\pi_n(x), n), \mathbb{Z}) \in \mathcal{C}$, so once again using the homological Serre sequence, $H_*(X_n)$ is computed using $E_{p,q}^2 = H_p(X_{n-1}, H_q(K(\pi_n(x), n)))$, and therefore $H_n(X_n; \mathbb{Z}) \in \mathcal{C}$. But this is $H_n(X)$. \square

There seems to be a duality, where things with complicated homology seem to have uncomplicated homotopy, and vice versa.

Now we can get to the reason we're doing this abstract nonsense.

In the following theorems, *isomorphic mod \mathcal{C}* means that the kernel and cokernel of the map are both in \mathcal{C} .

Theorem 2.10 (Mod \mathcal{C} Hurewicz). *If X has $\pi_i(X) \in \mathcal{C}$ for $i < n$, then $h : \pi_n(X) \rightarrow H_n(X)$ is an isomorphism mod \mathcal{C} .*

Corollary 2.11. *If $\pi_i(X) \in \mathcal{C}$ for all i , then $\pi_n(X) \rightarrow H_n(X)$ is isomorphic mod \mathcal{C} for all i .*

Corollary 2.12. *If $\pi_i(X) \in FG$, then $\pi_i(X) \otimes \mathbb{Q} \rightarrow H_i(X) \otimes \mathbb{Q}$ is an isomorphism.*

Proof of Theorem 2.10. $\pi_n(X) \rightarrow H_n(X)$ is the same as $\pi_n(X_n) \rightarrow H_n(X_n)$, where $\{X_i\}$ is the Postnikov tower. Look at the fibration (2.2), and use the five-term exact sequence

$$0 \longrightarrow H_{n+1}(X_{n-1}) \longrightarrow H_n(K(\pi_n(X), n)) \longrightarrow E_{0,n}^\infty \longrightarrow H_n(X_n) \longrightarrow H_n(X_{n-1}).$$

The composition of the middle maps is the inclusion of the fiber, so we get a four-term exact sequence

$$0 \longrightarrow H_{n+1}(X_{n-1}) \longrightarrow H_n(K(\pi_n(X), n)) \longrightarrow H_n(X_n) \longrightarrow H_n(X_{n-1}) \longrightarrow 0.$$

By induction, the first and last terms are in \mathcal{C} , which is exactly what we need for the middle arrow to be an isomorphism mod \mathcal{C} .

Now, we can put that into the following diagram.

$$\begin{array}{ccc} H_n(K(\pi_n(X), n)) & \xrightarrow{\cong} & \pi_n(X_n) \\ \downarrow h \cong & & \downarrow \\ H_n(K(\pi_n(X), n)) & \xrightarrow{\cong \text{ mod } \mathcal{C}} & H_n(X_n) \end{array}$$

The isomorphism on the left follows from the usual Hurewicz theorem, and the one on the top is from the construction of the Postnikov tower. Then, we just showed the result for the bottom arrow, so the result is that the arrow on the right is an isomorphism mod \mathcal{C} . \square

You can see the game: Postnikov tower, fibration, Serre sequence, and then hope and pray that there's an equivalence (e.g. one coarse enough such as \mathcal{C}).

One corollary here is that homology and homotopy don't differ much over \mathbb{Q} , which is pretty nice.

Lecture 3.

Rational Homotopy Groups of Spheres: 9/3/15

"I looked through my notes the other day, and I had at least five spellings of Hurewicz."

Though we'll do some significant computations with rational homotopy groups today, we'll start by clarifying a few things from last time.

Proposition 3.1. *Let X be a space with a Postnikov tower $\{X_i\}$. Then, $H_n(X) = H_n(X_n)$.*

Proof. There's a map $X \rightarrow X_n$; consider it as a fibration, so we get a fiber $X^{>n} \rightarrow X \rightarrow X_n$. So we can either use the long exact sequence of a fibration or the Serre spectral sequence, and the former isn't so useful here.

We have $E_2^{p,q} = H_p(X_n, H_q(X^{>n}, \mathbb{Z}))$. We know $X^{>n}$ doesn't have homotopy in degrees n or lower, and therefore it doesn't have homology there either. This implies that $E_\infty^{0,n} = H_n(X_n)$, and for $p+q=n$ otherwise, $E_\infty^{p,q} = 0$. Since $E_\infty^{p,n-p}$ filters $H_n(X)$ and there's only one nonzero term in the filtration, $H_n(X) = H_n(X_n)$. \square

This is a common technique with spectral sequences: things work because there end up being large gaps. We've even seen it a few times before.

Another loose end from last time is the following corollary of Hurewicz mod \mathcal{C} .

Corollary 3.2. *Suppose that X is a space such that $\pi_1(X)$ acts trivially on X (i.e. X has a Postnikov tower). Then, $H_*(X; \mathbb{Q}) = 0$ iff $\pi_*(X) \otimes \mathbb{Q} = 0$.*

Now, we may compute the rational homotopy groups of spheres. It's pretty amazing that knowledge of the homotopy of Eilenberg-MacLane spaces and the Serre spectral sequence will do it. It's really clever (well of course it is; it's Serre).

We know $\pi_n(S^n) = \mathbb{Z}$, and therefore $\pi_n(S^n) \otimes \mathbb{Q} = \mathbb{Q}$.

Theorem 3.3. $\pi_i(S^n)$ for $i > n$ are finite except when $n = 2k$, in which case $\pi_{4k+1}(S^{2k}) = \mathbb{Z} \oplus A$ for a finite A .

Proof. The whole thing is based on the following: consider a map $S^n \rightarrow K(\mathbb{Z}, n)$ inducing an isomorphism on π_n , i.e., this is the generator of $\pi_n(K(\mathbb{Z}, n)) = \mathbb{Z}$. As usual, we'll convert it into a fibration $F \rightarrow S^n \rightarrow K(\mathbb{Z}, n)$. Repeating, we get a fibration $K(\mathbb{Z}, n-1) \rightarrow F \rightarrow S^n$.² The point is, $\pi_i(F)$ and $\pi_i(S^n)$ agree when $i > n$, and F fits nicely into these two fibrations.

Obviously, we're going to use the Serre spectral sequence on this, but first we need to understand the cohomology of $K(\mathbb{Z}, n-1)$.

When n is odd, $E_{p,q}^2 = H^p(S^n; \mathbb{Q}) \otimes H^q(K(\mathbb{Z}, n-1), \mathbb{Q}) = \Lambda(x) \otimes \mathbb{Q}[y]$, where $\deg x = n$ and $\deg y = n-1$. We computed this in the first lecture. This means that in row 0, there's only a \mathbb{Q} at $(0,0)$ and a $\mathbb{Q}x$ at $(0,n)$; in $n-1$, we have $\mathbb{Q}y$ at column 0 and $\mathbb{Q}xy$ in column n , and then y^2 in place of y at row $2n-2$, and so on.

Since F is $(n-1)$ -connected, then $H^{n-1}(F; \mathbb{Q})$ must vanish. Thus, $\mathbb{Q}y$ has to die, and this can only happen if $d_{n-1}: \mathbb{Q}y \rightarrow \mathbb{Q}x$ is an isomorphism. Then, by multiplicativity, the other differentials have to be isomorphisms: if $y \rightarrow x$, then $\mathbb{Q}y^2 \rightarrow \mathbb{Q}xy$, and so on. Thus, $H^*(F; \mathbb{Q}) = 0$, and therefore that $\pi_i(F) \otimes \mathbb{Q} = 0$ for all i . Thus, $\pi_i(F)$ (and therefore $\pi_i(S^n)$) is finite for $i > n$ when n is odd.

²This is because a fibration fits into a longer sequence $\cdots \rightarrow \Omega E \rightarrow \Omega B \rightarrow F \rightarrow E \rightarrow B$.

When n is even, then E^2 page has only \mathbb{Q} at $(0,0)$, $\mathbb{Q}x$ at $(0,n)$, $\mathbb{Q}y$ at $(n-1,0)$, and $\mathbb{Q}xy$ at $(n-1,n)$. As before, $H^{n-1}(F; \mathbb{Q}) = 0$, so $d_{n-1}\mathbb{Q}y \rightarrow \mathbb{Q}x$ is an isomorphism. That means that $H^*(F; \mathbb{Q}) = H^*(S^{2n-1}; \mathbb{Q})$, i.e. there's a generator at degree $2n-1$, and nothing else. This tells us that $\pi_{2n-1}(S^n) \otimes \mathbb{Q} = \mathbb{Q}$ by Hurewicz, and that everything below it vanishes.

Finally, we have to address $i > 2n-1$, where we want the homotopy to vanish. Let $F \rightarrow F^{<2n-1}$ be obtained by killing all homotopy above $i \geq 2n-1$. Turn this into a fibration and take the fiber to get $F^{\geq 2n-1} \rightarrow F \rightarrow F^{<2n-1}$.

We know $\pi_i(F^{<2n-1})$ is finite, since it agrees with $\pi_i(F)$ on this range, and therefore $H^*(F^{<2n-1}; \mathbb{Q}) = 0$. Using the Serre spectral sequence, we have that $H^*(F^{\geq 2n-1}; \mathbb{Q}) = H^*(F; \mathbb{Q}) = H^*(S^{2n-1}; \mathbb{Q})$. Now, consider $F^{\geq 2n-1} \rightarrow K(\mathbb{Z}, 2n-1)$ which induces an isomorphism mod FG on π_{2n-1} . Once again, we'll take the fiber to get $K(\mathbb{Z}, 2n-2) \rightarrow \tilde{F} \rightarrow F^{\geq 2n-1}$, and as in the previous case, the homotopy of \tilde{F} and of $F^{\geq 2n-1}$ are equal in degrees $2n$ and above.

Once again, we use the Serre spectral sequence: $E_2^{p,q} = H^p(F^{\geq 2n-1}; \mathbb{Q}) \otimes H^q(K(\mathbb{Z}, n-2); \mathbb{Q})$, but the first part of the tensor product is $H^p(S^{2n-1}; \mathbb{Q})$, so we have exactly the same case as when n was odd. Thus, $H^*(\tilde{F}; \mathbb{Q})$ vanishes for the same reason. \square

Though this proof looks frightful, what happened was that we set up the Serre spectral sequence, and then took the one fact we knew, that F is $(n-1)$ -connected, and used it to set up just enough calculations to figure things out.

Again, it's incredible, even though the Serre spectral sequence is a big machine, it's human-understandable (and even admits a really quick proof with simplicial sets), and surprisingly few techniques are needed. We do need that the homotopy groups of the spheres are finitely generated, but this isn't too bad either.

Rational Spaces and Localization. Now that we've done some rational calculations and seen that they're easier than regular ones, we can see that rational equivalence might be a nicer idea (which is true; the category ends up being completely algebraic).

Almost all of the theory from here on out will deal with simply connected spaces. This is because π_1 of non-simply-connected groups might not be abelian, and so we usually can't tensor it with \mathbb{Q} . You can sort of do this with nilpotent groups, but we won't delve into that.

Definition. A simply connected space X is called \mathbb{Q} -local or a \mathbb{Q} -space if

- (1) $\pi_*(X)$ is a \mathbb{Q} -vector space, or
- (2) $H_*(X; \mathbb{Z})$ is a \mathbb{Q} -vector space.

Proposition 3.4. *The above two criteria are equivalent.*

Proof. Once again, we'll use Postnikov towers and the Serre spectral sequence. It's almost boring, but it's also interesting how useful they are.

For homology implying homotopy, start induction at $X_2 = K(\pi_2(X), 2)$. We know $H_*(X_2, \mathbb{Z})$ is a \mathbb{Q} -vector space, by computation of $H^*(K(\mathbb{Q}, n); \mathbb{Z})$.

The inductive step looks at the fibration

$$\begin{array}{ccc} K(\pi_n(X), n) & \longrightarrow & X_n \\ & & \downarrow \\ & & X_{n-1}. \end{array}$$

Then, $E_2^{p,q} = H_p(X_{n-1}; H_q(K(\pi_n(X), n)))$, but all of these are \mathbb{Q} -vector spaces, and therefore $H_*(X_n)$ is a \mathbb{Q} -vector space; then, take the limit.

The other direction isn't any more interesting; we assume $H_*(X; \mathbb{Z})$ is a \mathbb{Q} -vector space. Suppose we know that $\pi_*(X_{n-1})$ is a \mathbb{Q} -vector space. Then, $H_*(X_{n-1}; \mathbb{Z})$ is, and so is $H_*(X, X_{n-1})$, and $\pi_n(X) = H_{n+1}(X, X_{n-1})$. \square

This is what we mean by \mathbb{Q} -local. In other words, there's only field-level information, and there's no torsion.

Definition. For "nice" topological spaces X and Y , we say that X and Y are \mathbb{Q} -equivalent if there's a map $X \rightarrow Y$ that induces either

- (1) $H_*(X; \mathbb{Q}) \xrightarrow{\sim} H_*(Y; \mathbb{Q})$ or
- (2) $\pi_*(X) \otimes \mathbb{Q} \xrightarrow{\sim} \pi_*(Y) \otimes \mathbb{Q}$.

Equivalently, the fiber of $X \rightarrow Y$ is \mathbb{Q} -trivial (also called \mathbb{Q} -acyclic, meaning it has no rational homology or homotopy).

We'll want to introduce the notion of a localization, producing a map from a space into a simpler topological space that preserves all the information over \mathbb{Q} .

Definition. A map $j : X \rightarrow X_{\mathbb{Q}}$ with $X_{\mathbb{Q}}$ a \mathbb{Q} -local space is a *localization* if one of the following is true:

- (1) $j_* : H_*(X; \mathbb{Q}) \rightarrow H_*(X_{\mathbb{Q}}; \mathbb{Q})$ is an isomorphism;
- (2) $j_* : \pi_*(X) \otimes \mathbb{Q} \rightarrow \pi_*(X_{\mathbb{Q}}) \otimes \mathbb{Q}$ is an isomorphism; or
- (3) j is universal in that if Y is \mathbb{Q} -local and $f : X \rightarrow Y$ is continuous, then there exists a unique $f' : X_{\mathbb{Q}} \rightarrow Y$ such that the following diagram commutes.

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow j & \nearrow f' \\ & X_{\mathbb{Q}} & \end{array}$$

We'll be able to construct a localization for any X by inducting on a Postnikov tower, tensoring everything with \mathbb{Q} .

Lemma 3.5. Let G be an abelian group and $K(G, n)$ be an Eilenberg-MacLane space for it. Then, its localization is $K(G, n) \rightarrow K(G \otimes \mathbb{Q}, n)$.

Proof. Follows from the definitions. ⊠

Theorem 3.6. If X is a simply connected topological space, then there exists a localization $X \rightarrow X_{\mathbb{Q}}$.

Proof. We'll induct on Postnikov towers; assume we've created a localization $X_{n-1} \rightarrow (X_{n-1})_{\mathbb{Q}}$, and we'd like to construct an $(X_n)_{\mathbb{Q}}$. How do we do that? We know how to construct X_n out of X_{n-1} : it fits into a homotopy fiber square

$$\begin{array}{ccc} X_n & & \\ \downarrow & & \\ X_{n-1} & \xrightarrow{k_n} & K(\pi_n(X), n+1), \end{array}$$

where k_n is a k -invariant. Compose this map with our localization of $K(\pi_n(X), n+1)$, so by the universal property of localizations, we get maps $X_{n-1} \rightarrow (X_{n-1})_{\mathbb{Q}} \rightarrow K(\pi_n(X) \otimes \mathbb{Q}, n+1)$; then, this pulls back up the fiber diagram, giving us an $(X_n)_{\mathbb{Q}}$. ⊠

At the start of the next lecture, knowing that localization of spheres exists produces a much more conceptual calculation of the homotopy groups of spheres.

Example 3.7. Chern classes give us maps $\varphi : BU \rightarrow \prod_k K(\mathbb{Z}, 2k)$; representability says that a map $BU \rightarrow K(\mathbb{Z}, 2k)$ is equivalent to a class in $H^{2k}(BU)$. Thus, computation of $H^*(BU, \mathbb{Q})$ tells us that φ^* is an isomorphism on cohomology, and therefore $BU_{\mathbb{Q}} \simeq \prod_k K(\mathbb{Q}, 2k)$. In particular, $\Omega^2 BU_{\mathbb{Q}} \simeq BU_{\mathbb{Q}}$. In other words, this is rational Bott periodicity.

Lecture 4.

Commutative Differential Graded \mathbb{Q} -algebras and Model Categories: 9/8/15

First, we'll talk a little bit about localization; then, we'll move to something completely different.

Proposition 4.1. $S_{\mathbb{Q}}^{2k+1}$ is a $K(\mathbb{Q}, 2k+1)$, and there's a fibration $K(\mathbb{Q}, 4k-1) \rightarrow S_{\mathbb{Q}}^{2k} \rightarrow K(\mathbb{Q}, 2k)$.

Once again, we have an even and an odd case.

Proof. $S_{\mathbb{Q}}^{2k+1} \rightarrow K(\mathbb{Q}, 2k+1)$ induces an isomorphism on homology, and therefore rational homotopy.

For the fibration, consider $S_{\mathbb{Q}}^{2k} \rightarrow K(\mathbb{Q}, 2k)$ acting on $H_{2k}(-)$. Then, we get a fiber $F \rightarrow S_{\mathbb{Q}}^{2k} \rightarrow K(\mathbb{Q}, 2k)$; to figure out what the fiber is, we'll use the Serre spectral sequence.

$$\begin{array}{c|cccc} 4k-1 & \mathbb{Q} & \mathbb{Q} & & \\ \hline & \mathbb{Q} & \mathbb{Q} & \mathbb{Q} & \mathbb{Q} \\ \hline & 0 & 2k & 4k & 6k \end{array}$$

Then, $E_{p,q}^2 = H^*(F) \otimes H^*(K(\mathbb{Q}, 2k))$, so $H^*(F; \mathbb{Q}) = H^*(S_{\mathbb{Q}}^{4k-1})$. That is, F is an $M(\mathbb{Q}, 4k-1)$, but Moore spaces and Eilenberg-MacLane spaces are the same in rational homotopy. \square

Remark. A Moore space is the homology analogue of an Eilenberg-MacLane space; an $M(G, n)$ has homology equal to G in degree n and 0 otherwise.

Right now, we have lots of nice notions of equivalence in topology, including homotopy equivalence, weak homotopy equivalence, and rational homotopy equivalence. We don't have that going on for commutative differential graded algebras. The reason to introduce the abstraction of model categories is to allow for very natural notions of homotopy in categories such as this one.

Definition. A commutative differential graded algebra is a graded algebra A^* over a field k , i.e. $A^* = \bigoplus_{p \geq 0} A^p$ with

- (1) a differential $d : A^* \rightarrow A^{*+1}$ with $d^2 = 0$ and $d(ab) = da \cdot b + (-1)^{\deg a} a \cdot db$, and
- (2) a multiplication $A^p \otimes A^q \rightarrow A^{p+q}$ such that $ab = (-1)^{\deg a \deg b} ba$.

You could formalize this as a chain complex, but we've chosen cochain complexes. The idea is that the multiplication and differential should both respect the grading, and otherwise it's exactly what the name suggests. Moreover, a morphism of CDGAs is a morphism of k -algebras commuting with the differential and grading. Thus, we get a category of k -CDGAs.

In our case, of course, we'll usually take $k = \mathbb{Q}$. In this case, we'll more or less get a categorical equivalence between this category and the category of rational homotopy types, which is incredible!

Example 4.2.

- (1) An important example over \mathbb{R} is $\Omega^*(X; \mathbb{R})$ when X is a manifold, the algebra of differential forms. Here, multiplication is given by the wedge product.
- (2) Over \mathbb{Q} , we could take $\mathbb{Q}[x] \otimes \Lambda(y)$, where $dx = y$, $\deg x = 1$, and $\deg y = 2$. Then, $d(x^n y) = 0$, which you can check, and that $H^*(-) = k$ (i.e. taken with d), and this field k has degree 0. This algebra is, in some mysterious sense we'll clarify, "the interval," notice at least that it has similar cohomology.
- (3) The Koszul complex $K(x_1, \dots, x_n) = \Lambda(x_1) \otimes \dots \otimes \Lambda(x_n)$, with a complicated mess of differentials.
- (4) Another great example, but one that is important, is cohomology $H^*(X; \mathbb{Q})$ with trivial differential. This is important because equivalences between CDGAs will be quasi-isomorphisms, which can send nontrivial differentials to trivial ones. Things quasi-isomorphic to CDGAs with trivial differential are called *formal*, and include important classes such as the rational homology of Kähler manifolds.

A significant non-example is $C^*(X; \mathbb{Q})$. This is because multiplication isn't commutative. In some cases, e.g. finite fields, Steenrod operations have nice structure, and one generally hopes that cochains are a good representative for your space, containing lots of information about your space; so it would've been nice if there was a commutative structure akin to $\Omega^*(M; \mathbb{R})$ for manifolds M . Thus, it doesn't work, but its failure to work is motivational.

Moreover, there isn't really a "free" CDGA, only semi-free ones; the issue is that you cannot freely choose the differential. But given a graded vector space V^* , the semi-free CDGA is

$$F(V^*) = k[V_{\text{even}}] \otimes \Lambda(V_{\text{odd}}).$$

We have a forgetful functor from CDGAs over \mathbb{Q} to \mathbb{Q} -chains and then to \mathbb{Q} -graded vector spaces; this isn't adjoint to the latter, but to the former. Note that the category of CDGAs is probably abelian; certainly, it has kernels, and all of the structure that we need.

We implicitly used the cohomology in Example 4.2; here's the explicit definition.

Definition.

- The *cohomology* of A^\bullet , denoted $H^*(A^\bullet)$, is defined by $H^n(A^\bullet) = (\ker d : A^n \rightarrow A^{n+1}) / (\operatorname{Im} d : A^{n-1} \rightarrow A^n)$.
- Let C^\bullet and D^\bullet be CDGAs, with a morphism $f : C^\bullet \rightarrow D^\bullet$, and define $M_f^n = C^n \oplus D^{n-1}$ with differential $d_M : M_f^n \rightarrow M_f^{n+1}$ given by $\begin{pmatrix} d_C & 0 \\ f & -d_D \end{pmatrix}$. Then, the *relative cohomology* is $H^*(D^\bullet, C^\bullet) = H^*(M_f^\bullet)$.

Notice that we have things like mapping cylinders, relative cohomology, etc., so CDGAs do look somewhat like topological spaces, which will be nice.

Lemma 4.3. *There exists a long exact sequence*

$$\cdots \longrightarrow H^n(C^\bullet) \longrightarrow H^n(D^\bullet) \longrightarrow H^{n+1}(D^\bullet, C^\bullet) \longrightarrow H^{n+1}(C^\bullet) \longrightarrow \cdots$$

Definition. $f : C^\bullet \rightarrow D^\bullet$ is a *quasi-isomorphism* if $H^*(f)$ is an isomorphism.

What we'll do now is spend some time codifying what it means for a category to have homotopy theory; that both rational homotopy and CDGAs live in the same world. This is because both topological spaces and CDGAs are model categories.

If you do much more homotopy theory, model categories will be really useful. Even where they're not explicitly talked about, they'll clarify ideas and thoughts in papers and books.

A model category can be thought of as a model for homotopy; in such a category, there will be things that act like cylinders (or, dually, paths), and therefore we can speak of when objects are homotopic. This will follow from some impressively minimal axioms, because Quillen was extremely clever. We'll define the category, and then show that these induce notions of homotopy equivalence and weak equivalence, and some correspondences between the two.

First, recall a category-theoretic definition.

Definition. Let $f : A \rightarrow B$ and $g : A' \rightarrow B'$ be morphisms in a category \mathcal{C} . Then, f is said to be a *retract* of g if the following diagram commutes.

$$\begin{array}{ccccc} & & \text{id} & & \\ & \curvearrowright & & \curvearrowleft & \\ A & \longrightarrow & A' & \longrightarrow & A \\ \downarrow f & & \downarrow g & & \downarrow f \\ B & \longrightarrow & B' & \longrightarrow & B \\ & \curvearrowleft & & \curvearrowright & \\ & & \text{id} & & \end{array}$$

Definition. A *model category* is a category \mathcal{C} closed under small³ limits and colimits, together with three subcategories:

- $\operatorname{co}(\mathcal{C})$ (sometimes denoted $\operatorname{cof}(\mathcal{C})$), the subcategory of *cofibrations*,
- $\operatorname{fib}(\mathcal{C})$, the subcategory of *fibrations*, and
- $w(\mathcal{C})$, the subcategory of *weak equivalences*.⁴

We'll also call $w(\mathcal{C}) \cap \operatorname{co}(\mathcal{C})$ the *trivial cofibrations* (as in homotopically trivial), and $w(\mathcal{C}) \cap \operatorname{fib}(\mathcal{C})$ the *trivial cofibrations*. If $f : A \rightarrow B$ is a morphism in $\operatorname{co}(\mathcal{C})$, then it will often be denoted $f : A \hookrightarrow B$, and similarly if $f \in \operatorname{fib}(\mathcal{C})$, then it'll be denoted $f : A \twoheadrightarrow B$. Weak equivalences will be denoted with \simeq .

This data is subject to the following axioms.

- (1) The *two-out-of-three axiom*: if $f : A \rightarrow B$ and $g : B \rightarrow C$ are morphisms in \mathcal{C} such that two of gf , g , and f are in $w(\mathcal{C})$, then the third is.
- (2) Retracts of cofibrations, fibrations, and weak equivalences are cofibrations, fibrations, and retracts, respectively; that is, each of these three categories is closed under retracts.

³We really don't care about set-theoretic issues in this class, as evidenced when the first day didn't begin with "In the beginning, there was a Grothendieck universe." We'll primarily use finite limits and colimits, and requiring pushouts to exist is good enough for us.

⁴Sometimes $w(\mathcal{C})$ is called the "water closet," albeit not in serious literature.

- (3) The *lifting axiom*: let $i : A \rightarrow B$ be a trivial cofibration and $p : X \rightarrow Y$ be a fibration. If the following diagram commutes, then the following dotted arrow exists.

$$\begin{array}{ccc} A & \xrightarrow{\quad} & X \\ \downarrow i & \nearrow \text{dotted} & \downarrow p \\ B & \xrightarrow{\quad} & Y \end{array}$$

Similarly, if i is a cofibration and p is a trivial fibration, then the following dotted arrow exists.

$$\begin{array}{ccc} A & \xrightarrow{\quad} & X \\ \downarrow i & \nearrow \text{dotted} & \downarrow p \\ B & \xrightarrow{\quad} & Y \end{array}$$

- (4) *Functorial factorization*: Every morphism $f : X \rightarrow Y$ can be factored as $X \hookrightarrow Y' \xrightarrow{\sim} Y$, where the first map is in $\text{co}(\mathcal{C})$ and the second in $w(\mathcal{C}) \cap \text{fib}(\mathcal{C})$; it can also be factored as $X \xrightarrow{\sim} X' \rightarrow Y$, where the first map is a trivial fibration and the second is a cofibration. Moreover, these factorizations are functorial in that (for the first factorization; the second is similar) there exists a functor $(F, G) : \text{Mor } \mathcal{C} \rightarrow \text{Mor } \mathcal{C} \times \text{Mor } \mathcal{C}$ such that $F(f)$ is a cofibration, $G(f)$ is a trivial fibration, and $G(f) \circ F(f) = f$.

Remark. \mathcal{C} has an initial and final object, since pushouts exist. These will be denoted 0 and $*$, respectively.

The lifting axiom looks pretty specifically homotopical, but it's still surprisingly minimal for all of this homotopy theory to exist in a category. And functorial factorization is basically cellular approximation.

It should ease your mind that topological spaces form a model category, with fibrations given by fibrations, cofibrations given by cofibrations, and weak equivalences given by weak homotopy equivalences. This is nonetheless hard to show; simplicial sets are also a category, but this is even harder to show.

Definition.

- An object B is called *cofibrant* if $0 \rightarrow B$ is a cofibration.
- Similarly, Y is called *fibrant* if $Y \rightarrow *$ is a fibration.

These objects tend to have nicer properties. Moreover, every object is weakly equivalent to a cofibrant and a fibrant object.

Definition. Given any object B , $0 \rightarrow B$ factors as $0 \hookrightarrow QB \xrightarrow{\sim} B$, so that $QB \simeq B$ and QB is cofibrant. QB is called the *cofibrant replacement* of B .

Similarly, $Y \rightarrow *$ factors as $Y \xrightarrow{\sim} RY \rightarrow *$, with $RY \simeq Y$ and RY fibrant; this RY is called the *fibrant replacement* of Y .

There are lots of terms here, but we'll need to use all of them to talk about stuff. Topological spaces are already all fibrant, so intuition can be challenging here.

The lifting property means that fibrations and cofibrations mutually define each other, so there's sort of too much information in these problems.

Definition. Let $X \in \mathcal{C}$ be an object. Then, a *cylinder* for X , denoted $\text{Cyl}(X)$, is a factorization of $X \sqcup X \rightarrow X$ as $X \sqcup X \hookrightarrow \text{Cyl}(X) \xrightarrow{\sim} X$.

Remark. The above factorization need not be the functorial factorization guaranteed by our category. But we can use that, and therefore there's a cylinder functor.

The abstract definition can be realized topologically as something weakly equivalent to X (which is true of $X \times [0, 1]$), but that fills in the area between two copies of X .

We'll see that everything in model categories has a dual. The notion of a path object is dual to the cylinder.

Definition. If $Y \in \mathcal{C}$, then a *path object* for Y , denoted PY , is one that arises from a factorization $Y \xrightarrow{\sim} PY \rightarrow Y \times Y$ of the diagonal map $Y \rightarrow Y \times Y$.

Again, this can be made functorial. This notion, making a space into a path space, also appears in homotopy theory. But we can go the other way, making avatars of our familiar notions in homotopy theory inside a more general model category. For example, we'll define a homotopy!

Note: next lecture is next Tuesday, so don't come to class on Thursday.

Lecture 5.

Homotopy in Model Categories: 9/15/15

We've defined cylinders and path objects in model categories, and today will define homotopies in two flavors, left and right. The next few lectures will be some relatively boring but useful abstract manipulations in model categories; then we'll get to some cool stuff. But you have to take your medicine.

Throughout this lecture, \mathcal{C} will denote a model category.

Definition. Let $f_0, f_1 : X \rightarrow Y$ be maps in \mathcal{C} .

- (1) f_0 is *left homotopic* to f_1 , written $f_0 \simeq_\ell f_1$, if there exists an $H : \text{Cyl}(X) \rightarrow Y$ such that $H \circ i_0 = f_0$ and $H \circ i_1 = f_1$, where $i_0 \sqcup i_1 : X \sqcup X \hookrightarrow \text{Cyl}(X)$.
- (2) f_0 is *right homotopic* to f_1 , written $f_0 \simeq_r f_1$, if there is a map $K : X \rightarrow PY$ such that $p_0 K = f_0$ and $p_1 K = f_1$, where $p_0 \times p_1 : PY \rightarrow Y \times Y$.
- (3) f_0 is *homotopic* to f_1 , written $f_0 \simeq f_1$, if $f_0 \simeq_\ell f_1$ and $f_0 \simeq_r f_1$.
- (4) $f : X \rightarrow Y$ is a *homotopy equivalence* if there exists a $g : Y \rightarrow X$ such that $f \circ g \simeq \text{id}_Y$ and $g \circ f \simeq \text{id}_X$.

This is at once very abstract and somewhat familiar from topology; we'll show that in nice case left and right homotopy are identical, which is why we don't usually talk about them as distinct in topology. The usual topological notion is usually defined as left homotopy, but homotopies of chain complexes are examples of right homotopies.

Again, this isn't the most earthshattering definition, but it's a useful model to hold in your head. Nonetheless, the following results aren't super exciting to prove.

Lemma 5.1.

- (1) Let $f, g : B \rightarrow X$ be maps in \mathcal{C} . If $f \simeq_\ell g$ and $h : X \rightarrow Y$ is any other map, then $h \circ f \simeq_\ell h \circ g$.
- (2) If $e : A \rightarrow B$ is a map and $f \simeq_r g$, then $f \circ e \simeq_r g \circ e$.

Proof. Since $f \simeq_\ell g$, then there's an $H : \text{Cyl}(B) \rightarrow X$ with $H i_0 = f$ and $H i_1 = g$. Then, $h \circ H : \text{Cyl}(B) \rightarrow X \rightarrow Y$ is a left homotopy between $h \circ f$ and $h \circ g$. (2) is similar. \square

A lot of the work we'll be doing will be to get rid of these "handedness issues."

Proposition 5.2. Suppose $f, g : B \rightarrow X$ are two maps in \mathcal{C} .

- (1) If X is fibrant, $f \simeq_\ell g$, and $h : A \rightarrow B$ is any map, then $f \circ h \simeq_\ell g \circ h$.
- (2) If B is cofibrant, $f \simeq_r g$, and $e : X \rightarrow Y$ is any map, then $e \circ f \simeq_r e \circ g$.

Proof. We'll prove (1); then, (2) is similar.

We have a left homotopy $H : \text{Cyl}(B) \rightarrow X$ where $B \sqcup B \hookrightarrow \text{Cyl}(B) \xrightarrow{\sim} B$. We'd prefer a map $\text{Cyl}(B) \xrightarrow{\twoheadrightarrow} B$ (i.e. a trivial fibration).

Factor $\text{Cyl}(B) \xrightarrow{\sim} B$ as $\text{Cyl}(B) \hookrightarrow \text{Cyl}'(B) \xrightarrow{\sim} B$; by the two-out-of-three rule, $\text{Cyl}(B) \hookrightarrow \text{Cyl}'(B)$ is a weak equivalence, and so we have maps

$$B \sqcup B \hookrightarrow \text{Cyl}(B) \hookrightarrow \text{Cyl}'(B) \xrightarrow{\sim} B,$$

which presents $\text{Cyl}'(B)$ as a cylinder object.

The homotopy H can be considered to be a map $H' : \text{Cyl}'(B) \rightarrow X$: by two-out-of-three $\text{Cyl}(B) \xrightarrow{\sim} \text{Cyl}'(B)$ is a trivial cofibration. Since X is fibrant, we can draw the diagram

$$\begin{array}{ccc} \text{Cyl}(B) & \xrightarrow{\quad} & X \\ \downarrow & \nearrow H' & \downarrow \\ \text{Cyl}'(B) & \xrightarrow{\quad} & *, \end{array}$$

so by the path lifting property we get the desired H' . All of this was just to move the homotopy to come out of a different object!

Now, we can form the diagram

$$\begin{array}{ccccccc}
 A \sqcup A & \xrightarrow{h \sqcup h} & B \sqcup B & \longrightarrow & \text{Cyl}'(B) & \longrightarrow & X \\
 \downarrow & \nearrow \text{dashed} & & & \downarrow \cong & & \\
 \text{Cyl}(A) & \longrightarrow & A & \xrightarrow{h} & B & &
 \end{array}$$

This completes the proof, because it gives us a map $\text{Cyl}(A) \rightarrow X$ through which h factors. \square

These proofs look very similar: we don't have very many rules. We can factor and compose, and not very much more.

As an aside, we'll prove Ken Brown's lemma.⁵ It's invoked a lot, often even without a name. Note that no structure on F is assumed below.

Lemma 5.3 (Ken Brown). *Let \mathcal{C} and \mathcal{D} be model categories and $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Suppose F takes trivial cofibrations between cofibrant objects to weak equivalences; then, F takes weak equivalences between cofibrant objects to weak equivalences.*

This seems small, but is extremely useful: if we know what F does to trivial cofibrations on cofibrant objects, we recover it on all weak equivalences of cofibrant objects. The proof is a fun game of model-categorical axioms.

Proof. Suppose $f : A \xrightarrow{\sim} B$ is a weak equivalence of cofibrant objects. Consider $A \sqcup B \rightarrow B$ given by $f \sqcup \text{id}$, and factor it as

$$A \sqcup B \xrightarrow{p} C \xrightarrow{q} B.$$

Consequently, we have

$$0 \hookrightarrow A \hookrightarrow A \sqcup B \xrightarrow{p} C \xrightarrow{q} B \quad (5.1)$$

by the 2/3 rule, $A \hookrightarrow C$ is a trivial cofibration, since C is a cofibrant object.

By assumption, $F(A \hookrightarrow C)$ is a weak equivalence, but $F(5.1) = F(\text{id})$. By 2/3, $B \hookrightarrow C$ is a weak equivalence, and therefore it's a trivial cofibration between cofibrant objects. Then, since $F(B \hookrightarrow C) \cdot F(C \xrightarrow{\sim} B) = F(\text{id})$, then $F(C \xrightarrow{\sim} B)$ is a weak equivalence. Thus, $F(A \hookrightarrow A \sqcup B \hookrightarrow C \rightarrow B) = F(A \xrightarrow{\sim} B)$, so it factors as $F(A \hookrightarrow C) \circ F(C \rightarrow B)$; both of these are weak equivalences, and therefore the composite is to. \square

Again, we'll get back to rational homotopy theory soon. But that lemma will be useful.

Lemma 5.4. *Let B be cofibrant in \mathcal{C} . Then, left homotopy is an equivalence relation in $\text{Map}_{\mathcal{C}}(B, X)$*

We're working towards a goal: we've defined homotopy equivalence and weak equivalence in unrelated ways, and we want to prove that they're related; in the category of topological spaces, this result is just as consciously proven from differing definitions, and is known as Whitehead's theorem.

Proof. We'll leave reflexivity and symmetry to the reader, and they're not hard.

Suppose $f \simeq_{\ell} g$ and $g \simeq_{\ell} h$, where $f, g, h : B \rightarrow X$ via homotopies $H_{f,g} : B' \rightarrow X$ and $H_{g,h} : B'' \rightarrow X$, where B' and B'' are cylinder objects.

We want to "glue" these two cylinders together, and do so in topology. But in category theory we use pushouts. So let's consider the following diagram.

$$\begin{array}{ccccc}
 B & \xrightarrow{i''_0} & B' & \xleftarrow{i'_1} & B \\
 \downarrow i'_1 & & \downarrow & & \\
 B' & \longrightarrow & L & & \\
 \uparrow i'_0 & & & & \\
 B & & & &
 \end{array}$$

⁵People older than Ken Brown call it Kenny Brown's lemma. Never Kenny's lemma.

By the universal property of pushouts, we get a map $\tilde{H} : L \rightarrow X$. However, since L might not be a cylinder object, then \tilde{H} might not be a homotopy. We solve this the only way we know how: by factoring stuff.

We do know that $B \sqcup B \rightarrow K \xrightarrow{\sim} B$, and factor $B \sqcup B \rightarrow K$ as $B \sqcup B \hookrightarrow L' \tilde{B}$, so L' is a cylinder object and therefore gives us a homotopy $\tilde{H}' : L' \rightarrow X$. \square

Lemma 5.5. *Dually, if X is fibrant, then right homotopy is an equivalence relation on $\text{Map}_{\mathcal{C}}(B, X)$.*

We're in the middle of proving that weak equivalences and homotopy equivalences are reasonable things, and behave the way we want them to.

Lemma 5.6. *Let B be cofibrant and $h : X \rightarrow Y$ be a trivial fibration or a weak equivalence between fibrant objects. Then, $\text{Map}_{\mathcal{C}}(B, X) / \simeq_{\ell} \xrightarrow{\cong} \text{Map}(B, Y) / \simeq_{\ell}$.*

Proof. The equivalence of the two hypotheses on h follows from Ken Brown's lemma, so we'll suppose $h : X \xrightarrow{\sim} Y$ is a trivial fibration.

For surjectivity, suppose $g : B \rightarrow Y$ is given. Then, we can lift it in the following diagram.

$$\begin{array}{ccc} 0 & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow \sim \\ B & \longrightarrow & Y \end{array}$$

Injectivity isn't much harder. Suppose $hf \simeq_{\ell} hg$ for two $f, g : B \rightarrow X$, and this left homotopy is via $H : \text{Cyl}(B) \rightarrow Y$. Then, we can lift in the following diagram, which provides a homotopy between f and g .

$$\begin{array}{ccc} B \sqcup B & \xrightarrow{f \sqcup g} & X \\ \downarrow & \nearrow & \downarrow \sim \\ \text{Cyl}(B) & \xrightarrow{H} & Y \end{array}$$

\square

There's a dual statement, given by switching fibrant and cofibrant, and precomposition and postcomposition. It turns out that model categories are sort of self-dual. The proof is the same, with path objects instead of cylinder objects. This means that a lot of books on model categories could often be half as long! Hirshhorn is particularly representative of this.

The next lectures will return to using this in rational homotopy theory, including showing that chain complexes are a model category.

Lemma 5.7. *Let $f, g : B \rightarrow X$. Then, if B is cofibrant, then $f \simeq_{\ell} g$ implies $f \simeq_r g$, and dually, if X is fibrant, then $f \simeq_r g$ implies $f \simeq_{\ell} g$.*

In topological spaces and chain complexes, everything is fibrant, so we're not used to having two kinds of homotopy, and in simplicial sets, everything is cofibrant. So secretly there's two kinds of homotopy, but we haven't needed to use them.

Another way to think about Lemma 5.7 is as a generalization of the adjunction between cylinders and path spaces on the category of topological spaces.

Definition. A *fibrant-cofibrant* object of \mathcal{C} is, unsurprisingly, an object of \mathcal{C} that's both fibrant and cofibrant. The subcategory of fibrant-cofibrant objects and their maps is denoted \mathcal{C}_{cf} .

These are the “nice” objects, where homotopy equivalence is at its nicest, and all the lemmas we've given so far apply.

The following theorem has a somewhat boring proof, which can be found in the lecture notes, but wasn't given in class.

Theorem 5.8. *If \mathcal{C} is a model category, then a map in \mathcal{C}_{cf} is a weak equivalence iff it is a homotopy equivalence.*

This is really important, and here are some consequences.

- We can take the category $\mathcal{C}_{\text{cf}} / \sim$, modding out by weak equivalence. There are weird set-theoretic issues in general unless we take an equivalence relation (which, in this case, weak equivalence is).

- The homotopy category $\mathrm{Ho}(\mathcal{C})$, defined by formally inverting weak equivalences, is equivalent as categories to $\mathcal{C}_{\mathrm{cf}}/\sim$. This is why model categories are nice: homotopy theory is in some sense about notions of equivalence, and this equates two of them.

Exercise 2. Read about ordinals and transfinite composition, and be very uncomfortable with the foundations of mathematics, as you should be.

This will be used next time for the small object argument, but it's not fun to talk about in lecture. We'll be able to actually make a model category, though! And you'll realize that attaching cells in infinite constructions in algebraic topology requires ordinals too.

Lecture 6.

Today is the Cofibrantly Generated Model Categories Day: 9/17/15

"Now we get to do the small object argument. It's fun! I can see you share my enthusiasm, but I'm enjoying myself, and that's all that matters here."

Cofibrantly generated model categories are nice; they can be built and realized relatively nicely; by showing that the category of differential graded \mathbb{Q} -algebras is a cofibrantly generated model category, we'll be able to do some stuff faster.

We'll start off with some fun stuff about the sizes of... things. We'll consider ordinals as categories by analogy, with $a \rightarrow b$ if $a \leq b$, so it makes sense to talk about functors from ordinals to categories.

Definition. Let λ be an ordinal. Then, a λ -sequence in a category \mathcal{C} is a colimit-preserving functor $X : \lambda \rightarrow \mathcal{C}$. Then, the map $X_0 \rightarrow \varinjlim_{\beta < \lambda} X_\beta$ is called a *transfinite composition*.

That is, λ has a diagram associated to it, and X preserves this diagram.

It may be boring to play around with set theory like this, but we make infinite constructions like this (e.g. infinite cell complexes) in topology all the time, and we want to know that we're not breaking set theory.

Definition. Let κ be a cardinal. Then, an ordinal λ is κ -filtered if

- (1) λ is a limit ordinal, and
- (2) for any $S \leq \lambda$ such that $|S| < \kappa$, $\sup(S) < \lambda$.

This is a weird definition, but if κ is finite, being, for example, 10-filtered means being an infinite limit ordinal (a supremum of 10 things is finite, but could be arbitrarily large).⁶

Definition. Let \mathcal{C} be a category with small colimits (i.e. colimits indexed by sets) and $M \subseteq \mathrm{Mor}(\mathcal{C})$ be a collection of morphisms. We then say that $A \in \mathcal{C}$ is κ -small with respect to M if for any κ -filtered ordinal λ and $X : \lambda \rightarrow \mathcal{C}$ with $X_\beta \rightarrow X_{\beta+1}$ in M , we have that as sets,

$$\mathrm{colim}_{\beta < \lambda} \mathrm{Map}_{\mathcal{C}}(A, X_\beta) \xrightarrow{\cong} \mathrm{Map}_{\mathcal{C}}(A, \mathrm{colim}_{\beta < \lambda} X_\beta).$$

This is really saying that a map $A \rightarrow \mathrm{colim} X_\beta$ factors through some X_β .

Exercise 3. Think about why this is equivalent to the definition.

Remark. If we don't specify κ , then saying that A is *small* with respect to M implies that there exists some κ making it true, and if $M = \mathrm{Mor}(\mathcal{C})$, we simply say that A is *small* (sometimes *compact*).

This will allow us to get control over colimits, which is important.

Definition. Let I be a class of maps in \mathcal{C} . We say that $f \in \mathrm{Mor}(\mathcal{C})$ is *I-injective* if f has the right lifting property with respect to I , i.e. $I \circ^R f$.⁷ Similarly, we say that f is an *I-cofibration* if $f \circ^L A$ whenever A is *I-injective*.

That is, $I \circ^R I\text{-inj.}$, and $I\text{-cof} \circ^L I\text{-inj.}$

The next definition looks reasonably topological, but is abstracted enough to be useful for us.

⁶Recall that we built ordinals by von Neumann's model: $1 = \{\bullet\}$, $2 = \{\bullet, \{\bullet\}\}$, and in general $\mathrm{succ}(S) = S \cup \{S\}$; then, the first infinite ordinal is the union of all of the finite ones, and therefore is called a *limit ordinal*, i.e. it's not the successor of some ordinal.

⁷This should really be a box with a slash, but I get package conflicts when I try to make that work.

Definition. If $I \subseteq \text{Mor}(\mathcal{C})$ has all small colimits, then a *relative I -cell complex* is a transfinite composition of pushouts along I of the form

$$\begin{array}{ccc} A & \longrightarrow & X \\ \downarrow \varphi & & \downarrow \\ B & \longrightarrow & X \sqcup_A B, \end{array}$$

where $\varphi \in I$.

Exercise 4. Show that I -cells are I -cofibrant.

The next argument, though still formal, is at least a little closer to topology or model categories.

Theorem 6.1. Let \mathcal{C} be a category with all small colimits and $I \subseteq \text{Mor}(\mathcal{C})$, and suppose the domains of I are all small relative I -cells. Then, there is a functorial factorization

$$(F, G) : \text{Mor}(\mathcal{C}) \longrightarrow \text{Mor}(\mathcal{C}) \times \text{Mor}(\mathcal{C}),$$

such that for all f , $F(f)$ is a relative I -cell and $G(f)$ is I -injective.

That is, every map $A \rightarrow B$ factors as $A \rightarrow C \rightarrow B$, where the first is an I -cell and the second is I -injective.

Proof. This is the celebrated small object argument.

Assume there exists some κ such that all domains are κ -small, and let λ be a κ -filtered ordinal. Our goal will be to define a λ -sequence $Z^\lambda : \lambda \rightarrow \mathcal{C}$ such that $X \rightarrow \text{colim}_{\beta < \lambda} Z^\beta \rightarrow Y$.

Start with $Z_0 = Y$, and if we've defined Z_α for all $\alpha < \beta$, where β is a limit ordinal, then define $Z_\beta = \text{colim}_{\alpha < \beta} Z_\alpha$. If β is a successor ordinal, let S be the set of all diagrams of the form

$$\begin{array}{ccc} A & \longrightarrow & Z_\beta \\ \downarrow \varphi & & \downarrow \\ B & \longrightarrow & Y, \end{array}$$

with $\varphi \in I$, and define $Z_{\beta+1}$ to be the pushout

$$\begin{array}{ccc} \coprod_{s \in S} A_s & \longrightarrow & Z_\beta \\ \downarrow & & \downarrow \\ \coprod_{s \in S} B_s & \longrightarrow & Z_{\beta+1}. \end{array}$$

Then, let Z be the colimit of this sequence. There's obviously an induced $Z \rightarrow Y$, so we want to provide a lift in the following diagram.

$$\begin{array}{ccc} A & \longrightarrow & Z \\ \downarrow \in I & & \downarrow \\ B & \longrightarrow & Y \end{array}$$

but our map $A \rightarrow Z$ must factor through some Z_β . Then, smallness gives us a map $B \rightarrow Z_{\beta+1}$, and therefore into the colimit, which is our desired lift. \square

This seems pedantic, but if you want to make infinite constructions of cells, which you do all the time in homotopy theory, you need to make some kind of smallness. This is the category-theoretic statement that if you map a compact space into some cell complex, the image is compact.

The phrase “ I permits the small object argument” means that the domains of I are all small relative I -cells (which is what made this argument work).

Corollary 6.2. If I permits the small object argument and $f : A \rightarrow B$ is an I -cofibration, then there is an I -cell $g : A \rightarrow C$ such that f is a retract of

$$\begin{array}{ccccc} A & \xlongequal{\quad} & A & \xlongequal{\quad} & A \\ g \downarrow & & f \downarrow & & g \downarrow \\ C & \longrightarrow & B & \longrightarrow & C \end{array}$$

That is, cofibrations are I -cells or retracts thereof. This will be important for characterizing cofibrations later, and the proof is example of an important kind of argument.

Proof. Factor $A \rightarrow B$ as $A \rightarrow C \rightarrow B$, where the intermediate maps are, respectively, an I -cell and I -injective. Then, we get the following diagram.

$$\begin{array}{ccc} A & \xrightarrow{g} & C \\ f \downarrow & \nearrow & \downarrow I\text{-inj.} \\ B & \xlongequal{\quad} & B \end{array}$$

⊠

Remark. This is an example of the “retract argument.”

Now, we’ll give a whole bunch more definitions, avoiding some extremely dry, formal proofs. The upshot will be to show that \mathbb{Q} -CDGAs are a cofibrantly generated model category; then, over the next few lectures, we’ll begin to build their homotopy theory, which will feel less formal and more topological.

Definition. \mathcal{M} is a *cofibrantly generated model category* (CGMC) if there exist $I, J \subseteq \text{Mor}(\mathcal{M})$ such that

- the domains of I and J are small with respect to the sets of I -cells and J -cells, respectively
- $\text{fib}(\mathcal{M}) = J\text{-inj.}$
- $w(\mathcal{M}) \cap \text{co}(\mathcal{M}) = I\text{-inj.}$

Such a structure will be denoted $(\mathcal{M}, w(\mathcal{M}), I, J)$. I is called the *generating cofibrations*, and J is called the *generating trivial cofibrations*.

The first point means that we can (and do!) use the small object argument on I and J .

Proposition 6.3. *Let $(\mathcal{M}, w(\mathcal{M}), I, J)$ be a CGMC. Then,*

- (1) $\text{co}(\mathcal{M}) = I\text{-cof.}$
- (2) *every cofibration is a retract of an I -cell, and*
- (3) *the domains of I are small relative to cofibrations.*

Moreover, all three statements hold with “trivial fibrations” in place of “cofibrations,” and J replacing I .

Theorem 6.4. *Let \mathcal{C} be a category with subcategories $w(\mathcal{C})$, I , and J . Then, $(\mathcal{C}, w(\mathcal{C}), I, J)$ is a CFMC if the following are true.*

- (1) $w(\mathcal{C})$ has the 2/3 property.
- (2) I and J permit the small object argument.
- (3) $J\text{-cell} \subseteq w(\mathcal{C}) \cap I\text{-cof.}$
- (4) $I\text{-inj} \subseteq w(\mathcal{C}) \cap J\text{-inj.}$
- (5) *Either $w(\mathcal{C}) \cap I\text{-cof} \subseteq J\text{-cof}$ or $w(\mathcal{C}) \cap I\text{-inj} \subseteq J\text{-inj.}$*

You should look up a proof, though we won’t prove it ourselves. The point is, if you prove a bunch of lifting properties, you get a CGMC more or less for free.

Let $\text{Ch}_{\mathbb{Q}}$ denote the category of chain complexes over \mathbb{Q} . Let S^{n-1} denote a chain complex with \mathbb{Q} in degree n and nothing else, and D^n denote one with degrees $n+1$ and n with $d: \mathbb{Q} \rightarrow \mathbb{Q}$ given by isomorphism. Then, let $I = \{S^{n-1} \rightarrow D^n \mid n \geq 0\}$ and $J = \{0 \rightarrow D^n \mid n \geq 0\}$.

Theorem 6.5. $(\text{Ch}_{\mathbb{Q}}, w(\text{Ch}_{\mathbb{Q}}), I, J)$ has a cofibrantly generated model category structure, where the weak equivalences are given by quasi-isomorphisms.

Corollary 6.6. *The fibrations in $\text{Ch}_{\mathbb{Q}}$ are surjections; the cofibrations are retracts of relative I -cells. In particular, the cofibrant objects are I -cellular objects, and this is illuminating in general: cofibrant objects are built out of small numbers of cells.*

Topological spaces, Top , form another important example. Let $I = \{S^{n-1} \rightarrow D^n \mid n \geq 1\}$ and $J = \{D^n \rightarrow D^n \times I \mid n \geq 0\}$, and let $w(\text{Top})$ denote the weak homotopy equivalence (i.e. $X \xrightarrow{\sim} Y$ if $\pi_i(X) \cong \pi_i(Y)$ for all i).

Theorem 6.7. $(\text{Top}, w(\text{Top}), I, J)$ is also a cofibrantly generated model category.

A different category of topological spaces appears for the Quillen adjunction between topological spaces and simplicial sets.

You really should read the proofs, which are boring but instructive.

One cool thing is that, not only can we use adjunctions to induce model category structures, but in many cases it's really easy.

Theorem 6.8. *Let $F : \mathcal{C} \rightleftarrows \mathcal{D} : U$ be an adjunction between a CDMC \mathcal{C} and \mathcal{D} satisfying some closure axioms (e.g. \mathcal{D} has all small colimits). If FI and FJ permit the small object argument and U takes relative FJ -cell complexes to weak equivalences, then \mathcal{D} is a CGMC with (FI, FJ) as generating (trivial) cofibrations, and weak equivalences those maps $f \in \mathcal{D}$ such that $U(f) \in w(\mathcal{C})$.*

In practice, these conditions are quite easy to check, so we get a model category structure for free.

There is a free forgetful adjunction $\Lambda : \text{Ch}_{\mathbb{Q}} \rightleftarrows \text{CDGA}_{\mathbb{Q}} : U$ given by $\Lambda(C) = \Lambda_{\mathbb{Q}}[C^{\text{odd}}] \otimes \mathbb{Q}[C^{\text{even}}]$, with differential uniquely determined.

Theorem 6.9. *There is a model category structure on $\text{CDGA}_{\mathbb{Q}}$ induced from $\text{Ch}_{\mathbb{Q}}$ such that $I = \{\Lambda S^{n-1} \rightarrow \Lambda D^n \mid n \geq 1\}$, $J = \{0 \rightarrow \Lambda D^n \mid n \geq 0\}$, and the weak equivalences are quasi-isomorphisms. Moreover, cofibrations are retracts of relative I -cells and fibrations are degree-wise surjections.*

This tells us a lot about homotopies of CDGAs, and in particular the things that are well-behaved. Sometimes, people will avoid this high-level language, and “Sullivan algebra” is a word for a cofibrant object in some model category. Then, a lot of their manipulations to get things to factor come from stuff we’ve already done in model categories.