

MSRI: QUANTUM SYMMETRIES INTRODUCTORY WORKSHOP

ARUN DEBRAY
JANUARY 27–31, 2020

These notes were taken at MSRI’s [introductory workshop on quantum symmetries](#) in Spring 2020. I live-T_EXed them using `vim`, and as such there may be typos; please send questions, comments, complaints, and corrections to a.debray@math.utexas.edu.

CONTENTS

Part 1. Monday, January 27	1
1. Sarah Witherspoon: Hopf algebras, I	1
2. Victor Ostrik, Introduction to fusion categories, I	4
3. Eric Rowell, An introduction to modular tensor categories I	6
4. Emily Peters, Subfactors and planar algebras I	8
Part 2. Tuesday, January 28	10
5. Victor Ostrik, Introduction to fusion categories, II	10
References	12

Part 1. Monday, January 27

1. SARAH WITHERSPOON: HOPF ALGEBRAS, I

Our perspective on Hopf algebras, their actions on rings and modules, and the structures on their categories of rings and modules, will be to think of them as generalizations of group actions and representations; groups actions are symmetries in the usual sense, and Hopf algebra actions are often related to “quantum symmetries.”

We’re not going to give the full definition of a Hopf algebra, because it would require drawing a lot of commutative diagrams, but we’ll say enough to give the picture.

Throughout this talk we work over a field k ; all tensor products are of k -vector spaces.

Definition 1.1. A *Hopf algebra* is an algebra A together with k -linear maps $\Delta: A \rightarrow A \otimes A$, called *comultiplication*; $\varepsilon: A \rightarrow k$, called the *counit*; and $S: A \rightarrow A$, called the *coinverse*. These maps must satisfy some properties, including that ε is an algebra homomorphism and that S is an *anti-automorphism*, i.e. that $S(xy) = S(y)S(x)$.

The definition is best understood through examples.

Example 1.2.

- (1) Let G be a group. Then the group algebra $k[G]$ is a Hopf algebra, where for all $g \in G$, $\Delta(g) := g \otimes g$, $\varepsilon(g) := 1$, and $S(g) := g^{-1}$. This is a key example that allows us to generalize ideas from group actions to Hopf algebra actions: whenever we define a notion for Hopf algebras, when we implement it for $k[G]$ it should recover that notion for groups.
- (2) Let \mathfrak{g} be a Lie algebra over k . Then its universal enveloping algebra $\mathcal{U}(\mathfrak{g})$ is a Hopf algebra, where for all $x \in \mathfrak{g}$, $\Delta(x) := x \otimes 1 + 1 \otimes x$, $\varepsilon(x) := 0$, and $S(x) := -x$. Since ε is an algebra homomorphism, $\varepsilon(1_{\mathcal{U}(\mathfrak{g})}) = 1$.

For example,

$$(1.3) \quad \mathcal{U}(\mathfrak{sl}_2) = k\langle e, f, h \mid ef - fe = h, he - eh = 2e, hf - fh = -2f \rangle,$$

given explicitly by the basis of \mathfrak{sl}_2

$$(1.4) \quad e := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad f := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad h := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad \blacktriangleleft$$

Both of these examples are classical, in that they've been known for a long time. But more recently, in the 1980s, people discovered new examples, coming from quantum groups.

Example 1.5 (Quantum \mathfrak{sl}_2). Let $q \in k^\times \setminus \{\pm 1\}$. Then, given a simple Lie algebra \mathfrak{g} , we can define a “quantum group,” $\mathcal{U}_q(\mathfrak{g})$, which is a Hopf algebra. For example, for \mathfrak{sl}_2 ,

$$(1.6) \quad \mathcal{U}_q(\mathfrak{sl}_2) = k \left\langle E, F, K^{\pm 1} \mid EF - FE = \frac{K - K^{-1}}{q - q^{-1}}, KE = q^2 EK, KF = q^{-2} EK \right\rangle,$$

with comultiplication

$$(1.7a) \quad \Delta(E) := E \otimes 1 + K \otimes E$$

$$(1.7b) \quad \Delta(F) := F \otimes K^{-1} + 1 \otimes F$$

$$(1.7c) \quad \Delta(K^{\pm 1}) := K^{\pm 1} \otimes K^{\pm 1}$$

and counit $\varepsilon(E) = \varepsilon(F) = 0$ and $\varepsilon(K) = 1$. This generalizes to other simple \mathfrak{g} , albeit with more elaborate data. \blacktriangleleft

Example 1.8 (Small quantum \mathfrak{sl}_2). Let q be an n^{th} root of unity. Then, as before, given a simple Lie algebra \mathfrak{g} , we can define a Hopf algebra $u_q(\mathfrak{g})$, called the *small quantum group* for \mathfrak{g} and q , which is a finite-dimensional vector space over k ; for \mathfrak{sl}_2 , this is

$$(1.9) \quad u_q(\mathfrak{sl}_2) = \mathcal{U}_q(\mathfrak{sl}_2) / (E^n, F^n, K^n - 1). \quad \blacktriangleleft$$

Before we continue, we need some useful notation for comultiplication, called *Sweedler notation*. Let A be a Hopf algebra and $a \in A$; then we can symbolically write

$$(1.10) \quad \Delta(a) = \sum_{(a)} a_1 \otimes a_2.$$

Comultiplication in a Hopf algebra is *coassociative*, in that as maps $A \rightarrow A \otimes A \otimes A$,

$$(1.11) \quad (\Delta \otimes \text{id}) \circ \Delta = (\text{id} \otimes \Delta) \circ \Delta.$$

Therefore when we iterate comultiplication, we can symbolically write

$$(1.12) \quad (\text{id} \otimes \Delta) \circ \Delta(a) = \sum_{(a)} a_1 \otimes a_2 \otimes a_3$$

without worrying about parentheses.

Actions on rings. Hopf algebra actions on rings generalize group actions on rings by automorphisms and actions of Lie algebras on rings by derivations. If a group G acts on a ring R , then for all $g \in G$ and $r, r' \in R$,

$$(1.13a) \quad g(rr') = (gr)(gr')$$

$$(1.13b) \quad g(1_R) = 1_R.$$

In $k[G]$, our Hopf algebra avatar of G , $\Delta(g) = g \otimes g$, and $\varepsilon(g) = 1$.

If a Lie algebra \mathfrak{g} acts on a ring R by derivations, then for all $x \in \mathfrak{g}$ and $r, r' \in R$,

$$(1.14a) \quad x \cdot (rr') = (x \cdot r)r' + r(x \cdot r')$$

$$(1.14b) \quad x \cdot (1_R) = 0.$$

In $\mathcal{U}(\mathfrak{g})$, our Hopf algebra avatar of \mathfrak{g} , $\Delta(x) = x \otimes 1 + 1 \otimes x$, and $\varepsilon(x) = 0$. These two examples suggest how we should implement a general Hopf algebra action on a ring: comultiplication tells us how to act on the product of two elements, and the counit tells us how to act on 1.

Definition 1.15. Let A be a Hopf algebra and R be a k -algebra. An A -module algebra structure on R is data of an A -module structure on R such that for all $a \in A$ and $r, r' \in R$,

$$(1.16a) \quad a \cdot (rr') = \sum_{(a)} (a_1 \cdot r)(a_2 \cdot r')$$

$$(1.16b) \quad a \cdot (1_R) = \varepsilon(a)1_R.$$

Thus a group action as in (1.13) defines an action of the Hopf algebra $k[G]$, and a Lie algebra action as in (1.14) defines an action of the Hopf algebra $\mathcal{U}(\mathfrak{g})$.

Example 1.17. The quantum analogue of the \mathfrak{sl}_2 -action on $k[x, y]$, thought of as (functions on the) plane, there is an action of $\mathcal{U}_q(\mathfrak{sl}_2)$ on the *quantum plane*

$$(1.18) \quad R := k\langle x, y \mid xy = qyx \rangle.$$

This is a deformation of $k[x, y]$, which is the case $q = 1$. The explicit data of the action is

$$(1.19) \quad E \cdot x = 0 \quad F \cdot x = y \quad K^{\pm 1} \cdot x = q^{\pm 1}x$$

$$(1.20) \quad E \cdot y = x \quad F \cdot y = 0 \quad K^{\pm 1} \cdot y = q^{\mp 1}y.$$

One has to check that this extends to an action satisfying Definition 1.15, but it does, and R is an A -module algebra. Here E and F act as *skew-derivations*, e.g.

$$(1.21) \quad E \cdot (rr') = (E \cdot r)r' + (K \cdot r)(E \cdot r')$$

for all $r, r' \in R$. ◀

Given a Hopf algebra action of A on R in this sense, we can construct two useful rings: the *invariant subring*

$$(1.22) \quad R^A := \{r \in R \mid a \cdot r = \varepsilon(a) \cdot r \text{ for all } a \in A\},$$

and the *smash product ring* $R \# A$, which as a vector space is $R \otimes A$, with multiplication given by

$$(1.23) \quad (r \otimes a)(r' \otimes a') := \sum_{(a)} r(a_1 \cdot r') \otimes a_2 a'.$$

The smash product ring knows the A -module algebra structure on R . Often, rings we're interested in for other reasons are smash product rings of interesting Hopf algebra actions, and identifying this structure is useful.

Example 1.24. The *Borel subalgebra* of $\mathcal{U}_q(\mathfrak{sl}_2)$ is $k\langle E, K^{\pm 1} \mid KE = q^{-2}K \rangle$. This is isomorphic to the smash product $k[E] \# k\langle K \rangle$, where $k\langle K \rangle$ is the group algebra of the free group on the single generator K .

In fact, there's a sense in which $\mathcal{U}_q(\mathfrak{sl}_2)$ is a deformation of $k[E, F] \# k\langle K \rangle$: in this smash product ring, E and F commute, and we deform this to $\mathcal{U}_q(\mathfrak{sl}_2)$, in which they don't commute. ◀

Modules. Given a Hopf algebra A , what is the structure of its category of modules? The first thing we can do is take the tensor product of A -modules U and V using comultiplication: for $a \in A$, $u \in U$, and $v \in V$,

$$(1.25) \quad a \cdot (u \otimes v) = \sum_{(a)} a_1 \cdot u \otimes a_2 \cdot v.$$

Moreover, k has a canonical A -module structure via the counit: $a \cdot x := \varepsilon(a)x$ for $a \in A$ and $x \in k$. Finally, if U is an A -module, its vector space dual $U^* := \text{Hom}_k(U, k)$ has an A -module structure via S : for all $a \in A$, $u \in U$, and $f \in U^*$, $(a \cdot f)(u) := f(S(a)u)$.

The existence of tensor products, duals, and the ground field in the world of Hopf algebra modules is a nice feature: these aren't always present for a general associative algebra. Moreover, these constructions interact well with each other.

- (1) Coassociativity of Δ implies the tensor product is associative: for A -modules U , V , and W , we have a natural isomorphism $U \otimes (V \otimes W) \xrightarrow{\cong} (U \otimes V) \otimes W$.

(2) In any Hopf algebra A , we have the condition

$$(1.26) \quad \sum_{(a)} \varepsilon(a_1) a_2 = \sum_{(a)} a_1 \varepsilon(a_2)$$

for any $a_1, a_2 \in A$. This implies k , as an A -module, is the unit for the tensor product: we have natural isomorphisms $k \otimes U \cong U \cong U \otimes k$ for an A -module U .

(3) Suppose U is an A -module which is a finite-dimensional k -vector space. Then it comes with data of a *coevaluation map* $c: k \rightarrow U \otimes U^*$ sending

$$(1.27) \quad 1 \mapsto \sum_i u_i \otimes u_i^*,$$

where $\{u_i\}$ is a basis for U over k and $\{u_i^*\}$ is its dual basis; this map turns out to be independent of basis. We also have an *evaluation map* $e: U^* \otimes U \rightarrow k$ sending $f \otimes u \mapsto f(u)$. Now, not only are these A -module homomorphisms, but the composition

$$(1.28) \quad U \xrightarrow{c \otimes \text{id}_U} U \otimes U^* \otimes U \xrightarrow{\text{id}_U \otimes e} U$$

is the identity map.

Definition 1.29. A *tensor category*, or *monoidal category* is a category \mathcal{C} together with a functor $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$, an object $\mathbf{1} \in \mathcal{C}$ called the *unit*, and natural isomorphisms $U \otimes (V \otimes W) \cong (U \otimes V) \otimes W$ and $\mathbf{1} \otimes U \cong U \cong U \otimes \mathbf{1}$ for all objects U, V , and W in \mathcal{C} , subject to some coherence conditions.

Our key examples of tensor categories are the category of modules over a Hopf algebra A , as well as the subcategory of finite-dimensional modules.

If the coinverse of A is invertible, which is always the case when A is finite-dimensional over k , then $\mathcal{C} = \text{Mod}_A$ is a *rigid* tensor category, meaning that every object U has a *right dual* ${}^*U := \text{Hom}_k(U, k)$, which means the composition (1.28) is the identity.

Remark 1.30. Notations for left and right duals differ. We're following [EGNO15], but Bakalov-Kirillov [BK01] use a different convention; be careful! \blacktriangleleft

Some Hopf algebras' categories of modules have additional structure or properties: they might be semisimple, or braided, or even symmetric. This amounts to additional information on the Hopf algebra itself.

2. VICTOR OSTRIK, INTRODUCTION TO FUSION CATEGORIES, I

In the world of classical symmetries, i.e. those given by group actions, there is a particularly nice subclass: finite groups. If you know your symmetry group is finite, you can take advantage of many simplifying assumptions. Likewise, in the setting of quantum symmetries, given by, say, \mathbb{C} -linear tensor categories, fusion subcategories form a very nice subclass for which many simplifying assumptions hold. And indeed, if G is a finite group, its category of finite-dimensional representations is a fusion category.

Recall that a *monoidal category* is a category \mathcal{C} together with a functor $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ and a distinguished object $\mathbf{1} \in \mathcal{C}$ called the *unit*, together with natural isomorphisms implementing associativity of \otimes , via $(X \otimes Y) \otimes Z \xrightarrow{\cong} X \otimes (Y \otimes Z)$; and unitality of $\mathbf{1}$, via $\mathbf{1} \otimes X \xrightarrow{\cong} X \xrightarrow{\cong} X \otimes \mathbf{1}$. These must satisfy some axioms which we won't discuss in detail here; the most important one is the *pentagon axiom* on the associator.

Today, we work over an algebraically closed field k , not necessarily closed. Recall that a *k -linear category* \mathcal{C} is one for which for all objects $x, y \in \mathcal{C}$, $\text{Hom}_{\mathcal{C}}(x, y)$ is a k -vector space, such that composition is bilinear. A *k -linear monoidal category* is a monoidal category that is also a k -linear category — and we also impose the consistency condition that the tensor product is a k -linear functor. we will impose a few more niceness conditions before arriving at the definition of a fusion category — in fact, as many as we can such that we still have examples!

In particular, we will only consider k -linear monoidal categories \mathcal{C} such that

- all Hom-spaces are finite-dimensional over k ,
- \mathcal{C} is semisimple,¹
- \mathcal{C} has only finitely many isomorphism classes of simple objects,

¹A k -linear category is *semisimple* if it's equivalent to the category of modules over $k \oplus \cdots \oplus k$, where there is a finite number of summands.

- $\mathbf{1}$ is indecomposable, and
- \mathcal{C} is *rigid*, a condition on duals of objects.

A category satisfying all of these axioms is a *fusion category*. (TODO: double-check)

There are three ways we can come to an understanding of these categories: through the definition, through realizations and examples, and through diagrammatics. We will also heavily use semisimplicity, through the principle that *k-linear functors out of \mathcal{C} are determined by their values on simple objects, and all choices are allowed*.

Example 2.1. Our running example is $\mathcal{V}ec_{\mathbb{Z}/n}^\omega$, where n is a natural number and ω is a degree-3 cocycle for \mathbb{Z}/n , valued in k^\times .

The objects of $\mathcal{V}ec_{\mathbb{Z}/n}^\omega$ are the elements of \mathbb{Z}/n , with the tensor product $i \otimes j := i + j$. If $\omega = 1$, then we use the obvious associator, i.e. the isomorphism

$$(2.2) \quad (i \otimes j) \otimes k \xrightarrow{\cong} i \otimes (j \otimes k)$$

which corresponds to the identity under the identifications with $i + j + k$.² But in general, we can do something different: choose the map (2.2) which is $\omega(i, j, k)$ times the standard one.

A priori you can use any function $\mathbb{Z}/n \times \mathbb{Z}/n \times \mathbb{Z}/n \rightarrow k^\times$, but the pentagon axiom on associativity imposes the condition that ω is a cocycle.

Exercise 2.3. If you have not seen this before, verify that the pentagon axiom forces $\partial\omega = 1$.

The simplest nontrivial example³ is for $n = 2$ and

$$(2.4) \quad \omega(i, j, k) := \begin{cases} 1, & \text{if } i = 0, j = 0, \text{ or } k = 0 \\ -1, & \text{otherwise.} \end{cases} \quad \blacktriangleleft$$

\mathbb{Z}/n was not special here — given any finite group G and a cocycle $\omega \in Z^3(G; k^\times)$, we obtain a fusion category $\mathcal{V}ec_G^\omega$ in the same way.

With ω as in (2.4), $\mathcal{V}ec_{\mathbb{Z}/2}^\omega$ looks like a new example, not equivalent to $\mathcal{V}ec_G^0$ for any G — but in order to understand that precisely, we need to discuss when two tensor categories are equivalent.

Definition 2.5. A *tensor equivalence* of tensor categories \mathcal{C} and \mathcal{D} is a monoidal functor $F: \mathcal{C} \rightarrow \mathcal{D}$, i.e. a functor together with data of natural isomorphisms $F(X \otimes Y) \xrightarrow{\cong} F(X) \otimes F(Y)$ satisfying some axioms.

Choose cocycles ω and ω' for \mathbb{Z}/n , and let's consider tensor functors $F: \mathcal{V}ec_{\mathbb{Z}/n}^\omega \rightarrow \mathcal{V}ec_{\mathbb{Z}/n}^{\omega'}$. Furthermore, let's assume F is the identity on objects, so the data of F is the natural isomorphism $F(X \otimes Y) \cong F(X) \otimes F(Y)$. This is a choice of an element of k^\times for every pair of objects, subject to some additional conditions:

Proposition 2.6. F is a tensor functor iff $\omega = \omega' \cdot \partial\psi$.

Corollary 2.7. $\mathcal{V}ec_{\mathbb{Z}/n}^\omega \simeq \mathcal{V}ec_{\mathbb{Z}/n}^{\omega'}$ if ω and ω' are cohomologous.

Recall that $H^3(\mathbb{Z}/n; k^\times) \cong \mathbb{Z}/n$, so we have n possibilities, some of which might coincide. If F isn't the identity on objects, it's fairly easy to see that as a function on objects, identified with a function $\mathbb{Z}/n \rightarrow \mathbb{Z}/n$, we must get a group homomorphism; if F is to be an equivalence, this homomorphism must be an isomorphism. One can run a similar argument as above and obtain a nice classification result.

Proposition 2.8. The tensor equivalence classes of tensor categories $\mathcal{V}ec_{\mathbb{Z}/n}^\omega$ are in bijection with the orbits $H^3(\mathbb{Z}/n; k^\times) / \text{Aut}(\mathbb{Z}/n)$, via the map sending ω to its class in cohomology.

The action of $\text{Aut}(\mathbb{Z}/n) = (\mathbb{Z}/n)^\times$ on $H^3(\mathbb{Z}/n; k^\times) \cong \mathbb{Z}/n$ is not the first action you might write down! Given $a \in (\mathbb{Z}/n)^\times$ and $s \in H^3(\mathbb{Z}/n; k^\times)$, the action is

$$(2.9) \quad a \cdot s = a^2 s.$$

This is a standard fact from group cohomology.

²These multiplication rules are really special, in that we were able to just write down an associator. This is generally not true; for general multiplication rules you're interested in, you'll have to work a little harder.

³This is nontrivial provided $\text{char}(k) \neq 2$.

Now let's discuss some realizations of fusion categories. If H is a semisimple Hopf algebra, then $\mathcal{C} := \mathcal{R}ep_H^{fd}$ is a fusion category. Let $F: \mathcal{C} \rightarrow \mathcal{V}ec$ denote the forgetful functor to finite-dimensional vector spaces. It turns out that one can reconstruct \mathcal{C} as a fusion category from F , and in fact any fusion category \mathcal{C} with a tensor functor to $\mathcal{V}ec$ is equivalent to $\mathcal{R}ep_H^{fd}$ for some Hopf algebra H . The data of the tensor functor to $\mathcal{V}ec$ is crucial!

Example 2.10. For example, $\mathcal{V}ec_{\mathbb{Z}/n} \simeq \mathcal{R}ep_{\mathbb{Z}/n}^{fd}$; we saw in the previous lecture that representations of \mathbb{Z}/n are equivalent to modules over the Hopf algebra $k[\mathbb{Z}/n] := k[x]/(x^n - 1)$, with comultiplication $\Delta(x) := x \otimes x$.

However, if ω is nontrivial, $\mathcal{V}ec_{\mathbb{Z}/n}^\omega$ admits no tensor functor to $\mathcal{V}ec$, and therefore cannot be seen using Hopf algebras. One can try to generalize the reconstruction program, using quasi-Hopf algebras, weak Hopf algebras, etc. \blacktriangleleft

Bimodules provide another approach to realizations: we look for a ring R and a tensor functor $F: \mathcal{C} \rightarrow \mathcal{B}imod_R$. Applying this to $\mathcal{V}ec_{\mathbb{Z}/n}^\omega$, we get (R, R) -bimodules $F(i)$ for each $i \in \mathbb{Z}/n$ and isomorphisms $F(i) \otimes_R F(j) \xrightarrow{\cong} F(i+j)$. In particular, each $F(i)$ is (tensor-)invertible.

Example 2.11. An *inner automorphism* of a ring R is conjugation by some $r \in R^\times$. Inner automorphisms form a normal subgroup of $\text{Aut}(R)$, and the quotient is called the *outer automorphism group* of R and denoted $\text{Out}(R)$. An *outer action* of a group G on a ring R is a group homomorphism $\varphi: G \rightarrow \text{Out}(R)$.

Given an outer automorphism θ of R , one obtains an (R, R) -bimodule R_θ , whose left action is the R -action on R by left multiplication, and whose right action is $r \cdot x = r\theta(x)$. We need to choose an element in $\text{Aut}(R)$ mapping to θ to make this definition, but different choices lead to isomorphic bimodules.

Anyways, given an outer action of \mathbb{Z}/n on R , we obtain (R, R) -bimodules $R_{\varphi(i)}$ indexed by the objects $i \in \mathcal{V}ec_{\mathbb{Z}/n}$ and isomorphisms between $R_{\varphi(i)} \otimes R_{\varphi(j)} \xrightarrow{\cong} R_{\varphi(i+j)}$. This data stitches together into a tensor functor $\mathcal{V}ec_{\mathbb{Z}/n} \rightarrow \mathcal{B}imod_R$. \blacktriangleleft

Diagrammatics represents the objects of a fusion category \mathcal{C} as points, and morphisms as lines. One can then impose relations on certain morphisms, and therefore diagrammatics provide a generators-and-relations approach to the structure of a given fusion category. Next time, we'll see how to do this for $\mathcal{V}ec_{\mathbb{Z}/n}^\omega$, and see more examples.

3. ERIC ROWELL, AN INTRODUCTION TO MODULAR TENSOR CATEGORIES I

In this lecture, we'll begin with definitions and basic examples of modular tensor categories, and then use them in the next lecture. But first, let's discuss the whys of modular tensor categories.

We're often interested in knot and link invariants which are pictorial in nature, e.g. computed using a diagram. Another seemingly unrelated application is to study statistical-mechanical systems. Witten introduced TQFT into this story, extending the Jones polynomial to 3-manifold invariants using physics. Lately, there are interesting condensed-matter phenomena in topological phases. All of these are governed by modular tensor categories in different ways, and in related ones.

(TODO: list of references, via handout)

Definition 3.1. Let \mathcal{C} be a fusion category. A *braiding* on \mathcal{C} (after which it's called a *braided fusion category*) is data of a natural transformation $c_{X,Y}: X \otimes Y \xrightarrow{\cong} Y \otimes X$ satisfying some relations called the *hexagon identities*.

You can think of $c_{X,Y}$ as taking strands labeled by the objects X and Y , and laying the X strand over the Y strand. The hexagon identities arise by comparing the two strands

$$(3.2) \quad \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \quad \text{and} \quad \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array}.$$

Because the braiding is implemented via a natural transformation, it is functorial: we can braid morphisms as well as objects.

Example 3.3. Given a finite group G , \mathcal{Rep}_G is a braided fusion category. Let V and W be representations; then the braiding $c_{V,W}(v \otimes w) := w \otimes v$. \blacktriangleleft

Definition 3.4. Let \mathcal{C} be a braided fusion category. The *symmetric center* or *Müger center* of \mathcal{C} is the subcategory \mathcal{C}' of $x \in \mathcal{C}$ such that $c_{X,Y}c_{Y,X} = \text{id}_X$ for all $Y \in \mathcal{C}$.

For example, the symmetric center of \mathcal{Rep}_G is once again \mathcal{Rep}_G .

Exercise 3.5. Why is the symmetric center of \mathcal{C} a braided fusion category? In particular, why is it closed under tensor products?

Definition 3.6. If the symmetric center of \mathcal{C} is itself, we call \mathcal{C} *symmetric*.⁴ If the symmetric center of \mathcal{C} is generated by the unit object (equivalently, $\mathcal{C}' \simeq \mathcal{Vect}$), we call \mathcal{C} *nondegenerate*.

Here, “generated by the unit object” means every object is isomorphic to a direct sum of copies of the unit.

Now let’s put some more adjectives in front of these structures. These will make the structure nicer, as usual, but are interesting enough to have examples.

Definition 3.7. Let \mathcal{C} be a braided fusion category. A *twist* on \mathcal{C} is a choice of $\theta \in \text{Aut}(\text{id}_{\mathcal{C}})$.

Diagrammatically, we think of the twist as acting by the diagram in the first Reidemeister move, except we place right over left, not left over right. By looking at a picture of the twist on $X \otimes Y$, and untangling the picture, you can prove the *balancing equation*

$$(3.8) \quad \theta_{X \otimes Y} = c_{X,Y} \circ \theta_X \otimes \theta_Y.$$

Diagrams make it easier to picture these relations, but aren’t strictly necessary. For example, the evaluation map $d_X: X^* \otimes X \rightarrow \mathbf{1}$ is represented by a diagram \frown labeled by X , and coevaluation $b_X: \mathbf{1} \rightarrow X^* \otimes X$ is represented by a diagram \smile labeled by X . Since braided categories aren’t necessarily symmetric, one must be careful with left versus right duals.

Definition 3.9. A *ribbon structure* on a braided fusion category \mathcal{C} is a twist such that $(\theta_X)^* = \theta_{X^*}$.

TODO: picture goes here. Here’s where it’s useful to use ribbon diagrams rather than string diagrams: really we want to keep track of the normal framings of the strings in our diagrams (thought of as embedded in \mathbb{R}^3), and ribbons provide a clean way to understand that.

Let $\text{Irr}(\mathcal{C})$ denote the set of isomorphism classes of irreducible objects in \mathcal{C} . This is always a finite set; the *rank* of \mathcal{C} is $\#\text{Irr}(\mathcal{C})$. Choose representatives x_1, \dots, x_r of the isomorphism classes of simple objects; then, by Schur’s lemma, $\text{Aut}(X_i) \cong \mathbb{C}^\times$. Let $\theta_i \in \mathbb{C}^\times$ denote the twist of X_i .

Now we have all the words we need to define modular tensor categories.

Definition 3.10. A *modular tensor category* is a nondegenerate ribbon fusion category.

There are other, equivalent definitions.

Definition 3.11. A *pivotal structure* on a fusion category \mathcal{C} is a natural isomorphism $j: X \xrightarrow{\cong} X^{**}$.

If a pivotal structure satisfies a certain niceness condition, it’s called *spherical*. Then:

- A braided fusion category with a pivotal structure automatically has a twist.
- If that pivotal structure is spherical, the twist defines a ribbon structure.
- A nondegenerate braided fusion category with a spherical structure is a modular tensor category.

This still hasn’t quite made contact with the usual definition.

If \mathcal{C} is a ribbon fusion category, it has a canonical trace on $\text{End}(X)$, valued in $\text{End}(\mathbf{1}) \cong \mathbb{C}$. The *dimension* of an object $X \in \mathcal{C}$ is $\text{tr}(\text{id}_X)$.

Definition 3.12. The *S-matrix* of a ribbon fusion category is the matrix with entries $S_{ij} := \text{tr}(c_{X_i, X_j} \circ c_{X_j, X_i})$ for $X_i, X_j \in \text{Irr}(\mathcal{C})$.

Theorem 3.13 (Brugières-Müger). *A ribbon tensor category \mathcal{C} is modular if and only if the S-matrix is invertible.*

Now let’s turn to examples.

⁴Notice that being symmetric is a property of braided fusion categories.

Example 3.14. Let G be a finite abelian group and $\mathcal{V}ec_G$ be the category of G -graded vector spaces. These were discussed previously in Example 2.1, albeit in a slightly different way.

Let $c: G \times G \rightarrow \mathbb{C}^\times$ be a *bicharacter* of G , i.e. for all $g, h, k \in G$,

$$(3.15) \quad c(gh, k) = c(g, k)c(h, k).$$

Then we obtain a braiding on $\mathcal{V}ec_G$ by $c: g \otimes h \rightarrow h \otimes g$ by

$$(3.16) \quad \theta_g(v \otimes w) = c(g, h)w \otimes v.$$

For the twist, use $\theta_g := c(g, g)$. This defines a ribbon tensor category, and it is modular iff $\det((c(g, h)c(h, g))_{g, h}) \neq 0$.

Exercise 3.17. In particular, let $G := \mathbb{Z}/3$ and w be a generator. Show that $c(w, w) = \exp(2\pi i/3)$ extends to a bicharacter that defines a modular tensor structure on $\mathcal{C} := \mathcal{V}ec_G$. Show that we cannot obtain a modular structure on $\mathcal{V}ec_{\mathbb{Z}/2}$ in this way, however.

We can produce a modular structure on $\mathcal{V}ec_{\mathbb{Z}/2}$ in a different way: let z be a generator, and define $c(z, z) := i$ and $c(1, z) = c(z, 1) = c(1, 1) = 1$. This defines a modular tensor category structure on $\mathcal{V}ec_{\mathbb{Z}/2}^\omega$ whenever ω is cohomologically nontrivial; this category is of considerable interest in physics, where it's known as the *semion category*. ◀

If you tried to generalize this to G nonabelian, you would not be able to write down a braiding, because $g \otimes h \not\cong h \otimes g$.

If all simple objects in \mathcal{C} are invertible, \mathcal{C} is called a *pointed fusion category*. It turns out these have been classified, and the underlying monoidal tensor category is $\mathcal{V}ec_G^\omega$ for some finite group G and some cocycle ω . If in addition \mathcal{C} is braided, then G is abelian, and we can ask about the converse.

Theorem 3.18. If $|G|$ is odd, $\mathcal{V}ec_G^\omega$ admits a braiding iff ω is cohomologically trivial.

When $|G|$ is even, things are more complicated, as we saw above, but the answers are known. For $\mathbb{Z}/2$, we can get $\mathcal{R}ep_{\mathbb{Z}/2}$, and for $c(z, z) = -1$, we obtain $s\mathcal{V}ec$. Both of these are symmetric. One can generalize: Deligne [Del02] classified symmetric fusion categories, showing they're all equivalent to $\mathcal{R}ep_G$ or $\mathcal{R}ep_G(z)$, where $z \in G$ is central and order 2 (giving a super-vector space structure on G -representations). Symmetric fusion categories equivalent to $\mathcal{R}ep_G$ are called *Tannakian*; those equivalent to $\mathcal{R}ep_G(z)$ are called *super-Tannakian*.

4. EMILY PETERS, SUBFACTORS AND PLANAR ALGEBRAS I

Note: I (Arun) didn't fully understand this talk, and there are a lot of **TODOs**. Hopefully I can fix some of them soon. I'm sorry about that.

In the subject of planar algebra, one can do a lot of math by drawing pictures and reasoning carefully about them. So these talks will have plenty of pictures.

References for today's talk:

- Jones, "Planar algebras I," [Jon99] the original reference.
- The speaker's thesis.
- Heunen and Vicary, "Categories for quantum theory."

Definition 4.1. A *Temperley-Lieb diagram* of size n is an embedding of n disjoint copies of $[0, 1]$ into $[0, n] \times [0, 1]$, such that the boundaries of the embedded intervals lie on integer-valued points.

That is, we take an $n \times 2$ rectangle of points, and draw lines pairing them, such that no two lines cross. We identify two Temperley-Lieb diagrams which are isotopic.

Let \mathcal{TL}_n denote the complex vector space spanned by Temperley-Lieb diagrams of size n . Addition is formal. \mathcal{TL}_n acquires an algebra structure by *stacking*: place one diagram on top of another.

(**TODO**: some pictures)

The identity operator for multiplication is (**TODO**: diagram that looks like $||||$).

This algebra has some additional interesting structure.

- There's a trace $\mathcal{TL}_n \rightarrow \mathbb{C}$: given a Temperley-Lieb diagram, close up the embedded intervals in a process akin to a braid closure. Then **TODO**. (Also, **TODO**: a picture) I think there is a parameter δ , and if the result has n circles, we get δ^n .

- A $*$ -structure, by reversing the diagram horizontally.
- This defines a Hermitian form on TL_n , by $\langle x, y \rangle := \mathrm{tr}(y^* x)$. This is an inner product if $\delta \geq 2$.

Since the trace depends on δ , we will write $\mathrm{TL}_n(\delta)$ for the Temperley-Lieb algebra with trace given by δ .

There is an embedding $\mathrm{TL}_n \hookrightarrow \mathrm{TL}_{n+1}$, given by adding a single vertical interval on the right-hand side of a diagram. Call the colimit $\mathrm{TL}(\delta)$.

Exercise 4.2. Check that this inclusion respects multiplication, the identity, and the trace, assuming we use the same value of δ in both cases.

This is the basic example of a *planar algebra*. In general, a planar algebra is a collection of vector spaces V_0, V_1, V_2, \dots , together with an action by something called the *planar operad*. Fortunately, you don't need to know what an operad is to understand the planar operad. This operad is given by (TODO: in what sense?) *planar diagrams* (which TODO: I think are also called “spaghetti-and-meatballs diagrams”). These are diagrams of embeddings of compact 1-manifolds inside many-holed annuli, together with marked points on the boundaries of the annuli. These compose in a manner reminiscent of operator product expansion. TODO: a picture is more useful than a description here.

An action of the planar operad means, for each planar diagram, a multilinear map $\bigotimes V_i \rightarrow V_0$; we also ask for these maps to be compatible with compositions.

Example 4.3. The Temperley-Lieb algebra is a planar algebra, where the planar diagrams act by insertion. ◀

Example 4.4. The *graph planar algebra* on a simply laced graph Γ takes as its V_n the complex vector space spanned by the set of loops on Γ of length n .

There are a few different ways we can compose loops. Of course, we can concatenate loops with the same origin, as in algebraic topology; but there's another option. Assume both loops are of even length, and let p and p' be their respective halfway points; then, we can define their composition to be 0 if $p \neq p'$, and to be the first half of the first loop, then the second half of the second loop, if $p = p'$. TODO: I think that these two composition laws correspond to two planar diagrams, and these should give you the general story, so double-check this and then include those pictures.

TODO: planar diagrams enter this story somehow?

There's also a trace (TODO: picture goes here). This procedure is slightly ambiguous, so we simply sum up over all possibilities. ◀

We've seen in a few different talks so far the idea of a monoidal category, along with many variations of their definition.

Definition 4.5. A *monoidal category* is a category \mathcal{C} together with a functor $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ and a distinguished object $\mathbf{1} \in \mathcal{C}$ called the *unit*, together with data of an *associator*, a natural isomorphism $(- \otimes -) \otimes - \xrightarrow{\cong} - \otimes (- \otimes -)$ and *left and right unitors*, natural isomorphisms $\mathbf{1} \otimes - \xrightarrow{\cong} -$ and $- \otimes \mathbf{1} \xrightarrow{\cong} -$; these are subject to some coherence conditions.

The point of recalling this definition is that we'll relate it to all the pictures in not just this lecture, but also the other ones this week. This is a point that is often unclear to people — if you already know why you can do diagrammatics for various kinds of categories, it might feel not worth reviewing, but if not, it's certainly confusing.

The idea is, we can draw objects, morphisms, and equations in a monoidal category as diagrams in 2d. TODO: those diagrams.

- A morphism $f: A \rightarrow B$ is a box from a strand labeled by B to a strand labeled by A .
- Composition is stacking vertically.
- The tensor product is stacking horizontally, both of objects and of morphisms.
- The monoidal unit is the empty diagram.

Two diagrams which are related under planar isotopy are considered equal.

Theorem 4.6. A well-typed equation between morphisms in a monoidal category follows from the axioms of a monoidal category iff it holds true in the graphical language described above.

As a simple example, how do vertical and horizontal composition (namely, composition of morphisms, resp. tensor product) interact? If you do vertical, then horizontal, or horizontal, then vertical, you get the same

diagrams, and therefore they must be equal: given maps $f: A \rightarrow B$, $g: B \rightarrow C$, $h: D \rightarrow E$, and $k: E \rightarrow F$,

$$(4.7) \quad (f \circ g) \otimes (h \circ k) = (f \otimes h) \circ (g \otimes k)$$

as maps $A \otimes D \rightarrow C \otimes F$.

A monoidal category is *rigid* if it has left and right duals for all objects. Evaluation and coevaluation correspond to cups and caps; thus we obtain an identity (TODO: Zorro diagram, also called snake diagram). This allows us to freely do planar isotopy. (TODO: so do we need rigidity in order for Theorem 4.6 to hold?)

Now let \mathcal{C} be a rigid monoidal category and $X \in \mathcal{C}$; we will obtain a planar algebra by “zooming in” on this object X . Specifically, take $V_n := \text{End}(X^{\otimes n})$. (TODO: rest of data comes from diagrammatics, I think?)

Why care? Well, the formalism of planar algebras is different enough from that of monoidal categories to lend different tools to the study of things in their intersections. For example, monoidal categories and planar algebras have different notions of smallness. For example, in a semisimple rigid monoidal category, you might measure the number of simple objects. In a planar algebra generated by X as above, smallness is more traditionally measured with the *Frobenius-Perron dimension*. This can be understood in general semisimple rigid monoidal categories \mathcal{C} ; it is a map $K_0(\mathcal{C}) \rightarrow \mathbb{R}$ which is positive on simple eigenvalues. Specifically, suppose

$$(4.8) \quad X \otimes Y = \sum c_{XY}^Z Z,$$

where the sum is over isomorphism classes of simple objects Z of \mathcal{C} ; this defines a matrix in the entries X and Y ; its Frobenius-Perron eigenvalue is the Frobenius-Perron dimension of X .

Part 2. Tuesday, January 28

5. VICTOR OSTRIK, INTRODUCTION TO FUSION CATEGORIES, II

We begin by discussing diagrammatics for $\text{Vec}_{\mathbb{Z}/n}^\omega$, following work of Agustina Czenky.

The objects of $\text{Vec}_{\mathbb{Z}/n}^\omega$ are all tensor products of $1 \in \mathbb{Z}/n$, which in diagrammatics is denoted \uparrow ; thus, i is denoted $\uparrow \uparrow \dots \uparrow$. The generating morphisms are (TODO: two of them), and there are some relations: (TODO) and (TODO) are the identity, and (TODO) is ζ times (TODO), where ζ is some n^{th} root of unity determined by ω .⁵

This generators-and-relations description of $\text{Vec}_{\mathbb{Z}/n}^\omega$ allows us to uncover a universal property.

Proposition 5.1. *Let \mathcal{C} be a fusion category. Then there is a natural bijection between tensor functors $F: \text{Vec}_{\mathbb{Z}/n}^\omega \rightarrow \mathcal{C}$ and isomorphism classes of data of an object $X \in \mathcal{C}$ and an isomorphism $\theta: X^{\otimes n} \xrightarrow{\cong} \mathbf{1}$.*

The idea is that, looking at the diagrammatics of $X := F(1)$, we have two different isomorphisms $X^{\otimes(n+1)} \xrightarrow{\cong} X$, and one must be ζ times the other.

The slightly more sophisticated way to say this is that functors $\text{Vec}_{\mathbb{Z}/n}^\omega \rightarrow \mathcal{C}$ form a category, and the data (X, ϕ) as above forms a category, and the above bijection can be promoted to an equivalence of categories.

Example 5.2. Assume n is odd, so that we can use $2 \in \mathbb{Z}/n$ as a generator. Now you can compare (TODO: two diagrams), and one should be a multiple of another. It turns out the factor is ζ^4 , and this gives the action of $\text{Aut}(\mathbb{Z}/n)$ on $H^3(\mathbb{Z}/n; k^\times)$ from last time. \blacktriangleleft

Now let us use this to study tensor functors $F: \text{Vec}_{\mathbb{Z}/n}^\omega \rightarrow \mathcal{B}imod_R$, where R is a k -algebra. These are classified by (R, R) -bimodules X together with an isomorphism $X^{\otimes n} \xrightarrow{\cong} R$; in particular, X is invertible. Hence we can restrict our search to $\text{Pic}(R) \subset \mathcal{B}imod_R$, the subcategory of invertible bimodules. Inside $\text{Pic}(R)$, we also have $\text{Out}(R)$; as described last time, an outer automorphism defines an (isomorphism class of) (R, R) -bimodules.

Exercise 5.3. Assuming $Z(R) \cong k$, let $\theta \in \text{Out}(R)$. Show that $\theta(g) = \zeta g$ for $g \in R^\times$ and some root of unity ζ . If this is too difficult at first, take a look at some examples; see if you can give an example for any ζ .

One big open problem in this field is to classify all fusion categories. This is of course way too hard, given that it's more difficult than the classification of finite groups, but as with the classification of finite groups, intermediate results are interesting, possible, and useful.

⁵Conversely, any choice of ζ is given by some ω , though it's not always easy to work this out in practice.

Theorem 5.4 (Ocneanu rigidity (Etingof-Nikshych-Ostrik [ENO05])). *Fusion categories over \mathbb{C} have no deformations.*

This was originally conjectured by Ocneanu. We won't say precisely what a deformation of a fusion category is, but the data of associativity in a fusion category is matrices satisfying some equations, modulo the action of some symmetry group. Ocneanu rigidity amounts to there being only finitely many orbits of solutions under this group action.

Corollary 5.5. *There are countably many tensor equivalence classes of fusion categories over \mathbb{C} .*

For example, Vec_G^ω is classified by $H^3(G; \mathbb{C}^\times)$, which is a finite group.

Remark 5.6. Ocneanu rigidity is open in positive characteristic; the speaker expects it to be true, but for different reasons. The proof in characteristic zero uses a tool called *Davydov-Yetter cohomology* for fusion categories — this vanishes in characteristic zero, which implies Theorem 5.4, but is known to not vanish in characteristic p in general. ◀

Example 5.7. Another rich source of interesting examples of fusion categories are quantum groups at roots of unity. The construction is quite complicated. For example, fix an integer λ , called the *level*; then, one can build a fusion category $\mathcal{C}(\mathfrak{sl}_2, \lambda)$ as follows. There are $\lambda + 1$ simple objects L_0, \dots, L_λ , and the fusion rules are determined by

$$(5.8) \quad L_i \otimes L_1 = L_1 \otimes L_i = L_{i-1} \oplus L_{i+1},$$

where $L_{-1} = L_{\lambda+1} = 0$; this suffices to determine the rest of the fusion rules. This looks reminiscent of the representation theory of \mathfrak{sl}_2 , but “cut off” at λ . ◀

Any fusion category not equivalent to Vec_G^ω or a quantum group is called *exotic*. Examples of exotic fusion categories are constructed using subfactor theory; many are related to something called *near group categories*. Such a category has as its set of simple objects $G \amalg \{X\}$, where G is a finite group. The tensor product on elements of G is just multiplication, and the remaining rules are

$$(5.9a) \quad g \otimes X = X \otimes g = X$$

$$(5.9b) \quad X \otimes X = \bigoplus_{g \in G} g \oplus nX,$$

for some $n \in \mathbb{N}$.

It's not necessarily true that one can find an associator compatible with this data, but often one can. For $n = 0$, these are *Tambara-Yamagami categories*, which are relatively well-studied. For $n > 0$, Evans-Gannon [EG14] showed that either $n = \#G - 1$ or $\#G \mid n$. Moreover, if $n \geq \#G$, then G must be abelian, which is a nice simplification! Izumi-Tucker [IT19] considered the cases where $n = \#G - 1$ and $n = \#G$ — in the latter case, there are only finitely many examples, and for $n = 2\#G$, there's a single example with $G = \mathbb{Z}/3$. In general, it's open whether there are a finite number of examples for a fixed n as a function of $\#G$.

There are also various useful constructions which, given some fusion categories, produce more. One example is Deligne's tensor product $\mathcal{C} \boxtimes \mathcal{D}$ of fusion categories, which is again a fusion category.

Definition 5.10. An *associative algebra* in a fusion category \mathcal{C} is an object $A \in \mathcal{C}$ together with \mathcal{C} -morphisms $m: A \otimes A \rightarrow A$ and a unit $i: \mathbf{1} \rightarrow A$, subject to axioms guaranteeing associativity of m and that i is a unit for m .

Given an associative algebra A in \mathcal{C} , we can define A -module and (A, A) -bimodule objects in \mathcal{C} , analogous to the usual case.

Exercise 5.11. Write these definitions down.

The upshot is, given an associative algebra A in \mathcal{C} , the category of (A, A) -bimodule objects, denoted ${}_A\mathcal{C}_A$, is a tensor category. The monoidal product is \otimes_A , and the unit is A . This is generally not a fusion category — and indeed, even if A is just an algebra in vector spaces, its category of bimodules generally isn't fusion! But if A satisfies an assumption called *separability*, then ${}_A\mathcal{C}_A$ is semisimple and rigid, which is pretty close to being fusion — all we need is that $\mathbf{1} \in {}_A\mathcal{C}_A$ is indecomposable. A sufficient (but not necessary) condition for this is $\text{Hom}_{\mathcal{C}}(\mathbf{1}, A) \cong k$.

Remark 5.12. If we start with $\mathcal{C} = \text{Vec}_G^\omega$ and look for bimodule categories for algebras in \mathcal{C} , the fusion categories we obtain are called *group-theoretical fusion categories*. ◀

Another procedure to obtain fusion categories is called *graded extensions*, building a G -graded fusion category, where G is a finite group, out of an ungraded fusion category.

Here are two extremely useful tools for classifying fusion categories.

- (1) The Drinfeld center of a fusion category is a modular tensor category. Modular tensor categories have a lot of structure, and this is helpful for learning about fusion categories.
- (2) Diagrammatic methods are helpful, as we saw at the beginning of today's lecture.

REFERENCES

- [BK01] Bojko Bakalov and Alexander Kirillov, Jr. *Lectures on tensor categories and modular functors*, volume 21 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2001. 4
- [Del02] P. Deligne. Catégories tensorielles. volume 2, pages 227–248. 2002. Dedicated to Yuri I. Manin on the occasion of his 65th birthday. 8
- [EG14] David E. Evans and Terry Gannon. Near-group fusion categories and their doubles. *Adv. Math.*, 255:586–640, 2014. <https://arxiv.org/abs/1208.1500>. 11
- [EGNO15] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. *Tensor categories*, volume 205 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015. 4
- [ENO05] Pavel Etingof, Dmitri Nikshych, and Viktor Ostrik. On fusion categories. *Ann. of Math. (2)*, 162(2):581–642, 2005. 11
- [IT19] Masaki Izumi and Henry Tucker. Algebraic realization of noncommutative near-group fusion categories. 2019. <https://arxiv.org/abs/1908.01655>. 11
- [Jon99] Vaughan F. R. Jones. Planar algebras, I. 1999. <https://arxiv.org/abs/math/9909027>. 8