

Vitamin D₃ Protects against Cyclophosphamide-induced Neurotoxicity Via Modulation of Inflammatory Cytokines and Oxidative Stress Parameters

Dosunmu Damilare Paul¹, Babatunde Oluwatobiloba Bashirat², Onaolapo Olakunle James^{1,3}

1. Department of Pharmacology and Therapeutics, Ladoke Akintola University of Technology, Ogbomoso, Oyo State, Nigeria.

2. Department of Anatomy, Ladoke Akintola University of Technology, Ogbomoso, Oyo, State, Nigeria.

3. Behavioural Neuroscience and Neuropharmacology Unit, Department of Pharmacology and Therapeutics, Ladoke Akintola University of Technology, Ogbomoso, Oyo State, Nigeria

Correspondence:

Dr Dosunmu Damilare Paul.

Department of Pharmacology and Therapeutics,
Ladoke Akintola University of Technology,
Ogbomoso, Oyo State, Nigeria.

Email: damilaredvm@gmail.com. Tel: +2348165943694

ABSTRACT

Vitamin D₃'s antioxidative properties make it potentially beneficial in many conditions where oxidative stress is implicated in the pathological processes. The study investigated the possible protective effects of Vitamin D₃ on weight loss, feed intake, brain oxidative status, markers of inflammation and cerebral cortex histomorphology in rats administered Cyclophosphamide (CYP). Sixty rats were randomly assigned into six groups (n=10). Groups A and D served as Normal and Cyclophosphamide control respectively and were fed standard rat chow, groups B and E received Vitamin D₃ (300 IU/kg), while those in groups C and F received Vitamin D₃ (600 IU/kg). Animals in group A-C received intraperitoneal normal saline on day1, while groups D-F got intraperitoneal Cyclophosphamide (100 mg/kg/day on day1). Standard diet and Vitamin D₃ supplementation were administered daily for 15 days. After the experimental period, animals were euthanised and blood was taken for the assessment of biochemical parameters. Cerebral cortex was also processed for histological study. Vitamin D₃ mitigated CYP-induced weight loss. It also improved total antioxidant capacity while reducing CYP-induced lipid peroxidation. Interleukin-10 increased, while the CYP-induced increase in TNF- α were mitigated following Vitamin D₃ supplementation. Again, protective efforts in cerebral cortex histomorphology were observed. The results showed Vitamin D₃'s ability to protect against CYP-induced neurotoxicity in rats via modulation of inflammatory and oxidative stress parameters. It also highlighted Vitamin D₃'s potential as a possible adjunct in cancer chemotherapy-induced tissue damage. However, further research will be needed to specify its exact role in cancer chemotherapy.

KEYWORDS: Secosteroids; Chemotherapy; Inflammation, Histomorphology; Neuroprotection; Oxidative stress

1. Introduction

Cyclophosphamide (CYP) is an important chemotherapeutic, and efficient immunosuppressive agent that belongs to the class of alkylating agents [1-3]. Cyclophosphamide's therapeutic usage has also gained widespread acceptance in bone marrow transplantation [4-5] and autoimmune conditions like rheumatoid arthritis [1], systemic lupus erythematosus (SLE) (including treatment for widespread proliferative lupus nephritis, modifications in neurologic and psychiatric manifestations, enhancement of platelet counts; improvement of gastrointestinal vasculitis, interstitial lung disease and diffuse alveolar haemorrhage) [6-11]. Cyclophosphamide has also been applied in managing certain renal conditions such as nephritic syndrome resistance to corticosteroid and focal segmental glomerulonephritis [12, 13], it is also effective in

multiple sclerosis [3,14,15]. Other therapeutic applications include treating severe forms of other autoimmune diseases [1,11,16] and neuroblastoma [17,18].

Despite the significant pharmacological benefits of Cyclophosphamide, it has been associated with several adverse effects including neurotoxicity, bone marrow toxicity, bladder toxicity, cardiotoxicity, and gonadal damage among others, [19-24]. Though newer agents (platinum, taxanes, and targeted curatives) are sometimes used as substitutes for treating numerous solid tumours, cyclophosphamide is deemed relatively effective against many malignancies. This is often due to insufficient evidence for the superiority of the newer options over Cyclophosphamide usage [2]. However, the adverse effects associated with Cyclophosphamide have prompted researchers to hunt for adjuvants in form of drugs,

extracts, or supplements that have an ameliorative impact to mitigate these side effects.

Vitamin D is a fat-soluble secosteroid which is synthesised in two forms: Vitamin D₂ (ergocalciferol) and Vitamin D₃ (cholecalciferol) [25]. Plants and fungi produce Vitamin D₂ from ergosterol [26]; in contrast, human skin synthesises Vitamin D₃ from 7-dehydrocholesterol (7-DHC) upon exposure to UVB radiation that comes from sunlight [27] with contributing efforts from the liver [28] and the kidney [29]. Vitamin D₃ (1,25 (OH) 2D₃) is primarily known to maintain normal calcium levels and bone development by regulating phospho-calcium metabolism [30]. Given that Vitamin D₃ receptors (VDRs) are found in various tissues, including the prostate, brain, breast, pancreas, colon, and immune cells: it has been discovered to also play roles beyond bone health. Examples of such non-skeletal functions of Vitamin D₃ are: regulation of arterial blood pressure, prevention of cardiovascular complications, enhancement of immunological responses, regulation of insulin production to prevent diabetes, protection against certain cancers, and renoprotection. Vitamin D₃'s application includes the management of multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, and treatment-induced pain in cancer patients [31-33]. Also, approximately 3% of the human genome is regulated by Vitamin D₃ [34]. Several tissues, including pancreatic islets, the prostate, colon, breast, macrophages, and vascular smooth muscle cells, contain 1- α -hydroxylase, enabling them to produce Vitamin D₃ (1,25(OH)2D₃) locally [35]. This local synthesis positions Vitamin D₃ (1,25 (OH)2D₃) to have a role in cancer prevention and innate/adaptive immunity modulation, [36,37] as well as brain health [38]. Evidence from epidemiology studies linked Vitamin D₃ deficiency to increased risks of cancer, autoimmune diseases, diabetes, cardiovascular issues, and mental health disorders [39,40].

2. Materials and Methodology

2.1 Chemicals and Drugs

Vitamin D₃ tablets (1000IU) (Nature's field®), Cyclophosphamide injection (Endoxan Asta®), Normal saline.

2.2 Animals and Animal Welfare

Healthy Wistar rats used in this study were obtained from Empire breeders, Osogbo, Osun State, Nigeria. Rats were housed in wooden cages measuring 20 x 10 x 12 inches in temperature-controlled (22.5°C \pm 2.5°C) quarters with lights on at 7.00 am. Rats were allowed free access to food and water. All procedures were conducted in accordance with the approved protocols of the Faculty of Basic Medical Sciences, Ladoke Akintola University of Technology, within the

provisions for animal care and use prescribed in the scientific procedures on living animals, European Council Directive (EU2010/63).

2.2 Experimental Methods

A total of sixty Wistar rats weighing 190 – 200g each were used for the study. The rats were randomly assigned into six groups (n=10 each). Group A (Control Group) received a single intraperitoneal dose of Normal Saline on the first day and distilled water orally for 15 days. Group B (Vitamin D₃ Control Group1) received oral 300IU/kg/day of Vitamin D₃ for 15 days. Group C (Vitamin D₃ Control Group2) received 600IU/kg/day of Vitamin D₃ (oral) for 15 days [41,42]. Group D (Cyclophosphamide Control Group) received a single intraperitoneal (i.p.) dose of 100mg/kg Cyclophosphamide at the start of the experiment [43]. Group E (Cyclophosphamide + Vitamin D₃ Group1) received a single i.p. dose of 100mg/kg Cyclophosphamide on the first day and 300IU/kg/day of oral Vitamin D₃ for 15 days. Group F (Cyclophosphamide + Vitamin D₃ Group2) received a single i.p. dose of 100mg/kg Cyclophosphamide on the first day and 600IU/kg/day of oral Vitamin D₃ for 15 days. At the end of the experimental period, animals were euthanised by cervical dislocation and blood was taken through an intra-cardiac puncture for the levels of inflammatory markers (IL-10, TNF- α), lipid peroxidation and antioxidant status. The brain was removed, observed grossly, weighed and either homogenised or fixed in 10 % buffered formol-calcium. Sections of the cerebral cortex were processed for paraffin embedding, cut at 5 μ m and stained for histological study.

2.3 Determination of Body Weight and Feed Intake

Body weight and food intake were assessed weekly and daily respectively using an electronic weighing balance (Mettler Toledo Type BD6000, Greifensee, Switzerland) according to the protocol described [44]. Relative change in body weight or food intake was calculated for individual animals using the equation shown below, following which results for all animals were computed to find the statistical mean.

$$\frac{\text{Final body weight (or feed intake)} - \text{Initial body weight (or feed intake)}}{\text{Initial body weight (or feed intake)}} \times 100$$

2.4 Biochemical Tests

2.4.1 Lipid Peroxidation

Lipid peroxidation level was measured as Malondialdehyde content as described previously [45-47]. Change in colour was measured using a spectrophotometer at 532 nm.

2.4.2. Antioxidant Status

Total antioxidant capacity was determined using commercially available assay kit. Colour changes were measured as described previously [48,49].

2.4.3 Tumour Necrosis Factor- α and Interleukin (IL) -10

Tumour necrosis factor- α and interleukin IL-10 were measured using enzyme-linked immunosorbent assay (ELISA) techniques with commercially available kits (Enzo Life Sciences Inc. NY, USA) designed to measure the 'total' (bound and unbound) amount of the respective cytokines as previously described [50].

2.5 Tissue Histology

Rat brains were dissected, fixed in neutral-buffered formal-calcium. The cerebral cortex was removed and processed for paraffin-embedding, cut at 5 μ m and then stained with haematoxylin and eosin and cresyl fast violet for general histological study.

2.6 Photomicrography

Histologically processed sections of the cerebral cortex were examined microscopically using a Sellon Olympus trinocular microscope (XSZ-107E, China) with a digital camera, and photomicrographs were taken. Histopathological changes were assessed by a pathologist that was blind to the groupings.

2.7 Statistical Analysis

Data were analysed with Chris Rorden's ANOVA for windows (version 0.98). Data analysis was by One-way analysis of variance (ANOVA) and post-hoc test (Tukey HSD) was used for within and between group comparisons. Results were expressed as mean \pm S.E.M. and $p < 0.05$ was taken as the accepted level of significant difference from control.

3 Results

3.1 Effect of Vitamin D₃ on Body Weight

Figure 1 shows the effect of vitamin D₃ supplementation on percentage change in body weight in rats exposed to cyclophosphamide (CYP). There was a significant decrease in body weight in the CYP groups *CYP (-9.56%). CYP+VD300 (-3.33%) and CYP+ VD600 (-3.4%) compared to control (5.52%). Compared to CYP (-9.56%), there was a significant increase in body weight with CYP+VD300 (-3.33%) and CYP+ VD600 (-3.4%) respectively.

3.2 Effect of Vitamin D₃ on Feed Intake

Figure 2 shows the effect of vitamin D₃ supplementation on percentage change in feed intake in rats exposed to CYP. There

was a significant ($p < 0.05$) decrease in feed intake with CYP, CYP/VD 300 and 600 compared to control. Compared to CYP, feed intake increased with CYP/VD300 and CYP/VD600 respectively. Percentage change in feed intake decreased with CYP (2.17%). CYP+VD300 (4.02%) and CYP+VD600 (3.19%) as well as VD600 (5.93%) compared to control (12.61%). Compared to CYP (2.17%), feed intake increased with CYP+VD300 (4.02%) and CYP+VD600 (3.19%) respectively.




Figure 1: Effect of Vitamin D₃ (VD) on change in body weight in CYP treated rats. Each bar represents Mean \pm S.E.M, * $p < 0.05$ vs. control, # $p < 0.05$ significant difference from CYP, number of rats per treatment group =10. CONT: Control; CYP: Cyclophosphamide, VD: Vitamin D₃.




Figure 2: Effect of Vitamin D₃ (VD) on relative change in feed intake in CYP treated rats. Each bar represents Mean \pm S.E.M, * $p < 0.05$ vs. control, # $p < 0.05$ significant difference from CYP, number of rats per treatment group =10. CONT: Control; CYP: Cyclophosphamide, VD: Vitamin D₃.

3.3 Effect of Vitamin D₃ Supplementation on Oxidative Stress Parameters

Table 1 shows the effects of Vitamin D₃ supplementation on Total Antioxidant Capacity and Malondialdehyde levels in CYP-treated rats. Malondialdehyde (MDA) levels increased significantly with CYP, CYP/VD300 and CYP VD600 compared to control. Compared to CYP, MDA levels decreased

with CYP/VD300 and CYP/VD600. Total antioxidant capacity (TAC) increased significantly with VD300 and VD600 while it decreased significantly with CYP, CYP/VD300 and CYP/VD600 compared to control respectively. Compared to CYP, TAC increased with CYP/VD300 and CYP/VD600.

Table 1: Effect of Vitamin D₃ on Oxidative stress parameters

Groups	MDA (μmol/L)	TAC (mMTE)
Control	1.75 ± 0.02	4.58 ± 0.08
VD300	1.89 ± 0.39	5.04 ± 0.51
VD600	2.51 ± 0.17 *	5.23 ± 0.38
CYP	5.65 ± 0.3 *	2.06 ± 0.15 *
CYP/VD300	2.13 ± 0.05 *#	5.03 ± 0.17 #
CYP/VD600	1.68 ± 0.14 #	4.51 ± 0.01 #

Mean ± S.E.M, * p < 0.05 vs. control, #p < 0.05 significant difference from CYP, number of rats per treatment group = 10. CYP: Cyclophosphamide; VD: Vitamin D₃; MDA: Malondialdehyde; TAC: Total Antioxidant Capacity.

3.4 Effect of Vitamin D₃ supplementation on Levels of Inflammatory Cytokines

Table 2 shows the effect of vitamin D₃ supplementation on inflammatory markers; tumour necrosis factor-α and interleukin-10 in CYP-treated rats. Interleukin-10 levels increased with VD300, VD600 and CYP/VD600 while it decreased significantly with CYP, but slightly with CYP/VD300 compared to control. Compared to CYP, interleukin-10 levels increased significantly with CYP/VD300 and CYP/VD600. Tumour necrosis factor α levels increased significantly with CYP and decreased with CYP/VD600 compared to control. Compared to CYP, tumour necrosis factor α levels decreased significantly with CYP/VD300 and CYP/VD600.

3.5 Effect of Vitamin D₃ supplementation on the cerebral cortex histomorphology

Figure 3 show representative haematoxylin and eosin (figure 3a-f); while, figure 4 shows cresyl fast violet (figure 4a-f) stained sections of the rat cerebral cortex. Examination of the haematoxylin and eosin-stained slides showed features of the cerebral cortex with evidence of numerous granule cells, pyramidal cells and neuroglia, in rats administered vehicle (Fig. 5a-c).

Multipolar shaped pyramidal cells with large, rounded, vesicular nucleus are seen scattered throughout the neuropil; granular neurons with large open-faced nuclei, prominent nucleoli and scant cytoplasm are likewise observed. The pink-staining background which is the neuropil is better appreciated

in the cresyl fast violet-stained slides (figure 4a-c). In the group administered CYP, degenerating pyramidal cells with pale-staining nuclei, joined cells and granule cells are observed, these features are in keeping with neuronal injury. In groups supplemented with Vitamin D₃ (3d-e) was observed a protection against CYP-induced neuronal injury. Examination of cresyl-fast violet-stained sections of the cerebrum revealed well-delineated multipolar pyramidal cells, deeply-staining granule cells and neuroglia in the control group (Figure 4a) and groups administered Vitamin D₃ (Fig. 4b, 4c). However, degenerating pyramidal cells with pale staining nuclei were observed in the group administered CYP (Figure 4d), while a protection against neuronal injury was observed in the groups with Vitamin D₃ supplementation (Figure 4e, 4f).

Table 2: Effect of Vitamin D₃ on inflammatory cytokines

Groups	Interleukin-10 (pg/ml)	TNF-α (pg/ml)
Control	51.42 ± 14.59	515.25 ± 203.6
VD300	47.22 ± 12.67	473.42 ± 68.49
VD600	42.29 ± 2.86	372.01 ± 117.52
CYP	18.54 ± 1.8 *	982.01 ± 0.65 *
CYP/VD300	31.93 ± 8.95 #	536.46 ± 106.34 #
CYP/VD600	51.16 ± 1.51 #	491.02 ± 98.41 #

Mean ± S.E.M, * p < 0.05 vs. control, #p < 0.05 significant difference from CYP, number of rats per treatment group = 10. CYP: Cyclophosphamide; VD: Vitamin D₃; IL-10: Interleukin-10; TNF-α: Tumour Necrosis Factor-α

4 Discussion

This study investigated the possible protective effects of vitamin D₃ supplementation on changes in biochemical and histomorphological markers of brain integrity, oxidant-antioxidant status, appetite, weight, and markers of inflammation in healthy rats administered Cyclophosphamide. The results showed that vitamin D₃ protected against Cyclophosphamide-induced changes in weight, feed consumption, antioxidant status, and inflammatory markers and cerebral morphology.

In this study, weekly body weight and percentage change in weight increased in the control group and decreased significantly in the groups administered Cyclophosphamide. The effects on body weight observed with Cyclophosphamide (CYP) in this study are consistent with the results of a number of other studies that had also reported that the administration of CYP caused significant weight loss [51]. On the other hand, treatment with vitamin D₃ mitigated CYP-induced weight loss. Vitamin D₃ is a secosteroid which plays a crucial role in

calcium homeostasis and bone health, but it has also been studied for its potential effects on body weight and metabolism [52-54]. Results of studies have varied significantly, with a few reporting vitamin D₃'s abilities to mitigate weight gain [52], others observed no effect on body weight in normal sized animals [55] while other studies identified reduction in feed intake with Vitamin D₃ administration [56].




Figure 3: Photomicrographs of cerebral cortex sections stained with haematoxylin and eosin. Stained slides revealed distinct layers of the cerebral cortex with presence of numerous granule cells (Gc, black arrows), pyramidal cells (Pc, red arrows) and neuroglia (Ng, yellow arrows) in rats administered vehicle (a-c). Multipolar shaped pyramidal cells with large, rounded, vesicular nuclei are seen scattered throughout the neuropil; granular neurons with large open-faced nuclei, prominent nucleoli (more in D) and scant cytoplasm are also observed. **Mag. X 100.**




Figure 4: Photomicrographs of cerebral cortex-stained sections by cresyl fast violet. The pink-staining background which is the neuropil is better appreciated in the cresyl fast violet-stained slides (a-c). In the group administered CYP (d)

numerous degenerating pyramidal and granule cells are observed, degenerating neurons are pale staining with shrunken pale staining nuclei, these features are in keeping with neuronal injury, numerous neuroglial are also seen as indication of inflammatory responses. In groups treated with vitamin D₃ (e-f) was observed a protection against CYP-induced neuronal injury. **Mag. X 100.**

In this study, groups administered vitamin D₃ alone showed no significant difference in body weight from control, supporting studies that had reported such effects. This would suggest that vitamin D₃'s effect on weight only came into effect in situations of body weight abnormalities like disease related weight loss or conditions of obesity. As also observed in this study (with CYP related weight loss), vitamin D₃'s ability to reduce lipid peroxidation, increase antioxidant levels, and modulate the immune/inflammation system, via regulating the production of proinflammatory cytokines and inhibiting the proliferation of proinflammatory cells [57-60] are possible mechanisms through which it reversed CYP-induced weight loss. The administration of CYP in this study was associated with a significant reduction in feed intake while treatment with vitamin D₃ generally mitigated CYP induced decrease in feed intake. In groups administered vitamin D₃ alone, no significant difference in feed intake was observed, supporting earlier observations that vitamin D₃'s effect on the body was to restore normalcy. A few studies have also reported that vitamin D₃ did not increase muscle strength or power, energy consumption, resting metabolic rate and body composition [61]. In the CYP groups, vitamin D₃ supplementation reduced CYP induced decrease in feed intake and weight loss, consistent with the results of other studies that had revealed that lack of vitamin D₃ receptor (VDR) led to a lean phenotype with reduced body weight, decreased mass of white adipose tissue, promoted rate of β -oxidation, and enhanced energy expenditure leading to weight loss [62]; which Vitamin D₃ supplementation could potentially prevent, according to this study. Vitamin D₃'s effects on body weight and feed intake can also be attributed to its ability to influence central processes that regulate energy homeostasis [61,63]. There have also been reports that vitamin D₃ can scavenge superoxide, hydroxyl, peroxy and alkoxy radicals in tissues [59, 64], leading to a decrease in lipid peroxidation level and an increase in the level of the total antioxidant capacity. Studies have continued to report the benefits of vitamin D₃ in modulating inflammatory activity and in protecting against brain injury as well as that of other organs [58-66]. The beneficial effects of vitamin D₃ could be attributed to a decrease in proinflammatory cytokine (TNF- α) [67]; while levels of anti-inflammatory cytokines (IL-10) increased [68] similar to this study. The neuroprotective effects of vitamin D₃ have also been linked to its ability to inhibit the formation of malondialdehyde, scavenge reactive oxygen species. There have been reports that the ability of Cyclophosphamide to cross the blood brain barrier and induce oxidative stress is one of the many ways through which it induces neurotoxicity [69]. Also,

in this study, Cyclophosphamide was associated with neuronal injury and an increase in migrating neuroglial cells, a significant indicator of cerebral inflammation, as previously reported [70] following Cyclophosphamide administration. Vitamin D₃'s ability to protect against and reduce brain injury has been reported severally [65,71,66,72]. While most studies have examined its benefits in dementia and Alzheimer's disease [73-75], only a few studies have examined its potential benefits in chemotherapy induced neurotoxicity [76]; hence this study serves as an eye-opener to more potential effects of Vitamin D₃ in chemotherapy induced neurotoxicity.

Conclusion

Vitamin D₃ supplementation causes significant dose-related modifications in antioxidative stress markers (malondialdehyde, total antioxidant capacity) and inflammatory cytokines (tumor necrosis factor- α (TNF- α), and interleukin-10) in cyclophosphamide-induced neurotoxicity in rats. Considering its ever-increasing potential clinical applications and natural sources, these study findings would assist in making objective appraisals of the protective effect of vitamin D₃ in the course of Cyclophosphamide chemotherapy.

Acknowledgements

None

Funding.

None

Availability of data and materials

Data generated during and analysed during the course of this study are available from the corresponding author on request.

Declaration of ethical approval

Ethical approval for this study was granted by the Ethical Committee of the Faculty of Basic Medical Sciences ERC/FBMS/038/2024

Competing interests

None

Received 2nd December, 2024, Revised 31st December, 2024, Accepted 4th January, 2025,

Published online 25/01/25

References

1. Iqbal A, Iqbal MK, Sharma S, Ansari MohdA, Najmi AK, Ali SM, et al. Molecular mechanism involved in cyclophosphamide-induced cardiotoxicity: Old drug with a new vision. *Life Sciences* [Internet]. 2018 Dec 12;218:112–31. <https://doi.org/10.1016/j.lfs.2018.12.018>
2. Emadi A, Jones RJ, Brodsky RA. Cyclophosphamide and cancer: golden anniversary. *Nature Reviews Clinical Oncology* [Internet]. 2009 Sep 29;6(11):638–47. <https://doi.org/10.1038/nrclinonc.2009.146>
3. Weiner HL, Cohen JA. Treatment of multiple sclerosis with cyclophosphamide: critical review of clinical and immunologic effects. *Multiple Sclerosis Journal* [Internet]. 2002 Apr 1;8(2):142–54. <https://doi.org/10.1191/1352458502ms790oa>
4. Montoro J, Boumendil A, Finel H, Bramanti S, Castagna L, Blaise D, et al. Post-Transplantation Cyclophosphamide-Based Graft-versus-Host Disease Prophylaxis in HLA-Matched and Haploidentical Donor Transplantation for Patients with Hodgkin Lymphoma: A Comparative Study of the Lymphoma Working Party of the European Society for Blood and Marrow Transplantation. *Transplantation and Cellular Therapy* [Internet]. 2023 Dec 1;30(2):210.e1–210.e14. <https://doi.org/10.1016/j.jtct.2023.11.021>
5. Paviglianiti A, Ngoya M, Peña M, Boumendil A, Gülbas Z, Ciceri F, et al. Graft-versus-host-disease prophylaxis with ATG or PTCY in patients with lymphoproliferative disorders undergoing reduced intensity conditioning regimen HCT from one antigen mismatched unrelated donor. *Bone Marrow Transplantation* [Internet]. 2024 Feb 8;59(5):597–603. <https://doi.org/10.1038/s41409-024-02225-2>
6. Drie T, Alsamman MI, Tarcha R, Haidar G, Kudsi M. Successful pregnancy after cyclophosphamide therapy for systemic lupus erythematosus: A case report. *Annals of Medicine and Surgery* [Internet]. 2024 Jan 3; <https://doi.org/10.1097/ms9.0000000000001641>
7. Elaziz MMA, Gamal SM, Fayed A, Abu-Zaid MH, Ghoniem SA, Teieb DA. High- and low-dose cyclophosphamide in Egyptian lupus nephritis patients: a multicenter retrospective analysis. *Zeitschrift Für Rheumatologie* [Internet]. 2023 Aug 15;83(S1):115–23. <https://doi.org/10.1007/s00393-023-01386-7>
8. Martínez ADS, Ortiz MJT, Gómez LVZ, Llano CDM. Chronic inflammatory demyelinating polyneuropathy as the initial presentation of systemic lupus erythematosus successfully treated with cyclophosphamide. *Cureus* [Internet]. 2024 Jan 4; <https://doi.org/10.7759/cureus.5164>
9. Rengasamy P. Congenital malformations attributed to prenatal exposure to cyclophosphamide. *Anti-Cancer Agents in Medicinal Chemistry* [Internet]. 2017 Mar 2;17(9). <https://doi.org/10.2174/1871520616666161206150421>
10. Lehman TJA, Onel K. Intermittent intravenous cyclophosphamide arrests progression of the renal chronicity index in childhood systemic lupus erythematosus. *The Journal of Pediatrics* [Internet]. 2000 Feb 1;136(2):243–7. [https://doi.org/10.1016/s0022-3476\(00\)70109-0](https://doi.org/10.1016/s0022-3476(00)70109-0)
11. Ortmann RA, Klippel JH. update on cyclophosphamide for systemic lupus erythematosus. *Rheumatic Disease Clinics of North America* [Internet]. 2000 May 1;26(2):363–75. [https://doi.org/10.1016/s0889-857x\(05\)70143-5](https://doi.org/10.1016/s0889-857x(05)70143-5)
12. Linares-Fernández BE, Alfieri AB. Cyclophosphamide induced cystitis: role of nitric oxide synthase, cyclooxygenase-1 and 2, and NK 1 receptors. *The Journal of Urology* [Internet]. 2007 Mar 24;177(4):1531–6. <https://doi.org/10.1016/j.juro.2006.11.072>
13. Wong TM, Yeo W, Chan LW, Mok TSK. Hemorrhagic pyelitis, ureteritis, and cystitis secondary to cyclophosphamide: case report and Review of the literature. *Gynecologic Oncology* [Internet]. 2000 Feb 1;76(2):223–5. <https://doi.org/10.1006/gyno.1999.5680>
14. Rinaldi L, Perini P, Calabrese M, Gallo P. Cyclophosphamide as second-line therapy in multiple sclerosis: benefits and risks. *Neurological Sciences* [Internet]. 2009 Oct 1;30(S2):171–3. <https://doi.org/10.1007/s10072-009-0145-4>

15. Giesser BS. Primer on multiple sclerosis. Oxford University Press; 2016.
16. Kim J, Chan JJ. Cyclophosphamide in dermatology. Australasian Journal of Dermatology [Internet]. 2016 Jan 24;58(1):5–17. <https://doi.org/10.1111/ajd.12406>
17. Mangó K, Fekete F, Kiss ÁF, Erdős R, Fekete JT, Búdi T, et al. Association between CYP2B6 genetic variability and cyclophosphamide therapy in pediatric patients with neuroblastoma. Scientific Reports [Internet]. 2023 Jul 21;13(1):. <https://doi.org/10.1038/s41598-023-38983-0>
18. McCune JS, Salinger DH, Vicini P, Oglesby C, Blough DK, Park JR. Population Pharmacokinetics of cyclophosphamide and metabolites in children with neuroblastoma: a report from the Children's Oncology Group. The Journal of Clinical Pharmacology [Internet]. 2008 Oct 17;49(1):88–102. <https://doi.org/10.1177/0091270008325928>
19. Papanicolaou GA, Chen M, He N, Martens MJ, Kim S, Batista MV, et al. Incidence and impact of fungal infections in Post-Transplantation Cyclophosphamide-Based Graft-versus-Host disease prophylaxis and haploidentical hematopoietic cell transplantation: A Center for International Blood and Marrow Transplant Research analysis. Transplantation and Cellular Therapy [Internet]. 2023 Sep 28;30(1):114.e1-114.e16. <https://doi.org/10.1016/j.jtct.2023.09.017>
20. Iqbal A, Syed MA, Haque MM, Najmi AK, Ali J, Haque SE. Effect of nerolidol on cyclophosphamide-induced bone marrow and hematologic toxicity in Swiss albino mice. Experimental Hematology [Internet]. 2020 Jan 24;82:24–32. <https://doi.org/10.1016/j.exphem.2020.01.007>
21. Komolafe OA, Arayombo BE, Abiodun AA, Saka OS, Abijo AZ, Ojo SK, et al. Immunohistochemical and histological evaluations of cyclophosphamide-induced acute cardiotoxicity in wistar rats: The role of turmeric extract (curcuma). Morphologie [Internet]. 2020 Jan 9;104(345):133–42. <https://doi.org/10.1016/j.morpho.2019.10.047>
22. Barnes H, Holland AE, Westall GP, Goh NS, Glaspole IN. Cyclophosphamide for connective tissue disease-associated interstitial lung disease. Cochrane Library [Internet]. 2018 Jan 3;2018(1). <https://doi.org/10.1002/14651858.cd010908.pub2>
23. Clifton K, Barnett C, Martinez A, Karuturi M, Saleem S, Litton J. Severe hyponatremia following cyclophosphamide infusion in breast cancer patients. The Breast Journal [Internet]. 2018 Mar 2;24(4):691–2. <https://doi.org/10.1111/tbj.13026>
24. Ghobadi E, Moloudizargari M, Asghari MH, Abdollahi M. The mechanisms of cyclophosphamide-induced testicular toxicity and the protective agents. Expert Opinion on Drug Metabolism & Toxicology [Internet]. 2016 Dec 24;13(5):525–36. <https://doi.org/10.1080/17425255.2017.1277205>
25. Bikle DD. Vitamin D and the skin: Physiology and pathophysiology. Reviews in Endocrine and Metabolic Disorders [Internet]. 2011 Aug 17;13(1):3–19. <https://doi.org/10.1007/s11154-011-9194-0>
26. Jäpelt RB, Jakobsen J. Vitamin D in plants: a review of occurrence, analysis, and biosynthesis. Frontiers in Plant Science [Internet]. 2013 Jan 1;4. <https://doi.org/10.3389/fpls.2013.00136>
27. Bikle D, Christakos S. New aspects of vitamin D metabolism and action — addressing the skin as source and target. Nature Reviews Endocrinology [Internet]. 2020 Feb 6;16(4):234–52. <https://doi.org/10.1038/s41574-019-0312-5>
28. Zhu JG, Ochalek JT, Kaufmann M, Jones G, DeLuca HF. CYP2R1 is a major, but not exclusive, contributor to 25-hydroxyvitamin D production in vivo. Proceedings of the National Academy of Sciences [Internet]. 2013 Sep 9;110(39):15650–5. <https://doi.org/10.1073/pnas.1315006110>
29. Bikle DD, Patzek S, Wang Y. Physiologic and pathophysiologic roles of extra renal CYP27b1: Case report and review. Bone Reports [Internet]. 2018 Feb 26;8:255–67. <https://doi.org/10.1016/j.bonr.2018.02.004>
30. Delrue C, Speeckaert MM. Vitamin D and Vitamin D-Binding protein in health and disease. International Journal of Molecular Sciences [Internet]. 2023 Feb 28;24(5):4642. <https://doi.org/10.3390/ijms24054642>
31. Zarrati M, Sohoul M, Aleayyub S, Keshavarz N, Razmpoosh E, Găman MA, et al. The Effect of Vitamin D Supplementation on Treatment-Induced Pain in Cancer Patients: A Systematic review. Pain Management Nursing [Internet]. 2022 Mar 10;23(4):458–66. <https://doi.org/10.1016/j.pmn.2022.02.001>
32. Jagannath VA, Filippini G, Nascimento IJBD, Di Pietrantonj C, Robak EW, Whamond L. Vitamin D for the management of multiple sclerosis. Cochrane Library [Internet]. 2018 Sep 24;2023(6). <https://doi.org/10.1002/14651858.cd008422.pub3>
33. Urena-Torres P, Souberbielle J. Pharmacologic role of vitamin D natural products. Current Vascular Pharmacology [Internet]. 2014 Apr 1;12(2):278–85. <https://doi.org/10.2174/15701611113119990020>
34. Bouillon R, Carmeliet G, Verlinden L, Van Etten E, Verstuyf A, Luderer HF, et al. Vitamin D and Human Health: Lessons from Vitamin D Receptor Null Mice. Endocrine Reviews [Internet]. 2008 Oct 1;29(6):726–76. <https://doi.org/10.1210/er.2008-0004>
35. Andress DL. Vitamin D in chronic kidney disease: A systemic role for selective vitamin D receptor activation. Kidney International [Internet]. 2005 Dec 22;69(1):33–43. <https://doi.org/10.1038/sj.ki.5000045>
36. Li YC, Kong J, Wei M, Chen ZF, Liu SQ, Cao LP. 1,25-Dihydroxyvitamin D3 is a negative endocrine regulator of the renin-angiotensin system. Journal of Clinical Investigation [Internet]. 2002 Jul 15;110(2):229–38. <https://doi.org/10.1172/jci15219>
37. Holick MF. Vitamin D deficiency. New England Journal of Medicine [Internet]. 2007 Jul 18;357(3):266–81. <https://doi.org/10.1056/nejmra070553>
38. Banafshe HR, Khoshnoud MJ, Abed A, Saghaizadeh M, Mesdaghinia A. Vitamin D supplementation attenuates the behavioral scores of neuropathic pain in rats. Nutritional Neuroscience [Internet]. 2018 Feb 12;22(10):700–5. <https://doi.org/10.1080/1028415x.2018.1435485>
39. Holick MF. VITAMIN D IN HEALTH AND DISEASE: Vitamin D for health and in chronic kidney disease. Seminars in Dialysis [Internet]. 2005 Jul 1;18(4):266–75. <https://doi.org/10.1111/j.1525-139x.2005.18402.x>
40. Giovannucci E. Vitamin D and cardiovascular disease. Current Atherosclerosis Reports [Internet]. 2009 Nov 1;11(6):456–61. <https://doi.org/10.1007/s11883-009-0068-9>
41. Korewo-Labelle D, Karnia MJ, Myślińska D, Kaczor JJ. Supplementation with Vitamin D3 Protects against Mitochondrial Dysfunction and Loss of BDNF-Mediated Akt Activity in the Hippocampus during Long-Term Dexamethasone Treatment in Rats. International Journal of Molecular Sciences [Internet]. 2023 Sep 11;24(18):13941. <https://doi.org/10.3390/ijms241813941>

42. Dabak DO, Kuloglu T, Ozercan MR. Effects of vitamin D3(Cholecalciferol) on Adriamycin-Induced nephrotoxicity. Renal Failure [Internet]. 2009 Jan 1;31(5):400–5. <https://doi.org/10.1080/08860220902883020>
43. Terpinskaya TI. Dynamics of Antitumor Resistance after Cyclophosphamide Injection. Bulletin of Experimental Biology and Medicine [Internet]. 2012 Mar 1;152(5):613–4. <https://doi.org/10.1007/s10517-012-1589-8>
44. Erdenebayar O, Kato T, Kawakita T, Kasai K, Kadota Y, Yoshida K, et al. Effects of peripheral oxytocin administration on body weight, food intake, adipocytes, and biochemical parameters in peri- and postmenopausal female rats. Endocrine Journal [Internet]. 2020 Sep 1;68(1):7–16. <https://doi.org/10.1507/endocri.ej19-0586>
45. Onaolapo OJ, Odeniyi AO, Jonathan SO, Samuel MO, Amadiogwu D, Olawale A, et al. An investigation of the Anti-Parkinsonism potential of co-enzyme Q10 and co-enzyme Q10 /Levodopa-carbidopa combination in mice. Current Aging Science [Internet]. 2019 Nov 8;14(1):62–75. <https://doi.org/10.2174/1874609812666191023153724>
46. Olofinnade AT, Onaolapo TM, Oladimeji S, Fatoki AM, Balogun CI, Onaolapo AY, Onaolapo OJ. An Evaluation of the Effects of Pyridoxal Phosphate in Chlorpromazineinduced Parkinsonism using Mice. Cent Nerv Syst Agents Med Chem. 2020;20(1):13–25. doi: 10.2174/1871524920666200120142508.
47. Olofinnade AT, Onaolapo AY, Onaolapo OJ, Olowe OA. Hazelnut Modulates Neurobehaviour and Ameliorates Ageing-induced Oxidative Stress, and Caspase-3- Mediated Apoptosis in Mice. Curr Aging Sci. 2021;14(2):154–162. doi: 10.2174/1874609813666201228112349.
48. Olofinnade AT, Onaolapo AY, Onaolapo OJ, Olowe OA, Mollica A, Zengin G, Stefanucci A. Corylus avellana L. modulates neurobehaviour and brain chemistry following high-fat diet. Front Biosci (Landmark Ed). 2021 Jan 1;26(3):537–551. doi: 10.2741/4906.
49. Onaolapo OJ, Jegede OR, Adegoke O, Ayinde MO, Akeredolu OM, Onaolapo AY. Dietary zinc supplement militates against ketamine-induced behaviours by agedependent modulation of oxidative stress and acetylcholinesterase activity in mice. Pharmacol Rep. 2020 Feb;72(1):55–66. doi: 10.1007/s43440-019-00003-2.
50. Bienvenu J, Monneret G, Fabien N, Revillard JP. The clinical usefulness of the measurement of cytokines. Clinical Chemistry and Laboratory Medicine (CCLM) [Internet]. 2000 Apr 30;38(4):267–85. <https://doi.org/10.1515/cclm.2000.040>
51. Khazaei F, Ghanbari E, Khazaei M. Protective Effect of Royal Jelly against Cyclophosphamide-Induced Thrombocytopenia and Spleen and Bone Marrow Damages in Rats. DOAJ (DOAJ: Directory of Open Access Journals) [Internet]. 2020 Oct 1;22(3):302–9. <https://doaj.org/article/834c54cf77c64f0d8727ea4c74ca70b6>
52. Soares MJ, Murhadi LL, Kurpad AV, Ping-Delfos WLCS, Piers LS. Mechanistic roles for calcium and vitamin D in the regulation of body weight. Obesity Reviews [Internet]. 2012 Mar 2;13(7):592–605. <https://doi.org/10.1111/j.1467-789x.2012.00986.x>
53. Mason C, Xiao L, Imayama I, Duggan C, Wang CY, Korde L, et al. Vitamin D3 supplementation during weight loss: a double-blind randomized controlled trial. American Journal of Clinical Nutrition [Internet]. 2014 Apr 1;99(5):1015–25. <https://doi.org/10.3945/ajcn.113.073734>
54. Tobias DK, Luttmann-Gibson H, Mora S, Danik J, Bubes V, Copeland T, et al. Association of body weight with response to vitamin D supplementation and metabolism. JAMA Network Open [Internet]. 2023 Jan 17;6(1):e2250681. <https://doi.org/10.1001/jamanetworkopen.2022.50681>
55. Ekwaru JP, Zwicker JD, Holick MF, Giovannucci E, Veugeliers PJ. The importance of body weight for the dose response relationship of oral vitamin D supplementation and serum 25-Hydroxyvitamin D in healthy volunteers. PLoS ONE [Internet]. 2014 Nov 5;9(11):e111265. <https://doi.org/10.1371/journal.pone.0111265>
56. Karges K, Brooks JC, Gill DR, Breazile JE, Owens FN, Morgan JB. Effects of supplemental vitamin D3 on feed intake, carcass characteristics, tenderness, and muscle properties of beef steers. Journal of Animal Science [Internet]. 2001 Jan 1;79(11):2844. <https://doi.org/10.2527/2001.79112844x>
57. Agrawal D, Yin K. Vitamin D and inflammatory diseases. Journal of Inflammation Research [Internet]. 2014 May 1;69. <https://doi.org/10.2147/jir.s63898>
58. Ke CY, Yang FL, Wu WT, Chung CH, Lee RP, Yang WT, et al. Vitamin D3 reduces tissue damage and oxidative stress caused by exhaustive exercise. International Journal of Medical Sciences [Internet]. 2016 Jan 1;13(2):147–53. <https://doi.org/10.7150/ijms.13746>
59. Sepidarkish M, Farsi F, Akbari-Fakhrabadi M, Namazi N, Almasi-Hashiani A, Hagiagha AM, et al. The effect of vitamin D supplementation on oxidative stress parameters: A systematic review and meta-analysis of clinical trials. Pharmacological Research [Internet]. 2018 Nov 15;139:141–52. <https://doi.org/10.1016/j.phrs.2018.11.011>
60. Mastali VP, Hoseini R, Azizi M. The effect of short-term vitamin D on the antioxidant capacity following exhaustive aerobic exercise. African Health Sciences [Internet]. 2023 Apr 11;23(1):584–91. <https://doi.org/10.4314/ahs.v23i1.61>
61. Montenegro KR, Cruzat V, Melder H, Jacques A, Newsholme P, Ducker KJ. Vitamin D supplementation does not impact resting metabolic rate, body composition and strength in vitamin D sufficient physically active adults. Nutrients [Internet]. 2020 Oct 12;12(10):3111. <https://doi.org/10.3390/nu12103111>
62. Narvaez CJ, Matthews D, Broun E, Chan M, Welsh J. Lean Phenotype and Resistance to Diet-Induced Obesity in Vitamin D Receptor Knockout Mice Correlates with Induction of Uncoupling Protein-1 in White Adipose Tissue. Endocrinology [Internet]. 2008 Oct 10;150(2):651–61. <https://doi.org/10.1210/en.2008-1118>
63. Su H, Liu N, Zhang Y, Kong J. Vitamin D/VDR regulates peripheral energy homeostasis via central renin-angiotensin system. Journal of Advanced Research [Internet]. 2021 Feb 3;33:69–80. <https://doi.org/10.1016/j.jare.2021.01.011>
64. Razzaghi R, Pourbagheri H, Momen-Heravi M, Bahmani F, Shadi J, Soleimani Z, et al. The effects of vitamin D supplementation on wound healing and metabolic status in patients with diabetic foot ulcer: A randomized, double-blind, placebo-controlled trial. Journal of Diabetes and Its Complications [Internet]. 2016 Jun 25;31(4):766–72. <https://doi.org/10.1016/j.jdiacomp.2016.06.017>
65. Evans MA, Kim HA, Ling YH, Uong S, Vinh A, De Silva TM, et al. Vitamin D3 Supplementation Reduces Subsequent Brain Injury and Inflammation Associated with Ischemic Stroke. NeuroMolecular Medicine [Internet]. 2018 Feb 23;20(1):147–59. <https://doi.org/10.1007/s12017-018-8484-z>
66. Jiang H, Yang X, Wang Y, Zhou C. Vitamin D Protects against Traumatic Brain Injury via Modulating TLR4/MyD88/NF-κB Pathway-Mediated Microglial Polarization and Neuroinflammation. BioMed Research International [Internet]. 2022 Jul 15;2022:1–12. <https://doi.org/10.1155/2022/3363036>
67. Zhang Y, Leung DYM, Richers BN, Liu Y, Remigio LK, Riches DW, et al. Vitamin D inhibits Monocyte/Macrophage

- proinflammatory cytokine production by targeting MAPK phosphatase-1. *The Journal of Immunology* [Internet]. 2012 Feb 2;188(5):2127–35. <https://doi.org/10.4049/jimmunol.1102412>
68. Wöbke TK, Sorg BL, Steinhilber D. Vitamin D in inflammatory diseases. *Frontiers in Physiology* [Internet]. 2014 Jul 2;5. <https://doi.org/10.3389/fphys.2014.00244>
 69. Singh S, Kumar A. Protective effect of edaravone on cyclophosphamide induced oxidative stress and neurotoxicity in rats. *Current Drug Safety* [Internet]. 2019 May 7;14(3):209–16. <https://doi.org/10.2174/1574886314666190506100717>
 70. Dong YS, Wang JL, Feng DY, Qin HZ, Wen H, Yin ZM, et al. Protective Effect of Quercetin against Oxidative Stress and Brain Edema in an Experimental Rat Model of Subarachnoid Hemorrhage. *International Journal of Medical Sciences* [Internet]. 2014 Jan 1;11(3):282–90. <https://doi.org/10.7150/ijms.7634>
 71. Sharma S, Kumar A, Choudhary A, Sharma S, Khurana L, Sharma N, et al. Neuroprotective role of oral vitamin D supplementation on consciousness and inflammatory biomarkers in determining severity outcome in acute traumatic brain injury patients: a Double-Blind randomized clinical trial. *Clinical Drug Investigation* [Internet]. 2020 Mar 14;40(4):327–34. <https://doi.org/10.1007/s40261-020-00896-5>
 72. Lasoń W, Jantas D, Leśkiewicz M, Regulska M, Basta-Kaim A. Vitamin D3 and Ischemic Stroke: A Narrative review. *Antioxidants* [Internet]. 2022 Oct 27;11(11):2120. <https://doi.org/10.3390/antiox11112120>
 73. Littlejohns TJ, Henley WE, Lang IA, Annweiler C, Beauchet O, Chaves PHM, et al. Vitamin D and the risk of dementia and Alzheimer disease. *Neurology* [Internet]. 2014 Aug 7;83(10):920–8. <https://doi.org/10.1212/wnl.0000000000000755>
 74. Shen L, Ji HF. Vitamin D deficiency is associated with increased risk of Alzheimer's disease and dementia: evidence from meta-analysis. *Nutrition Journal* [Internet]. 2015 Jul 31;14(1). <https://doi.org/10.1186/s12937-015-0063-7>
 75. Lin CI, Chang YC, Kao NJ, Lee WJ, Cross TW, Lin SH. 1,25(OH)2D3 Alleviates Aβ(25-35)-Induced Tau Hyperphosphorylation, Excessive Reactive Oxygen Species, and Apoptosis Through Interplay with Glial Cell Line-Derived Neurotrophic Factor Signaling in SH-SY5Y Cells. *International Journal of Molecular Sciences* [Internet]. 2020 Jun 13;21(12):4215. <https://doi.org/10.3390/ijms21124215>
 76. Jennaro TS, Fang F, Kidwell KM, Smith EML, Vangipuram K, Burness ML, et al. Vitamin D deficiency increases severity of paclitaxel-induced peripheral neuropathy. *Breast Cancer Research and Treatment* [Internet]. 2020 Mar 12;180(3):707–14. <https://doi.org/10.1007/s10549-020-05584-8>

Cite as: Dosunmu DP, Babatunde OB, Onaolapo OJ. Vitamin D₃ Protects against Cyclophosphamide-induced Neurotoxicity Via the Modulation of Inflammatory Cytokines and Oxidative Stress Parameters *Acta Bioscientia* **2024**;1 (1);013-021 <https://doi.org/10.71181/actabioscientia12130>