

Pyridoxal Phosphate protects against aluminium chloride-induced neurobehavioural and neuromorphological alterations in the rat cerebral cortex

Onigbinde Oluwanisola Akanji^{1, 2}, Ojo Foluso Olamide^{1,3}, Onaolapo Olakunle James⁴, Onaolapo Adejoke Yetunde^{1,5}.

1. Department of Anatomy, Ladoke Akintola University of Technology, Ogbomoso, Oyo State, Nigeria.

2. Department of Anatomy, Nile University of Nigeria, Jabi, Abuja, Nigeria.

3. Department of Anatomy, University of Ilesa, Ilesa, Osun State, Nigeria

4. Behavioural Neuroscience and Neuropharmacology Unit, Department of Pharmacology and Therapeutics, Ladoke Akintola University of Technology, Ogbomoso, Oyo State, Nigeria

5. Behavioural Neuroscience and Neurobiology Unit, Department of Anatomy, Ladoke Akintola University of Technology, Ogbomoso, Oyo State, Nigeria

Correspondence:

Onigbinde, Oluwanisola Akanji

Department of Anatomy,

Ladoke Akintola University of Technology,

Ogbomoso, Oyo State, Nigeria.

Email: onigbindesola@gmail.com. Tel: 2348065801808

ABSTRACT

The increasing prevalence of neurodegenerative diseases has necessitated the search for affordable medicines for better management. Pyridoxal phosphate (PLP) has been reported to improve cognition in humans and rodents; however, its possible benefits in the management of neurodegenerative diseases like Alzheimer's disease has been scarcely examined. This study assessed the effects of PLP on aluminium Chloride-induced neurobehavioural and histomorphological changes in rat cerebral cortex. Forty male Wistar rats weighing 130-150 g each were distributed into four groups (n=10). Group A received distilled water (10 ml/kg), group B received AlCl₃ (50 mg/kg), group C and D received AlCl₃ with PLP at 100 mg/kg and 200 mg/kg respectively. Pyridoxal phosphate was administered by gavage on days 1-15 while AlCl₃ was administered orally on days 16-30. The results revealed a significant increase in horizontal locomotion, rearing and grooming in groups treated with PLP compared to AlCl₃. Spatial working memory increased significantly improved in groups treated with PLP compared to AlCl₃. An increase in MDA and inflammatory cytokines was observed with AlCl₃ compared to control with a reversal of these parameters with PLP supplementation. Results of histomorphological examination revealed evidence of neurodegeneration in the AlCl₃ groups, with PLP supplementation rendering some protection against neuronal injury. In conclusion this study revealed that PLP supplementation was protective against AlCl₃-induced neurobehavioural, oxidative stress, neuroinflammation and neuromorphological injury in rats. However, more studies are required to determine its benefits in humans.

KEYWORDS: Aluminium Chloride; Neurotoxicity; Neuroinflammation; Neuroprotection; Vitamin B6

1. Introduction

Neurodegenerative diseases (NDDs) are progressive disorders that include Ataxias, Alzheimer's, disease, Amyotrophic lateral sclerosis, Huntington's disease, Parkinson's disease, Multiple system atrophy and Progressive supranuclear palsy that are characterised by alteration of the structure and function of the peripheral and central nervous system resulting in nerve cell

death [1]. The global transition in our demographics, which is characterised by an increasing elderly population [2-4] and compounded by the emerging long-term side effects of the COVID-19 pandemic has resulted in a global increase in the prevalence of NDDs [2, 5]. This increased prevalence coupled with the overwhelming burden of NDDs on individuals, families, and communities [3] is necessitating an increased urgency of developing novel therapies that would better manage

the diseases and also address the social and public health burden of NDDs. Generally, the aetiology of NDDs have been attributed to a number of factors including environment, genetics and lifestyle changes [4, 6]. However, individual disorders like Alzheimer's disease also have additional causes including neuroinflammation, oxidative stress, neurotransmitter dysfunction and energy dysmetabolism [1, 5, 7, 8].

Alzheimer's disease (AD), which is the most prevalent form of age-related dementias is characterised by progressive loss of spatial awareness, memory loss and personality changes [7]. The increasing social and economic burden associated with the development of AD has continued to make it a focus of extensive preclinical and clinical research. Recent advances in our understanding of the pathophysiological mechanisms of AD has been aided by the use of animal models which also allow the assessment of novel therapies. These approaches which include the use of transgenic animal models, endogenous substance model, chemical models and heavy metal models of AD [9].

Heavy metal-induced models of AD have been developed based on the premise that exposure to heavy metals including cadmium, lead, manganese, fluoride and aluminium; a number of which are ubiquitous within environment or earth's crust but have been associated with increased risk of AD in the general population [9, 10].

Aluminium (Al) is a heavy metal which is widely distributed within the earth's crust and upon exposure readily crosses the blood brain barrier [9]. However, despite being ubiquitous in the environment, for aeons exposure to it has only been associated with deleterious effects in the body particularly in the brain [11]. Aluminium exposure has been associated with such as increased inflammation, oxidative stress amyloid β deposition, calcium dysmetabolism and increased formation of neurofibrillary tangles in the brain [11]; which are pathological hallmarks of AD. that are associated with the progression of AD [12]. Exposure to Al has also been linked with cholinergic and synaptic dysfunction that is now associated with memory impairment [9]. Several studies have demonstrated that prolonged exposure to Al is associated with accumulation of the heavy metal within the brain particularly the hippocampus and the induction of neuroinflammatory cytokines and oxidative stress [13-18] affirming its validity as a model for studying AD; however, inability to readily reproduce deposition of neurofibrillary tangles and amyloid- β plaques in this model could be a limitation to its use [19]. The use of AD disease models has enabled further understanding of the pathophysiological mechanism of AD. In addition to the previously known pathophysiological mechanisms, there are

reports that have also linked the development of AD to deficits in the intracellular levels of pyridoxal 5'-phosphate [20, 21].

Pyridoxal 5'-phosphate (PLP) is the biochemically active form of vitamin B6 and is often required for biochemical reactions and synthesis of several neurotransmitters including dopamine, epinephrine, norepinephrine, serotonin, and gamma (γ) aminobutyric acid (GABA) [22-25]. In developing economies, the reduction in the levels of PLP is often due to dietary insufficiency, especially in infants, children, and women [26-28]. While several studies have demonstrated the ability of PLP to reverse neurodegenerative changes, oxidative stress and neuroinflammatory changes [29-31], there is however a dearth of scientific information on its effect in aluminium chloride model of AD, hence this study. This study examined the effects of pyridoxal phosphate supplementation on aluminium chloride induced neurobehavioural, and histomorphological changes in the cerebral cortex of rats.

2. Materials and Methodology

2.1 Chemicals and Drugs

Pyridoxal phosphate (Nature's field, USA), Aluminium chloride (Sigma-Aldrich Corporation, USA), normal saline, assay kits for lipid peroxidation (Malondialdehyde (MDA), interleukin-6 (IL-6), interleukin-1 β (IL-1 β), Tumour necrosis factor alpha (TNF- α), and total antioxidant capacity (TAC) (Biovision Inc., Milpitas, CA, USA).

2.2 Animals

Forty male Wistar rats were obtained from the Empire Breeders, Osogbo, Osun State and housed in wooden cages at room temperature (25°C \pm 2.5°C) with 12 hours of light. Animals were acclimatised to the experimental conditions for two weeks.

2.3 Experimental Methods

Forty male Wistar rats weighing 130-150g each were distributed into four groups (n=10). Group A received distilled water (10 ml/kg), group B received AlCl₃ (50 mg/kg), group C and D received AlCl₃ and PLP at 100 mg/kg and 200 mg/kg respectively. Pyridoxal phosphate was administered on days 1-15 while AlCl₃ was administered on days 16-30. All agents were administered orally throughout the experiment. On the 31st day, animals were subjected to neurobehavioural paradigms including the open field box and Y-maze. On the 32nd day, animals were sacrificed by cervical dislocation and blood samples were taken via cardiac puncture for the assay of Total Antioxidant Capacity (TAC), Malondialdehyde (MDA), Tumour Necrosis Factor Alpha (TNF- α), Interleukin-1 β (IL-1 β) and Interleukin-6 (IL-6). The cerebral cortex was excised and processed for histological analysis using haematoxylin and eosin, and cresyl fast violet stains.

2.4 Behavioural Testing

At the end of the experimental period, animals were exposed to behavioural paradigms (open field arena and Y-maze) as previously described [32-35]. Behavioural tests were carried out daily between 7.00 am and 1.00 pm.

2.4.1 Open field Novelty induced Behaviours

Animals were exposed to the open field arena for ten minutes during which the following behaviours were observed and scored; horizontal locomotion, rearing and grooming. The open-field apparatus used was a square box with dimensions of 75 x 75 x 30 cm, featuring a hard floor. The interior of the box was divided into 16 equal squares by permanent red markings. Each rat was placed in the centre of the open field, and its behaviour was recorded as described [36, 37]. The recorded parameters within the 10-minute observation period included: total horizontal locomotion (number of floor units entered with all paws), rearing frequency (number of times the rat stood on its hind legs), and frequency of grooming (number of body-cleaning actions, licking of the body and pubis, and face-washing) which are indicative of stereotypic behaviour.

2.4.2 Memory (Y maze)

Spontaneous alternation is a measure of spatial working-memory. The Y-maze was used to measure short term memory, general locomotor activity and stereotypic behaviours. For each animal, the Y-maze testing was carried out for 5 minutes. The Y-maze consisted of three equally spaced arms, each with dimensions of 41cm in length and 15cm in height. Each arm's floor measured 5 centimeters in width. Observing spontaneous alternation behaviour which is a reflection of rodents' inclination to switch between non-reinforced options was used to gauge spatial working memory. After being put in one of the Y-maze's arm compartments, each rat was free to roam about until its tail fully entered another arm. The arms were labelled A, B, or C, and the arm entry sequence was manually recorded. Entry into each of the three arms in succession was referred to as an alternation. The total number of arms entered minus two was used to determine the number of maximal spontaneous alternations. Next, (actual alternations/maximum alternations) x 100 was used to get the percentage alternation. Between sessions, the equipment was cleaned with 5% alcohol following each test [38].

2.5 Biochemical Test

2.5.1. Interleukin (IL) -1 β and Interleukin-6

Interleukin (IL)-1 β and Interleukin-6 levels were measured using enzyme-linked immunosorbent assay methods with

commercially available kits procured from Biovision Inc., Milpitas, CA, USA.

2.5.2. Antioxidant Status

Total antioxidant capacity was measured using the Trolox Equivalent Antioxidant Capacity Assay that is based on the ability of antioxidants within a sample to react with oxidised products as previously described [15, 16].

2.5.3. Lipid Peroxidation

Lipid peroxidation levels were assessed by determining the malondialdehyde (MDA) content, which assays the levels of thiobarbituric acid reactive substance in samples. Thiobarbituric acid reactive substance combine with free malondialdehyde present in samples to form a coloured complex, the concentration of which is expressed as μmol [15, 16, 37].

2.6 Tissue Histology

Rat brains were dissected, sectioned and fixed in neutral buffered formol calcium. The prefrontal cortex was then processed for paraffin-embedding, cut at 5 μm and stained with haematoxylin/eosin and cresyl fast violet for general histological study.

2.7 Photomicrography:

Histological slides of the prefrontal cortex were examined under an Olympus binocular light microscope. Images were captured using a Canon PowerShot 2500 digital camera. Histopathological alterations were evaluated by a pathologist who was unaware of the group assignments.

2.8 Statistical Analysis

Data was analysed using Chris Rorden's ANOVA for Windows (version 0.98). Analysis of data was by One-way analysis of variance (ANOVA) and a post-hoc test (Tukey HSD) used comparisons. All results were expressed as mean \pm S.E.M. and $p < 0.05$ was taken

3. Results

3.1 Effects of pyridoxal phosphate on relative body and brain weights.

Figure 1 shows the effect of pyridoxal phosphate on changes in bodyweight. There was a significant decrease in body weight in the groups administered aluminium chloride (AlCl_3), AlCl_3 with pyridoxal phosphate (PLP) at 200 mg/kg and an increase with AlCl_3 /PLP at 100 mg/kg compared to control. Compared to AlCl_3 control, relative body weight increased with AlCl_3 /PLP at 100 and 200 mg/kg respectively.




Figure 1: Shows the effect of pyridoxal phosphate on relative change in body weight. Each bar represents Mean \pm S.E.M * $p < 0.05$ vs. control, # $p < 0.05$ significant difference from AlCl₃, number of rats per treatment group = 10. AlCl₃: Aluminium chloride, PLP: Pyridoxal Phosphate.

3.2 Effects of pyridoxal phosphate on relative food intake

Figure 2 shows the effect of pyridoxal phosphate on relative change in food intake. There was a significant decrease in food intake with AlCl₃, AlCl₃PLP at 100 and 200 mg/kg compared to control. Compared to AlCl₃ control, relative food intake increased with AlCl₃/PLP at 100 and 200 mg/kg respectively.




Figure 2: Shows the effect of pyridoxal phosphate on relative change in food intake. Each bar represents Mean \pm S.E.M * $p < 0.05$ vs. control, # $p < 0.05$ significant difference from AlCl₃, number of rats per treatment group = 10. AlCl₃: Aluminium chloride, PLP: Pyridoxal Phosphate

3.3 Effect of pyridoxal phosphate on relative brain weight

Figure 3 shows the effect of pyridoxal phosphate on relative brain weight. There was no significant difference in relative brain weight in any of the groups compared to control or AlCl₃ control.

3.5 Effect of Pyridoxal Phosphate on rearing

Figure 5 shows the effect of pyridoxal phosphate on rearing activity in the open field arena. There was a significant decrease in rearing activity with AlCl₃, AlCl₃PLP at 100 and 200 mg/kg compared to control. Compared to AlCl₃ control, rearing

activity increased with AlCl₃/PLP at 100 and 200 mg/kg respectively.




Figure 3: Shows the effect of pyridoxal phosphate on relative brain weight. Each bar represents Mean \pm S.E.M, number of rats per treatment group = 10. AlCl₃: Aluminium chloride, PLP: Pyridoxal Phosphate.

3.4 Effect of Pyridoxal phosphate on locomotor activity

Figure 4 shows the effect of pyridoxal phosphate on horizontal locomotion in the open field arena. There was a significant decrease in locomotor activity with AlCl₃, AlCl₃PLP at 100 and 200 mg/kg compared to control. Compared to AlCl₃ control, locomotor activity increased with AlCl₃/PLP at 100 and 200 mg/kg respectively.




Figure 4: Shows the effect of pyridoxal phosphate on horizontal locomotion in the open field arena. Each bar represents Mean \pm S.E.M * $p < 0.05$ vs. control, # $p < 0.05$ significant difference from AlCl₃, number of rats per treatment group = 10. AlCl₃: Aluminium chloride, PLP: Pyridoxal Phosphate

3.6 Effect of Pyridoxal Phosphate on self-grooming

Figure 6 shows the effect of pyridoxal phosphate on self-grooming behaviour in the open field arena. There was a significant decrease self-grooming behaviours with AlCl₃, and AlCl₃PLP at 200 mg/kg and a significant increase with AlCl₃PLP at 100 mg/kg compared to control. Compared to AlCl₃ control, self-grooming behaviour increased with AlCl₃/PLP at 100 and decreased with AlCl₃PLP at 200 mg/kg.




Figure 5: Shows the effect of pyridoxal phosphate on rearing activity in the open field arena. Each bar represents Mean \pm S.E.M * $p < 0.05$ vs. control, # $p < 0.05$ significant difference from AlCl₃, number of rats per treatment group =10. AlCl₃: Aluminium chloride, PLP: Pyridoxal Phosphate.




Figure 6: Shows the effect of pyridoxal phosphate on self-grooming behaviour in the open field arena. Each bar represents Mean \pm S.E.M * $p < 0.05$ vs. control, # $p < 0.05$ significant difference from AlCl₃, number of rats per treatment group =10. AlCl₃: Aluminium chloride, PLP: Pyridoxal Phosphate

3.7 Effect of pyridoxal phosphate on working memory

Figure 7 shows the effect of pyridoxal phosphate on spatial working memory in the Y-maze measured as % alternation. There was a significant decrease working memory with AlCl₃, AlCl₃PLP at 100 and 200 mg/kg compared to control. Compared to AlCl₃ control, working memory increased with AlCl₃/PLP at 100 and 200 mg/kg.

3.8 Effect of pyridoxal phosphate on oxidative stress parameters and inflammatory cytokines

Table 1 shows the effects of pyridoxal phosphate shows the effect of pyridoxal phosphate on oxidative stress parameters and inflammatory cytokines. Antioxidant status measured as Total antioxidant capacity (TAC) decreased significantly with AlCl₃ and AlCl₃PLP at 100 mg/kg and increased significantly with AlCl₃PLP at 200 mg/kg compared to control. Compared to AlCl₃ control, TAC levels increased with AlCl₃/PLP at 100 and 200 mg/kg. respectively

Oxidative stress measured as malondialdehyde (MDA) increased significantly with AlCl₃ and AlCl₃PLP at 100 mg/kg and decreased significantly with AlCl₃PLP at 200 mg/kg compared to control. Compared to AlCl₃ control, MDA levels decreased with AlCl₃/PLP at 100 and 200 mg/kg. respectively.




Figure 7: Shows the effect of pyridoxal phosphate on spatial working memory in the Y maze. Each bar represents Mean \pm S.E.M * $p < 0.05$ vs. control, # $p < 0.05$ significant difference from AlCl₃, number of rats per treatment group =10. AlCl₃: Aluminium chloride, PLP: Pyridoxal Phosphate.

Inflammatory cytokines measured were interleukin (IL)-1 β , IL-6 and Tumour necrosis factor (TNF)- α . interleukin (IL)-1 β , IL-6 and TNF - α levels increased significantly with AlCl₃ and AlCl₃PLP at 100 and 200 mg/kg compared to control. Compared to AlCl₃ control, levels of all three inflammatory cytokines decreased with AlCl₃/PLP at 100 and 200 mg/kg. respectively

Table 1: Effect of pyridoxal phosphate on oxidative stress and proinflammatory parameters

Groups	TAC (mM)	MDA (μ M)	IL-6 (pg/ml)	IL-1 β (pg/ml)	TNF- α (pg/ml)
Control	18.92 ± 0.10	5.17 ± 0.10	2.68 ± 1.20	6.94 ± 1.00	19.04 ± 0.34
AlCl ₃	9.81 $\pm 0.12^*$	14.0 $\pm 0.12^*$	15.23 $\pm 1.80^*$	17.99 $\pm 2.20^*$	56.59 $\pm 0.65^*$
AlCl ₃ /PLP 100	15.80 $\pm 0.12^{*#}$	9.72 $\pm 0.14^{*#}$	10.94 $\pm 2.15^{*#}$	17.97 $\pm 2.20^*$	30.56 $\pm 0.15^{*#}$
AlCl ₃ /PLP 200	24.65 $\pm 0.15^{*#}$	3.25 $\pm 0.30^{*#}$	7.64 $\pm 2.60^{*#}$	10.47 $\pm 2.20^{*#}$	26.5 $\pm 0.30^{*#}$

Mean \pm S.E.M, * $p < 0.05$ significant difference from control. # $p < 0.05$ significant difference from AlCl₃. Number of rats per treatment group =10. AlCl₃: Aluminium chloride, PLP: Pyridoxal phosphate, TNF- α : Tumor necrosis factor α . IL: Interleukin.

3.9 Effect of pyridoxal phosphate on the cerebral cortex histomorphology

Figure 8 (A-D) and 9 (A-D) represent the photomicrographs of haematoxylin and eosin (H&E) and cresyl fast violet (CFV) stained sections of the rat cerebral cortex respectively. Examination of the H&E-stained slides of rats in the control group revealed characteristic features of the inner pyramidal layer of the cortex. The CFV stain micrographs also revealed well preserved multipolar shaped pyramidal cells within the neuropil. Granular neurons with large open-faced nuclei, scanty cytoplasm and small sized neuroglial cells were also observed.

In the group administered AlCl_3 , numerous degenerating pyramidal and granule cells and giant cells with double nuclei were observed features which are in keeping with neuronal degeneration and neuronal injury. Also observed were pale-staining neurons with shrunken nuclei. which were interspersed between a few normal pyramidal and granule cells. In the groups administered AlCl_3 /PLP 100 mg/kg (8C & 9 C) and AlCl_3 /PLP 200 mg/kg (8 D and 9 D), reversal of neuronal injury and preservation of Nissl bodies were observed compared with the AlCl_3 group.




Figure 8: Representative photomicrograph of haematoxylin and Eosin-stained sections of the rat cerebral cortex. Photomicrographs revealed distinct layers of the cerebral cortex with presence of numerous granule cells (Gc), pyramidal cells (PYR), and neuroglial cells (NG). Also observed are degenerating pyramidal cell (dPYR), degenerating granule cell (dGc) and Giant cell G. Magnification: x400.

4. Discussion

This study investigated the possible protective effects of pyridoxal phosphate on aluminium chloride-induced neurobehavioural, biochemical and histomorphological changes in rats. The results revealed that pyridoxal phosphate protected against neurotoxic effects associated with aluminium chloride ingestion supporting other literature that had reported similar findings following aluminium chloride ingestion in rodents [15, 16, 39]- Significant changes were observed in food consumption, relative body weight, neurobehavioural

parameters, antioxidant status, oxidative stress and proinflammatory markers. In this study, animal exposure to AlCl_3 revealed a significant reduction in feed intake, body weight, antioxidant capacity and behavioural parameters. There was also an increase in the levels of MDA, IL-1 β , IL-6 and TNF- α in the AlCl_3 group compared to the control. Photomicrographs of the AlCl_3 group further revealed neuronal degeneration in the histopathological architecture of the brain.




Figure 9: Photomicrograph of cresyl fast violet stained sections of the rat cerebral cortex section by cresyl fast violet X100. The photomicrographs show the neuronal cell scattered within the Nissl substance in groups A (Control), C (AlCl_3 and pyridoxal Phosphate at 100 mg/kg) and D (AlCl_3 and pyridoxal Phosphate at 200 mg/kg) and neurodegenerative cells in group B (Aluminium-chloride control). Magnification x 400.

The administration of PLP however, protected against the development of the deleterious effects of AlCl_3 across different parameters including feed intake, body weight, neurobehaviour, oxidative stress marker and proinflammatory cytokines. The photomicrographs of the treated group further revealed a reversal of the damage to the pyramidal and granular neurons as observed in the AlCl_3 group. The reduction seen in feed intake and body weight following the administration of AlCl_3 corroborates reports of previous studies that had reported significant reduction in feed intake and body weight following exposure of animals to AlCl_3 [15, 16]. Moreover, the significant increase in weight of animals treated with AlCl_3 /PLP100 mg/kg and AlCl_3 /PLP200mg/kg, corroborates reports of the research findings by Olofinnade *et al* [29] that examined the effects of pyridoxal phosphate in chlorpromazine-induced parkinsonism rodent model. They reported that PLP administration at 100 mg/kg was associated with weight gain [29] similar to our observations in this study.

Furthermore, blood assays revealed a significant increase in the total antioxidant capacity (TAC) levels with pyridoxal phosphate treated group compared with AlCl_3 group while MDA, IL-6 and TNF- α levels decreased significantly with the treated group compared with AlCl_3 group. There was also a significant decrease in IL-1 β level in the group treated with AlCl_3 /PLP 200 compared with groups AlCl_3 and AlCl_3 /PLP100. Exposure to AlCl_3 have been reported by previous studies to

significantly elevate brain concentrations of some key pro-inflammatory cytokines like TNF- α , IL-1 β and IL-6 [15,16]. The upregulation of these cytokines stimulates leukocyte production which will further intensify the inflammatory response leading to more production of additional pro-inflammatory mediators to the site of call. According to Abulfadl *et al.* [40] accumulation of AlCl₃ triggers the release of cytochrome C from mitochondria, leading to an increased free radical production and elevated pro-inflammatory cytokine levels [41]. In this study PLP's ability to increase antioxidant status was observed which was evidenced by a reduction in MDA and an increase in TAC levels in groups that received PLP, corroborating the evidence from previous studies that demonstrated PLP's significant antioxidant effects [42-45]. These neuroprotective effects of PLP have also been linked to its ability to inhibit the formation of malondialdehyde and scavenge reactive oxygen species [44, 45]. In addition, PLP's ability to reduce lipid peroxidation, increase antioxidant levels, and decrease inflammation even at the molecular level are possible mechanisms through which it reversed the damaging effects of AlCl₃ [14].

Exploration of the neurobehavioural paradigms revealed a significant increase in the open field exploration activities in the treated group compared with AlCl₃ group. Previous studies had reported that animal exposed to AlCl₃ exhibited behaviours like impaired locomotion, self-grooming, and anxiety-like responses. This is as a result of AlCl₃'s interference with cholinergic and dopaminergic systems that play vital roles in locomotion, memory and self-awareness. Dopaminergic and cholinergic dysfunction are key to motor and self-maintenance deficits [46]. The increase observed in the open field exploration activities in the PLP treatment groups compared to AlCl₃ group suggests that PLP has a modulatory effect on locomotion, rearing and grooming. According to Olofinnade *et al* [29], PLP modulated behaviours possibly through its positive effects on brain neurotransmitters activity or receptor binding. Besides, the improved performance observed in the exploration activities in the Y-maze apparatus of the AlCl₃/PLP groups compared to AlCl₃ group also supported the work done by Jeanclos *et al* [47] and Olofinnade *et al* [29], that reported an increase in the levels of pyridoxal phosphate was associated with increased cognition.

Lastly, aluminium chloride administration had been reported to be associated with morphological evidence of neuronal injury due to its neurotoxic effects [15, 16]. Histopathological analysis using H&E and cresyl fast violet stains further corroborated these findings. In this study, the photomicrographs of animals exposed to AlCl₃ revealed notable cortical degeneration, loss of neuronal structure and depletion of Nissl body substances. This indicates possible disruption in protein synthesis and neuronal

function. However, animals treated with pyridoxal phosphate revealed a preserved brain architecture, nearly restoring normal pyramidal cell morphology and Nissl body presence.

Conclusion

In conclusion, this study demonstrated the possible protective effects of pyridoxal phosphate supplementation on aluminium chloride induced changes in behaviour, oxidative stress, proinflammation and cerebral cortex histomorphology in rats. However, more studies are required to determine its benefits in humans.

Funding.

None

Availability of data and materials

Data generated during and analysed during the course of this study are available from the corresponding author on request.

Declarations Ethics approval

Ethical approval for this study was granted by the Ethical Committee of the Faculty of Basic Medical Sciences.

Competing interests

All authors of this paper declare that there is no conflict of interest related to the content of this manuscript.

Received 4th February, 2025, Revised 15th February 2025, Accepted 19th February 2025,

Published online:25/02/25

References

1. Zaib S, Javed H, Khan I, Jaber F, Sohail A, Zaib Z, Mehboob T, Tabassam N, Ogaly HA, *Chemistry Select* 2023, 8, e202300225.
2. Wang S, Jiang Y, Yang A, Meng F, Zhang J The Expanding Burden of Neurodegenerative Diseases: An Unmet Medical and Social Need. *Aging and disease*. 2025;16: 5
3. GBD 2021 Nervous System Disorders Collaborators. Global, regional, and national burden of disorders affecting the nervous system, 1990-2021: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet Neurol*. 2024;23(4):344-381. doi: 10.1016/S1474-4422(24)00038-3.
4. Onaolapo AY, Ojo FO, Adeleye OO, Falade J, Onaolapo OJ. Diabetes Mellitus and Energy Dysmetabolism in Alzheimer's Disease: Understanding the Relationships and Potential Therapeutic Targets. *Curr Diabetes Res*. 2023; 19. (8): e020123212333..
5. Onaolapo A, Onaolapo O. COVID-19, the Brain, and the Future: Is Infection by the Novel Coronavirus a Harbinger of Neurodegeneration? *CNS Neurol Disord Drug Targets*. 2022;21(9):818-829. doi: 10.2174/1871527321666211222162811.
6. Relja M. Pathophysiology and Classification of Neurodegenerative Diseases. *EJIFCC*. 2004;15(3):97-99.
7. Onaolapo OJ, Olofinnade AT, Ojo FO, Onaolapo AY. Neuroinflammation and Oxidative Stress in Alzheimer's Disease; Can Nutraceuticals and Functional Foods Come to the

- Rescue? Antiinflamm Antiallergy Agents Med Chem. 2022;21(2):75-89. doi: 10.2174/1871523021666220815151559.
8. Onaolapo AY, Onaolapo OJ. Peripheral and Central Glutamate Dyshomeostasis in Neurodegenerative Disorders. *Curr Neuropsychopharmacol*. 2021;19(7):1069-1089. doi: 10.2174/1570159X18666201015161919.
 9. Dhapola, R., Kumari, S., Sharma, P. *et al*. Insight into the emerging and common experimental in-vivo models of Alzheimer's disease. *Lab Anim Res* **39**, 33 (2023). <https://doi.org/10.1186/s42826-023-00184-1>
 10. Bakulski KM, Seo YA, Hickman RC, Brandt D, Vadari HS, Hu H, et al. Heavy metals exposure and Alzheimer's disease and related dementias. *J Alzheimers Dis*. 2020;76:1215.
 11. Dey M, Singh RK. Neurotoxic effects of aluminium exposure as a potential risk factor for Alzheimer's disease. *Pharmacol Rep*. 2022 Jun;74(3):439-450. doi: 10.1007/s43440-022-00353-4.
 12. Abd El-Aziz NM, Shehata MG, Alsulami T, Badr AN, Elbakatoshy MR, Ali HS, El-Sohaimy SA. Characterization of Orange Peel Extract and Its Potential Protective Effect against Aluminum Chloride-Induced Alzheimer's Disease. *Pharmaceuticals (Basel)*. 2022 Dec 22;16(1):12. doi: 10.3390/ph16010012.
 13. Liaquat L, Sadir S, Batool Z, Tabassum S, Shahzad S, Afzal A, Haider S. Acute aluminum chloride toxicity revisited: Study on DNA damage and histopathological, biochemical and neurochemical alterations in rat brain. *Life sciences*. 2019 Jan 15;217:202-11
 14. Elhadidy ME, Sawie HG, Meguid NA, Khadrawy YA. Protective effect of ashwagandha (*Withania somnifera*) against neurotoxicity induced by aluminum chloride in rats. *Asian Pacific Journal of Tropical Biomedicine*. 2018 Jan 1;8(1):59-66.
 15. Akinsehinwa AF, Samson PO, Onaolapo OJ, Onaolapo AY. Quercetin/Donepezil co-administration mitigates Aluminium chloride induced changes in open field novelty induced behaviours and cerebral cortex histomorphology in rats. *Acta Bioscientia* **2025**;1(1):01-07 doi.org/10.71181/actabioscientia12110
 16. Ajao JA, Akinsehinwa AS, Onaolapo OJ, Onaolapo AY. Alcohol Extract of Muira Puama (*Ptychopetalum Olacoides*) ameliorates Aluminium Chloride-induced changes in Behaviour and Cerebral cortex Histomorphology in Wistar Rats. *Acta Bioscientia* **2025**;1(1):022-029 doi.org/10.71181/actabioscientia12140.
 17. Cao Z, Wang F, Xiu C, Zhang J, Li Y. Hypericum perforatum extract attenuates behavioral, biochemical, and neurochemical abnormalities in aluminum chloride-induced Alzheimer's disease rats. *Biomed Pharmacother*. 2017;91:931-7.
 18. Justin Thenmozhi A, Dhivyabharathi M, William Raja TR, Manivasagam T, Essa MM. Tannoid principles of *Embllica officinalis* renovate cognitive deficits and attenuate amyloid pathologies against aluminum chloride induced rat model of Alzheimer's disease. *Nutr Neurosci*. 2016;19:269-78.
 19. Xu Q-Q, Yang W, Zhong M, Lin ZX, Gray NE, Xian YF. Animal models of Alzheimer's disease: preclinical insights and challenges. *Acta Mater Medica*. 2023;2:192-215.
 20. Wilson MP, Plecko B, Mills PB, Clayton PT. Disorders affecting vitamin B6 metabolism. *Journal of inherited metabolic disease*. 2019 Jul;42(4):629-46.
 21. Ghatge MS, Al Mughran M, Omar AM, Safo MK. Inborn errors in the vitamin B6 salvage enzymes associated with neonatal epileptic encephalopathy and other pathologies. *Biochimie*. 2021 Apr 1;183:18-29.
 22. Barile A. Characterisation of the key enzymes involved in vitamin B6 Salvage Pathway in *Escherichia coli* and humans. Doctoral theis. Sapienza universita di Roma (2019). <https://hdl.handle.net/11573/1379435>.
 23. Al-Shekaili HH, van Karnebeek C, Leavitt BR. Vitamin B6 and related inborn errors of metabolism. In: *B-Complex Vitamins- Sources, Intakes and Novel Applications 2021 Sep 3*. IntechOpen.
 24. Wu L, Liu L, Jin Z, Zhang D. Advances and prospects in the biosynthesis of vitamin B6 and its bioengineering as a cofactor of other chemicals. *Food Bioengineering*. 2022 Jun;1(2):171-81.
 25. Onaolapo OJ, Onaolapo AY. Chapter 31 - B vitamins: Pyridoxal phosphate and parkinsonism, Editor(s): Colin R. Martin, Vinood B. Patel, Victor R. Preedy, *Vitamins and Minerals in Neurological Disorders*, Academic Press, 2023, Pages 527-542, ISBN 9780323898355, <https://doi.org/10.1016/B978-0-323-89835-5.00029-6>.
 26. Isabirye N, Ezeamama AE, Kyeyune-Bakyayita R, Bagenda D, Fawzi WW, Guwatudde D. Dietary micronutrients and gender, body mass index and viral suppression among HIV-infected patients in Kampala, Uganda. *International Journal of Maternal and Child Health and AIDS*. 2020;9(3):337.
 27. Ali MA, Hafez HA, Kamel MA, Ghamry HI, Shukry M, Farag MA. Dietary vitamin B complex: orchestration in human nutrition throughout life with sex differences. *Nutrients*. 2022 Sep 22;14(19):3940.
 28. Kiani AK, Dhuli K, Donato K, Aquilanti B, Velluti V, Matera G, Iaconelli A, Connelly ST, Bellinato F, Gisondi P, Bertelli M. Main nutritional deficiencies. *Journal of preventive medicine and hygiene*. 2022 Jun;63(2 Suppl 3):E93.
 29. Olofinnade AT, Onaolapo TM, Oladimeji S, Fatoki AM, Balogun CI, Onaolapo AY, Onaolapo OJ. An Evaluation of the Effects of Pyridoxal Phosphate in Chlorpromazine-induced Parkinsonism using Mice. *Cent Nerv Syst Agents Med Chem*. 2020;20(1):13-25. doi: 10.2174/1871524920666200120142508.
 30. Hashim A, Wang L, Juneja K, Ye Y, Zhao Y, Ming LJ. Vitamin B6s inhibit oxidative stress caused by Alzheimer's disease-related Cu(II)- β -amyloid complexes-cooperative action of phospho-moiety. *Bioorg Med Chem Lett*. 2011 Nov 1;21(21):6430-2. doi: 10.1016/j.bmcl.2011.08.123.
 31. Ko JW, Jeon S, Kwon YH. Dietary vitamin B6 restriction aggravates neurodegeneration in mice fed a high-fat diet. *Life Sci*. 2022 Nov 15;309:121041. doi: 10.1016/j.lfs.2022.121041.
 32. Onaolapo OJ, Adeyemi OI, Amujoyegbe OJ, Fasola EA, Olofinnade AT, Onaolapo AY. High dietary fat modulates neurobehavioural effect of lopinavir/ritonavir in mice. *Current Pharmaceutical Biotechnology*. 2020;21(2):158-68.
 33. Onaolapo, O.J., Olofinnade, A.T., Ojo, F.O., Falade, J. and Onaolapo, A.Y., 2023. Prepubertal continuous dietary folate fortification enhances the brain function of adult mice by modulating antioxidant status, inflammation, and brain neurotransmitter levels. *Anti-Inflammatory & Anti-Allergy Agents in Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Inflammatory and Anti-Allergy Agents)*, 22(3), pp.198-209.
 34. Olofinnade AT, Alawode A, Onaolapo AY, Onaolapo OJ. *Lepidium meyenii* supplemented diet modulates neurobehavioural and biochemical parameters in mice fed high-fat high-sugar diet. *Endocrine, Metabolic & Immune Disorders-Drug Targets (Formerly Current Drug Targets-Immune, Endocrine & Metabolic Disorders)*. 2021 Jul 1;21(7):1333-43.
 35. Saeed NM, Esmail MM, Abdel-Rahman H, Elgindy AM. Valsartan ameliorated cognitive decline, oxidative stress and inflammation in A β 1-3-induced Alzheimer's disease in rats. *ERU Research Journal*. 2024 Apr 1;3(2):1038-57.

36. Knight P, Chellian R, Wilson R, Behnood-Rod A, Panunzio S, Bruijnzeel AW. Sex differences in the elevated plus-maze test and large open field test in adult Wistar rats. *Pharmacology Biochemistry and Behavior*. 2021 May 1;204:173168.
37. Olofinnade AT, Onaolapo AY, Onaolapo OJ. Graded effects of dry feed added sodium benzoate/ascorbic acid combination on neurobehaviour, oxidative stress, and markers of inflammation in mice. *Curr Bioact Compd* 2024b; 20(1): e060723218474. <http://dx.doi.org/10.2174/1573407219666230706145> 15.
38. Onaolapo AY, Sulaiman H, Olofinnade AT, Onaolapo OJ. Antidepressant-like potential of silymarin and silymarin/sertraline combination in mice: Highlighting effects on behaviour, oxidative stress, and neuroinflammation. *World J Pharmacol* 2022; 11(3): 27-47. [DOI: 10.5497/wjp.v11.i3.27].
39. Shabbir A, Rehman K, Akash MS, Akbar M, Chaudhary Z, Panichayupakaranant P, Shah MA. Differential neuroprotective effect of curcuminoid formulations in aluminum chloride-induced Alzheimer's disease. *Environmental Science and Pollution Research*. 2022 Sep;29(45):67981-96.
40. Abulfadl YS, El-Maraghy NN, Ahmed AE, Nofal S, Abdel-Mottaleb Y, Badary OA. Thymoquinone alleviates the experimentally induced Alzheimer's disease inflammation by modulation of TLRs signaling. *Hum Exp Toxicol*. 2018;37(10):1092–104.
41. Milnerowicz H, Ściskalska M, Dul M. Pro-inflammatory effects of metals in persons and animals exposed to tobacco smoke. *J Trace Elem Med Biol*. 2015;29:1–10.
42. Choi JM, Park HS, He MT, Kim YS, Kim HY, Lee AY, Cho EJ. Membrane-Free Stem Cells and Pyridoxal 5'-Phosphate Synergistically Enhance Cognitive Function in Alzheimer's Disease Mouse Model. *Antioxidants*. 2022;11(3):601.
43. Almuhayawi MS, Ramadan WS, Harakeh S, Al Jaouni SK, Bharali DJ, Mousa SA, Almuhayawi SM. The potential role of pomegranate and its nano-formulations on cerebral neurons in aluminum chloride induced Alzheimer rat model. *Saudi journal of biological sciences*. 2020 ;27(7):1710-6.
44. Taş S, Sarandöl E, Dirican M. Vitamin B6 supplementation improves oxidative stress and enhances serum paraoxonase/arylesterase activities in streptozotocin-induced diabetic rats. *The Scientific World Journal*. 2014;2014(1):351598.
45. Hsu CC, Cheng CH, Hsu CL, Lee WJ, Huang SC, Huang YC. Role of vitamin B6 status on antioxidant defenses, glutathione, and related enzyme activities in mice with homocysteine-induced oxidative stress. *Food & Nutrition Research*. 2015 Jan 1;59(1):25702.
46. Choi HJ, Park JH, Jeong YJ, Hwang JW, Lee S, Lee H, et al. Donepezil ameliorates Aβ pathology but not tau pathology in 5xFAD mice. *Mol Brain*. 2022;15(1):63.
47. Jeanclos E, Albersen M, Ramos RJ, Raab A, Wilhelm C, Hommers L, Lesch KP, Verhoeven-Duif NM, Gohla A. Improved cognition, mild anxiety-like behavior and decreased motor performance in pyridoxal phosphatase-deficient mice. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*. 2019 Jan 1;1865(1):193-205.

Cite as: Onigbinde OA, Ojo FO, Onaolapo OJ, Onaolapo AY. Pyridoxal Phosphate protects against aluminium chloride-induced neurobehavioural and neuromorphological alterations in the rat cerebral cortex *Acta Bioscientia* **2025**;1(2);95-103 <https://doi.org/10.71181/actabioscientia12240>