

Testing of Statistical Hypothesis

Adejumo Ridwan Suleiman

2022-06-30

What is a Statistical Hypothesis

- ▶ This is a formal statement about the nature of a population.
- ▶ Example;
 - ▶ Are all Africans rich?
 - ▶ Do all footballers make above \$100k/week.
 - ▶ Do Ronaldo and Messi have the same goal ratio.

Hypothesis Testing

- ▶ This is the use of statistical tests to make an inference about a population.
- ▶ The method of inference used depends on the nature of the data and reason for analysis.

Steps in conducting a Statistical Hypothesis Test

- ▶ Stating the Null and Alternative Hypothesis
- ▶ Stating the Level of Significance
- ▶ Test Statistics to be used
- ▶ Critical Region
- ▶ Decision Rule

Null Hypothesis

- ▶ Known as the Hypothesis of no difference.
- ▶ This Hypothesis states that the relationship the researcher is trying to investigate does not exist.
- ▶ Our hope is always to prove the **Null hypothesis** wrong.
- ▶ The **Null Hypothesis** is represented mathematically as H_0

Alternative Hypothesis

- ▶ Opposite of the **Null Hypothesis**
- ▶ This is usually the hypothesis of the researcher.
- ▶ Represented as H_A

Case Study

- ▶ Let's say the mean price of diamonds is less than \$10000
- ▶ In this case the **Null Hypothesis** can be stated as;
"The mean price of diamonds is \$10000"
- ▶ The **Alternative hypothesis** can be stated as ;
"The mean price of diamonds is less than \$1000"
- ▶ The parameter of interest in this example is the **Mean**.

Type 1 and Type 2 error

- ▶ Situation when you reject the **Null Hypothesis** when it is actually true is known as **Type 1** error.
- ▶ Situation when you fail to reject the **Alternative Hypothesis** when it is actually false is known as **Type 2** error.

Level of Significance (α)

- ▶ Situation where an experiment occurred beyond chance, then it is said to **Statistically Significant**.
- ▶ If we are to perform an experiment a number of times and the result of that experiment does not occurs by chance the experiment is declared **Statistically Significant**
- ▶ The level of significance denoted by α is the probability of rejecting the null hypothesis when it is actually true.
- ▶ The level of significance is mostly used as **0.05**.
- ▶ We can say that this is the probability of committing a **Type 1** error.

p-Value

- ▶ This is the probability of having the result of an experiment by chance when reproduced many times.
- ▶ For example a p-value of 0.32 means that when the experiment is repeated a number of times, there is a 32% chance that the result of your experiment actually occurred by chance.
- ▶ The lower the p-value the higher the statistical significance of the test.

Correlation Test

- ▶ This statistics test if two continuous variables are related to each other.
- ▶ The value ranges from -1 to +1 and the higher the value the stronger the relationship.
- ▶ Let's say from the diamonds data example we want to see if there is a relationship between diamonds carat and price.

```
##
```

```
## Pearson's product-moment correlation
```

```
##
```

```
## data: diamonds$price and diamonds$carat
```

```
## t = 551.41, df = 53938, p-value < 2.2e-16
```

```
## alternative hypothesis: true correlation is not equal to 0
```

```
## 95 percent confidence interval:
```

```
## 0.9203098 0.9228530
```

```
## sample estimates:
```

```
## cor
```

```
## 0.9215913
```

t-test