

Oncology Clinical Pathways

Head and Neck Cancer

January 2023 – V1.2023



Choose **VA**



SHOULDER to SHOULDER
Every Step of the Way

VA



U.S. Department
of Veterans Affairs

Table of Contents

Head and Neck Cancer Presumptive Conditions	3
Oral Cavity T1-2, N0, M0	4
Oral Cavity T1-4, N1-3, M0 and T3-4, N0, M0	5
Oral Cavity Recurrence	6
Oropharynx HPV Positive or Negative T1-2, N0, M0	7
Oropharynx HPV Positive or Negative T1-2, N1	8
Oropharynx HPV Positive or Negative T1-2, N2-3 and T3-4, N0-3	9
Hypopharynx T1-2, N0, M0	10
Hypopharynx T1-4, N1-3 and T3-4, N0, M0	11
Larynx Supraglottis T1-2, N0, M0	12
Larynx Supraglottis T1-4, N1-3 and T3-4, N0, M0	13
Larynx Glottis T1-2, N0, M0	14
Larynx Glottis T1-4, N1-3 and T3-4, N0, M0	15
Larynx Subglottis T1-2, N0, M0	16
Larynx Subglottis T1-4, N1-3 and T3-4, N0, M0	17
Nasopharynx T1, N0, M0	18
Nasopharynx T0-1, N1, M0; T2, N0, M0; and T2, N1, M0	19
Nasopharynx T3-4, N0-3, M0	20
Local or Regional Recurrent Disease	21
Metastatic or Recurrent Disease	22
Unknown Primary	23

Head and Neck Cancer – Presumptive Conditions

VA automatically presumes that certain disabilities were caused by military service. This is because of the unique circumstances of a specific Veteran's military service. If a presumed condition is diagnosed in a Veteran within a certain group, they can be awarded disability compensation.

Gulf War and Post 9/11 Veterans


If the patient served on or after Sept. 11, 2001, in Afghanistan, Djibouti, Egypt, Jordan, Lebanon, Syria, Uzbekistan, or Yemen or if the patient served in the *Southwest Asia theater of operations, or Somalia, on or after Aug. 2, 1990, specific conditions include:

- Head cancer of any type
- Neck cancer of any type

* The Southwest Asia theater of operations refers to Iraq, Kuwait, Saudi Arabia, the neutral zone between Iraq and Saudi Arabia, Bahrain, Qatar, the United Arab Emirates, Oman, the Gulf of Aden, the Gulf of Oman, the Persian Gulf, the Arabian Sea, the Red Sea, and the airspace above these locations.

For more information, please visit [U.S. Department of Veterans Affairs - Presumptive Disability Benefits \(va.gov\)](https://www.va.gov)

Head and Neck Cancer – Oral Cavity T1-2, N0, M0



Clinical trial(s) always considered on pathway.


^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **Resection** consider neck dissection if depth of invasion is >3mm; always consider neck dissection with perineural or perivascular/vascular invasion

^c **Adverse features include** extranodal extension, positive margins, close margins, pT3 or pT4 primary, pN2 or pN3 nodal disease, nodal disease in levels IV or V, perineural invasion, perivascular/vascular invasion, or lymphatic invasion

^d **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

Head and Neck Cancer – Oral Cavity T1-4, N1-3, M0 and T3-4, N0, M0




Clinical trial(s) always considered on pathway.

^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

^c **Radiation** forgo radiation for N1 disease with a single lymph node if the only adverse pathological feature

Head and Neck Cancer – Oral Cavity Recurrence



^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **Plan for total cisplatin** dose $\geq 200 \text{ mg/m}^2$ (either as 100 mg/m^2 IV every 3 weeks for 3 cycles or 40 mg/m^2 IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel



Choose **VA**




SHOULDER to SHOULDER
Every Step of the Way

VA



U.S. Department
of Veterans Affairs

Head and Neck Cancer – Oropharynx HPV Positive or Negative T1-2, N0, M0




Clinical trial(s) always considered on pathway.

^a **Pathway takes into consideration** the difference in staging between HPV negative and positive disease

^b **Adverse features include** extranodal extension, positive margins, close margins, pT3 or pT4 primary, pN2 or pN3 nodal disease, nodal disease in levels IV or V, perineural invasion, perivascular/vascular invasion, or lymphatic invasion

^c **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

Head and Neck Cancer – Oropharynx HPV Positive or Negative T1-2, N1



Clinical trial(s) always considered on pathway.


^a **Pathway takes into consideration** the difference in staging between HPV negative and positive disease

^b **Adverse features include** extranodal extension, positive margins, close margins, pT3 or pT4 primary, pN2 or pN3 nodal disease, nodal disease in levels IV or V, perineural invasion, perivascular/vascular invasion, or lymphatic invasion

^c **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); vandidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

^d **If not platinum eligible**, prescribe cetuximab

Head and Neck Cancer – Oropharynx HPV Positive or Negative T1-2, N2-3 and T3-4, N0-3




Clinical trial(s) always considered on pathway.

^a **Pathway takes into consideration** the difference in staging between HPV negative and positive disease

^b **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

Head and Neck Cancer – Hypopharynx T1-2, N0, M0



Clinical trial(s) always considered on pathway.

^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **Adverse features include** extranodal extension, positive margins, close margins, pT3 or pT4 primary, pN2 or pN3 nodal disease, nodal disease in levels IV or V, perineural invasion, perivascular/vascular invasion, or lymphatic invasion

^c **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel.



Choose **VA**




SHOULDER to SHOULDER
Every Step of the Way

VA



U.S. Department
of Veterans Affairs

Head and Neck Cancer – Hypopharynx T1-4, N1-3 and T3-4, N0, M0



Clinical trial(s) always considered on pathway.


^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **Bulky nodal disease** includes N2, N3

^c **Adverse features include** extranodal extension, positive margins, close margins, pT3 or pT4 primary, pN2 or pN3 nodal disease, nodal disease in levels IV or V, perineural invasion, perivascular/vascular invasion, or lymphatic invasion

^d **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

Head and Neck Cancer – Larynx Supraglottis T1-2, N0, M0




Clinical trial(s) always considered on pathway.

^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **Adverse features include** extranodal extension, positive margins, close margins, pT3 or pT4 primary, pN2 or pN3 nodal disease, nodal disease in levels IV or V, perineural invasion, perivascular/vascular invasion, or lymphatic invasion

^c **Plan for total cisplatin** dose $\geq 200 \text{ mg/m}^2$ (either as 100 mg/m^2 IV every 3 weeks for 3 cycles or 40 mg/m^2 IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

Head and Neck Cancer – Larynx Supraglottis T1-4, N1-3 and T3-4, N0, M0




Clinical trial(s) always considered on pathway.

^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **Adverse features include** extranodal extension, positive margins, close margins, pT3 or pT4 primary, pN2 or pN3 nodal disease, nodal disease in levels IV or V, perineural invasion, perivascular/vascular invasion, or lymphatic invasion

^c **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

Head and Neck Cancer – Larynx Glottis T1-2, N0, M0




Clinical trial(s) always considered on pathway.

^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **Consider** voice quality, swallowing function, and ability to adhere to radiation protocols

^c **Surgical options include** cold steel versus laser

Head and Neck Cancer – Larynx Glottis T1-4, N1-3 and T3-4, N0, M0



Clinical trial(s) always considered on pathway.


^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **If patient T4** with obvious cartilage invasion, laryngectomy with nodal basis dissection is preferred

^c **Adverse features include** extranodal extension, positive margins, close margins, pT3 or pT4 primary, pN2 or pN3 nodal disease, nodal disease in levels IV or V, perineural invasion, perivascular/vascular invasion, or lymphatic invasion

^d **Plan for total cisplatin** dose $\geq 200 \text{ mg/m}^2$ (either as 100 mg/m^2 IV every 3 weeks for 3 cycles or 40 mg/m^2 IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

Head and Neck Cancer – Larynx Subglottis T1-2, N0, M0




Clinical trial(s) always considered on pathway.

^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **Patient preference and positive margins:** consider voice quality, swallowing function, ability to adhere to radiation protocols, and patient preference

^c **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

Head and Neck Cancer – Larynx Subglottis T1-4, N1-3 and T3-4, N0, M0



Clinical trial(s) always considered on pathway.

^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing

^b **Consider consultation** with thoracic surgery

^c **Adverse features include** extranodal extension, positive margins, close margins, pT3 or pT4 primary, pN2 or pN3 nodal disease, nodal disease in levels IV or V, perineural invasion, perivascular/vascular invasion, or lymphatic invasion

^d **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel


Head and Neck Cancer – Nasopharynx T1, N0, M0



Clinical trial(s) always considered on pathway.

^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing


Head and Neck Cancer – Nasopharynx T0-1, N1, M0; T2, N0, M0; and T2, N1, M0



Clinical trial(s) always considered on pathway.

- ^a **Evaluate need for support regarding** tobacco use/smoking cessation, dental health, nutrition, audiology, speech/swallowing
- ^b **High risk features** include bulky disease and elevated EBV titers
- ^c **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

Head and Neck Cancer – Nasopharynx T3-4, N0-3, M0



Clinical trial(s) always considered on pathway.

^a **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel


EBV Epstein-Barr Virus

Head and Neck Cancer – Local or Regional Recurrent Disease



Clinical trial(s) always considered on pathway.

Head and Neck Cancer – Metastatic or Recurrent Disease



Clinical trial(s) always considered on pathway.

^a **Patient not eligible** for localized therapies

^b **Palliative Care**, consider palliative radiation


^c **Candidacy** based on platinum toxicities such as adequate cell counts, severe neuropathy, hearing loss/tinnitus, renal failure toxicity, and/or need for rapid cytoreduction

^d **Pembrolizumab**, duration maximum of two years

^e **If not docetaxel eligible**, prescribe cetuximab

CPS Combined Positive Score

Head and Neck Cancer – Unknown Primary



Clinical trial(s) always considered on pathway.

^a **Directed biopsy** of bilateral base of tongue, lingual tonsils, palatine tonsils; if Level 5 node positive, include nasopharynx

^b **Tonsillectomy** should be considered unless patient has very small, soft palatine tonsils

^c **Appropriate Pathway** if HPV positive follow oropharynx pathway; if EBV positive follow the nasopharynx pathway

^d **Plan for total cisplatin** dose ≥ 200 mg/m² (either as 100 mg/m² IV every 3 weeks for 3 cycles or 40 mg/m² IV weekly with concurrent radiation); candidates for high-dose cisplatin should be ECOG PS 0-1 with minimal comorbidities, as toxicities may be anticipated (i.e. nephrotoxicity, neutropenia, nausea/vomiting, etc.); **if not cisplatin-eligible**, prescribe carboplatin-fluorouracil or carboplatin-paclitaxel

Questions?

Contact VHAOncologyPathways@va.gov



Choose **VA**



SHOULDER to SHOULDER
Every Step of the Way

VA



U.S. Department
of Veterans Affairs