

Math 340: Elementary Matrix and Linear Algebra

Final Exam

Monday, May 5th, 7:25pm-9:25pm

Circle your Instructor and your TA:

Dr. Lars Niedorf	Dr. Jose Rodriguez	Dr. Yassine Tissaoui	Dr. Ruhui Jin	Dr. Timur Yastrzhembskiy
Karthik Ravishankar	Kanav Madhura	Ankit Raheja	Joey Yu Luo	Yiyu Wang
Zaidan Wu	Dylan Jamner	Yijie He	Peter Wei	Tu Cao

Name: M. Underhill Wisc email: _____

I pledge that the work on this exam is entirely my own. I understand that the penalties for cheating may include an F in the course and referral to my dean for further action.

Student Signature: _____

READ THE FOLLOWING INFORMATION.

- This is a 120-minute exam. It consists of 9 problems for a total of 100 points; the exam is 7 sheets of paper, including this cover sheet. It is your responsibility to make sure that you have a complete exam.
- Books, notes, calculators, phones, and other aids are not allowed.
- Problems are spaced out to allow ample room for work. You may use the last page for scratch work, but it will not be graded. Do not unstaple or remove pages as they can be lost in the grading process. **An incomplete exam packet will result in an automatic zero.**
- Only complete, well written, and neat solutions will be awarded full credit. Remember that all claims must be supported. Responses which do not meet these qualifications may be awarded some partial credit.
- If making multiple attempts at a solution, indicate clearly which attempt you'd like to be graded by crossing out the other answers. Multiple attempts will not be graded.
- When doing multiple choice and true/false questions, make sure to fully fill in the chosen bubble.

Correct mark: ●

Incorrect marks: ○ ⊗ ⊘ ⊙

- Vector space notation reminders: M_{mn} is the vector space of $m \times n$ real matrices with standard matrix addition and scalar multiplication, P_d is the vector space of polynomials (in x) of degree at most d with standard polynomial addition and scalar multiplication, $\mathbf{0}$ denotes the zero vector in a vector space V . $\mathbb{R}^n$ uses the standard vector addition/scalar multiplication and dot product unless otherwise noted.
- Matrix notation reminders: O denotes the zero matrix, and I_n denotes the $n \times n$ identity matrix. $\text{tr}(A)$ denotes the trace of a matrix A .
- Additional notation reminders: $M_{DB}(T)$ denotes the matrix representation of the linear transformation $T : V \rightarrow W$ using ordered bases B for V and D for W . $M_B(T)$ denotes the matrix representation of the linear operator T with respect to the ordered basis B . $C_B(\mathbf{v})$ denotes the coordinate representation of the vector $\mathbf{v}$ with respect to the ordered basis B .

DO NOT BEGIN THIS EXAM UNTIL SIGNALLED TO DO SO.

1. (7 points) Given $A = \begin{bmatrix} 2 & 3 & 0 \\ 0 & -2 & 4 \\ 1 & 2 & -1 \end{bmatrix}$, $C^{-1} = \begin{bmatrix} 2 & 4 & 1 \\ 2 & 6 & 4 \\ 1 & 2 & 2 \end{bmatrix}$, and $D = \begin{bmatrix} 2 & 4 & 1 \\ 0 & 1 & 4 \\ 3 & 2 & 9 \end{bmatrix}$, find B if $A + BC = D$. Show all work.

$$BC = D - A = \begin{bmatrix} 2 & 4 & 1 \\ 0 & 1 & 4 \\ 3 & 2 & 9 \end{bmatrix} - \begin{bmatrix} 2 & 3 & 0 \\ 0 & -2 & 4 \\ 1 & 2 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 1 \\ 0 & 3 & 0 \\ 2 & 0 & 10 \end{bmatrix}$$

$$B = (D - A)C^{-1} = \begin{bmatrix} 0 & 1 & 1 \\ 0 & 3 & 0 \\ 2 & 0 & 10 \end{bmatrix} \begin{bmatrix} 2 & 4 & 1 \\ 2 & 6 & 4 \\ 1 & 2 & 2 \end{bmatrix} = \begin{bmatrix} 3 & 8 & 6 \\ 6 & 18 & 12 \\ 14 & 28 & 22 \end{bmatrix}$$

Incorrect mult

$$"B" = C^{-1}(D - A) = \begin{bmatrix} 2 & 4 & 1 \\ 2 & 6 & 4 \\ 1 & 2 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 & 1 \\ 0 & 3 & 0 \\ 2 & 0 & 10 \end{bmatrix} = \begin{bmatrix} 2 & 14 & 12 \\ 8 & 20 & 42 \\ 4 & 7 & 21 \end{bmatrix}$$

2. (13 points total) Let W be the vector space of 2×2 symmetric matrices (using standard matrix addition/scalar multiplication), and define $T : P_2 \rightarrow W$ by $T(p(x)) = \begin{bmatrix} p'(0) & p(0) \\ p(0) & 0 \end{bmatrix}$. You can assume (without proof) that T is a linear transformation.

a. (8 points) Using ordered basis $B = (4x+1, x^2, x^2-2)$ for P_2 and ordered basis

$D = \left(\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right)$ for W , find the matrix representation $M_{DB}(T)$. Show all work.

$$T(4x+1) = \begin{bmatrix} 4 & 1 \\ 1 & 0 \end{bmatrix} \xrightarrow{\vec{u}_1} C_D(\vec{u}_1) = \begin{bmatrix} 4 \\ 0 \\ 1 \end{bmatrix}$$

$$T(x^2) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \xrightarrow{\vec{u}_2} C_D(\vec{u}_2) = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \quad M_{DB}(T) = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & -2 \end{bmatrix} = M$$

$$T(x^2-2) = \begin{bmatrix} 0 & -2 \\ -2 & 0 \end{bmatrix} \xrightarrow{\vec{u}_3} C_D(\vec{u}_3) = \begin{bmatrix} 0 \\ 0 \\ -2 \end{bmatrix}$$

b. (5 points) Is T an isomorphism? Justify your answer.

No. Many justifications possible.
 (1) $\det(M) = 0$ (b/c row of zeros), so M is not invertible, so T is not an isomorphism.

(2) $\dim \text{im } T = \text{rank}(M) = 2 < 3 = \dim(W)$, so T is not onto.

(3) $\dim \ker T = \dim \text{null}(M) = 1 > 0$, so T is not one-to-one.

(4) We see that $T(x^2) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = \vec{0}$ in W , so T has a nontrivial kernel, so T is not one-to-one.

3. (20 points, 2 points each) Clearly mark the correct answer for each of the following by **completely** filling in the appropriate bubble. **No justification is needed.**

a. (True/False) If $\mathbf{v}$ is a 3×1 vector, and A is a 3×3 matrix, then $\mathbf{v}A$ is a 3×1 vector.

- ☐ True
☒ False

$$3 \times 1 \quad 3 \times 3 \quad \times.$$

b. (True/False) If A is a 4×4 matrix with $\text{tr}(A) = 0$, then A is invertible.

- ☐ True
☒ False

$$\text{ex: } A = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \det(A) = \text{tr}(A) = 0.$$

c. (True/False) The subspace $W = \left\{ p \text{ in } P_{10} \mid \int_0^1 p(x) dx = 0 \right\}$ of P_{10} is finite-dimensional.

- ☒ True
☐ False

$$\dim(P_{10}) = 11, \text{ so } \dim(W) < 11.$$

d. (True/False) The function $F: \mathbb{R} \rightarrow \mathbb{R}$ by $F(x) = 2x + 1$ is an isomorphism.

- ☐ True
☒ False

$$F \text{ is not a linear transformation.}$$

e. (True/False) Suppose A is a 3×3 matrix with $A = A^{-1}$. Then $\det(A) = 1$ or $\det(A) = -1$.

- ☒ True
☐ False

$$\det(A) = \det(A^{-1}) = \frac{1}{\det(A)} \\ \Rightarrow \det(A)^2 = 1.$$

f. (True/False) If $\mathbf{x}$ and $\mathbf{y}$ are orthogonal vectors in an inner product space, then $\mathbf{x}$ and $\mathbf{x} + \mathbf{y}$ must also be orthogonal.

- ☐ True
☒ False

$$\text{ex: use dot product.} \\ \vec{x} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{y} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad \vec{x} + \vec{y} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

g. (Multiple Choice-choose one) Which vector space is isomorphic to P_3 ?

- ☐ $\mathbb{R}^4$ ✓

☐ The nullspace of $A = \begin{bmatrix} 1 & 2 & 4 & 3 & -1 \\ 2 & 4 & 8 & 6 & -2 \end{bmatrix} \xrightarrow{-2r_1 + r_2 \rightarrow r_2} \begin{bmatrix} 1 & 2 & 4 & 3 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$

- ☐ M_{22} ✓

- ☒ All of the above.

$$\dim(P_3) = 4.$$

4 free variables

h. **(Multiple Choice-choose one)** Suppose A is a 3×3 matrix which is row-equivalent to

$$B = \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}. \text{ Which of the following must be true about } A?$$

- ☐ $A\mathbf{x} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$ has no solutions. *maybe, maybe not.*
☐ $\lambda = 1$ is an eigenvalue of A . *maybe, maybe not.*
☒ $\lambda = 0$ is an eigenvalue of A . *✓. $A\vec{x} = \vec{0}$ has a nontrivial soln.*
☐ The columns of A must be linearly independent. *X*
☐ All of the above.
☐ None of the above.

i. **(Multiple Choice-choose one)** Suppose $\{\mathbf{u}, \mathbf{v}\}$ is a basis for a vector space V . Which of the following is also a basis for V ? *$\hookrightarrow \dim(V) = 2$.*

- ☐ $\{\mathbf{u} + \mathbf{v}\}$ *X*
☒ $\{\mathbf{u} + \mathbf{v}, 2\mathbf{u} - \mathbf{v}\}$ *✓. not scalar mult.*
☐ $\{\mathbf{u} - \mathbf{v}, \mathbf{v} - \mathbf{u}\}$ *X $\vec{u} - \vec{v} = -(\vec{v} - \vec{u})$ not indep.*
☐ $\{\mathbf{u} + \mathbf{v}, \mathbf{u} + 2\mathbf{v}, \mathbf{v}\}$ *X*
☐ All of the above.
☐ None of the above.

j. **(Multiple Choice-choose one)** Let V be $\mathbb{R}^2$ with the following vector addition $\oplus$ and scalar multiplication $\odot$:

$$\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} \oplus \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} = \begin{bmatrix} x_1 + x_2 + 3 \\ y_1 + y_2 \end{bmatrix}, \quad c \odot \begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = \begin{bmatrix} cx_1 + 3c - 3 \\ cy_1 \end{bmatrix}$$

Which of the following is the zero vector $\mathbf{0}$ for this vector addition $\oplus$?

- ☐ $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$
☐ $\begin{bmatrix} 3 \\ 0 \end{bmatrix}$
☒ $\begin{bmatrix} -3 \\ 0 \end{bmatrix}$
☐ $\begin{bmatrix} 3 \\ 3 \end{bmatrix}$
☐ None of the above.

4. (12 points total) Consider the vector space P_1 , and define the function

$$\langle p(x), q(x) \rangle = \int_0^1 p(x)q(x)dx$$

for $p(x), q(x)$ in P_1 .

You can assume that $\langle p(x), q(x) \rangle$ is an inner product for P_1 .

- a. (4 points) With respect to this inner product, $\langle p(x), q(x) \rangle$, find the distance between $3x + 1$ and $x - 3$. Show all work, but you do not need to simplify your final numerical answer.

$$3x+1-(x-3)=2x+4.$$

$$\|2x+4\| = \sqrt{\int_0^1 \underbrace{(2x+4)^2}_{4x^2+16x+16} dx} = \sqrt{\left. \frac{4x^3}{3} + 8x^2 + 16x \right|_0^1} = \sqrt{\frac{4}{3} + 8 + 16}$$

- b. (4 points) With respect to this inner product, $\langle p(x), q(x) \rangle$, show that the polynomials $p = 3 - 3x$ and $q = 6x - 2$ are orthogonal. Show all work.

$$\begin{aligned} \langle p, q \rangle &= \int_0^1 \underbrace{(3-3x)(6x-2)}_{-18x^2+24x-6} dx \\ &= \left. -6x^3 + 12x^2 - 6x \right|_0^1 \\ &= -6 \cdot 1^3 + 12 \cdot 1^2 - 6 \cdot 1 = 12 - 12 = 0 \checkmark. \end{aligned}$$

- c. (4 points) With respect to this inner product, $\langle p(x), q(x) \rangle$, using $p = 3 - 3x$ and $q = 6x - 2$, create an orthonormal basis for P_1 . Show all work.

$$\|p\| = \sqrt{\int_0^1 \underbrace{(3-3x)^2}_{9-18x+9x^2} dx} = \sqrt{\left. 9x - 9x^2 + 3x^3 \right|_0^1} = \sqrt{9-9+3} = \sqrt{3}.$$

$$\|q\| = \sqrt{\int_0^1 \underbrace{(6x-2)^2}_{36x^2-24x+4} dx} = \sqrt{\left. 12x^3 - 12x^2 + 4x \right|_0^1} = \sqrt{12-12+4} = \sqrt{4} = 2.$$

$$\left\{ \frac{3-3x}{\sqrt{3}}, \frac{6x-2}{2} \right\} \text{ are an orthonormal basis for } P_1.$$

5. (16 points total) Let $T : P_1 \rightarrow P_1$ be a linear transformation with

$$M_B(T) = \begin{bmatrix} 6 & 2 \\ -2 & 10 \end{bmatrix} \text{ with respect to the ordered basis } B = (x+1, x-3).$$

a. (4 points) Compute $T(2x-1)$. Show all work.

$$\begin{aligned} a(x+1) + b(x-3) &= 2x-1 \rightarrow \begin{array}{l} a+b=2 \\ -(a-3b=-1) \end{array} \\ \hline 4b &= 3 \Rightarrow b = 3/4 \quad a = 2 - 3/4 = 5/4 \end{aligned}$$

$$\begin{bmatrix} 6 & 2 \\ -2 & 10 \end{bmatrix} \begin{bmatrix} 5/4 \\ 3/4 \end{bmatrix} = \begin{bmatrix} \frac{30+6}{4} \\ \frac{-10+30}{4} \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \end{bmatrix} \rightarrow \boxed{9(x+1) + 5(x-3)}$$

b. (5 points) Is T diagonalizable? Justify your answer.

$$\begin{aligned} \det(\lambda I - M) &= \det \begin{bmatrix} \lambda-6 & -2 \\ 2 & \lambda-10 \end{bmatrix} = (\lambda-6)(\lambda-10) + 4 \\ &= \lambda^2 - 16\lambda + 60 + 4 = \lambda^2 - 16\lambda + 64 \\ &= (\lambda-8)^2 \quad \lambda = 8. \end{aligned}$$

$$\begin{aligned} 8I - M &= \begin{bmatrix} 2 & -2 \\ 2 & -2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix} \quad \begin{array}{l} x_1 - x_2 = 0 \Rightarrow x_1 = t \\ x_2 = t \end{array} \\ \begin{bmatrix} t \\ t \end{bmatrix} &= t \begin{bmatrix} 1 \\ 1 \end{bmatrix} \text{ is the eigenspace.} \end{aligned}$$

This matrix has one eigenvalue w/ alg. mult 2,
but geo. mult 1.

$\Rightarrow M$ is not diagonalizable.

$\Rightarrow T$ is not diagonalizable.

(Problem 5 continued)

Again, let $T : P_1 \rightarrow P_1$ be a linear transformation with

$$M_B(T) = \begin{bmatrix} 6 & 2 \\ -2 & 10 \end{bmatrix} \text{ with respect to the ordered basis } B = (x+1, x-3).$$

- c. (3 points) Suppose $P_{B \leftarrow D} = \begin{bmatrix} 2 & 1 \\ 3 & 1 \end{bmatrix}$ is the change matrix from unknown basis $D = (p_1, p_2)$ to B . Find p_1 and p_2 .

$$C_B(p_1) = \begin{bmatrix} 2 \\ 3 \end{bmatrix} \Rightarrow p_1 = 2(x+1) + 3(x-3)$$

$$C_B(p_2) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \Rightarrow p_2 = 1(x+1) + 1(x-3)$$

- d. (4 points) For the same basis D as in part (c), use the Similarity Theorem find $M_D(T)$.
(Hint: You do NOT need to know D to solve this problem). Show all work.

$$M_D(T) = P_{D \leftarrow B} M_B(T) P_{B \leftarrow D}$$

$$P_{D \leftarrow B} = (P_{B \leftarrow D})^{-1} = \frac{1}{\underbrace{2 \cdot 1 - 3 \cdot 1}_{=-1}} \begin{bmatrix} 1 & -1 \\ -3 & 2 \end{bmatrix} = \begin{bmatrix} -1 & 1 \\ 3 & -2 \end{bmatrix}$$

$$M_D(T) = \begin{bmatrix} -1 & 1 \\ 3 & -2 \end{bmatrix} \begin{bmatrix} 6 & 2 \\ -2 & 10 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 1 \\ 3 & -2 \end{bmatrix} \begin{bmatrix} 18 & 8 \\ 26 & 8 \end{bmatrix}$$

$$= \begin{bmatrix} 8 & 0 \\ 2 & 8 \end{bmatrix}$$

$\uparrow$
 $18 \cdot 3 - 2 \cdot 26 = 2$

6. (8 points) Consider the vector space $\mathbb{R}^3$, and consider the set U of vectors $\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$ where $x_1 + 3x_2 = x_3$

and $x_1x_2 \geq 0$, i.e. $U = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \mid x_1 + 3x_2 = x_3, x_1x_2 \geq 0 \right\}$. Is U a subspace of $\mathbb{R}^3$? Fully justify your answer.

No. Consider

$$\begin{bmatrix} -1 \\ -4 \\ -13 \end{bmatrix} + \begin{bmatrix} 2 \\ 2 \\ 8 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ -5 \end{bmatrix}$$

$-1 + 3(-4) = -13$ and $(-1)(-4) = 4 \geq 0$ ✓
 in U

$2 + 3 \cdot 2 = 8$ and $(2)(2) = 4 \geq 0$ ✓
 in U

$1 + 3(-2) = -5$, but $(1)(-2) = -2 < 0$. ✗

U is not closed under addition, so U is not a subspace of $\mathbb{R}^3$.

7. (9 points total) Consider $\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ in $\mathbb{R}^2$, and define the function

$$\langle \mathbf{u}, \mathbf{v} \rangle = 2u_1v_1 + 3u_2v_2.$$

a. (4 points) Verify the following inner product property for $\langle \mathbf{u}, \mathbf{v} \rangle$:

$$\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle \text{ for all } \mathbf{u}, \mathbf{v} \text{ in } \mathbb{R}^2.$$

$$\begin{aligned} \left\langle \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}, \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} \right\rangle &= 2u_1v_1 + 3u_2v_2 \\ &= 2v_1u_1 + 3v_2u_2 \\ &= \left\langle \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}, \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \right\rangle \quad \text{!} \end{aligned}$$

For part (b), assume the remaining inner product properties are true for

$$\langle \mathbf{u}, \mathbf{v} \rangle = 2u_1v_1 + 3u_2v_2.$$

b. (5 points) With respect to this inner product $\langle \mathbf{u}, \mathbf{v} \rangle$, find a basis for W , where W is the subspace of $\mathbb{R}^2$ of vectors orthogonal to $\begin{bmatrix} 5 \\ -4 \end{bmatrix}$. Show all work.

$$\begin{aligned} \text{Let } \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} \text{ be in } W. \text{ Then} \\ \left\langle \begin{bmatrix} 5 \\ -4 \end{bmatrix}, \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} \right\rangle &= 5 \cdot 2v_1 + 3(-4)v_2 = 0 \\ &10v_1 - 12v_2 = 0 \end{aligned}$$

$$\begin{bmatrix} 10 & -12 & : & 0 \end{bmatrix} \xrightarrow{10 \rightarrow r_1} \begin{bmatrix} 1 & -6/5 & : & 0 \end{bmatrix}$$

$v_2 = t$ (free)
 $v_1 = 6/5 t$

$$W = \left\{ t \begin{bmatrix} 6/5 \\ 1 \end{bmatrix} \right\}, \text{ so a basis is}$$

$$\boxed{\left\{ \begin{bmatrix} 6/5 \\ 1 \end{bmatrix} \right\}}$$

8. (5 points) Let $A = \begin{bmatrix} 0 & 2 & 1 \\ 2 & 4 & 3 \\ 0 & 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 5 & -1 & 1 \\ 3 & -3 & 1 \\ c & -10 & 3 \end{bmatrix}$. If A and B are similar matrices, find c .

Show all work.

Same det: $\det(A) = -2(2 \cdot 1 - 1) = -2(1) = -2.$

$\det(B) = 5(-9 + 10) + 1(9 - c) + 1(-30 + 3c)$

$$-2 = 5 + 9 - c - 30 + 3c = -16 + 2c$$

$$14 = 2c \Rightarrow \boxed{c = 7}$$

9. (10 points total, 2 points each) Suppose A is a 3×7 matrix with $\text{rank}(A) = 3$. Answer the following questions about A , if possible. Justification is not required.

a. (Multiple Choice-choose one) What is the $\dim(\text{null } A)$?

- ☐ 0
☐ 1
☐ 2
☐ 3
☒ 4
☐ Not enough information to answer.

$$7 - 3 = 4$$

b. (Multiple Choice-choose one) Is $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ in the column space of A ?

- ☒ Yes
☐ No
☐ Not enough information to answer.

b/c $\text{rank} = 3 = \dim \text{col}(A)$, so all of $\mathbb{R}^3$ is in $\text{col}(A)$.

c. (Multiple Choice-choose one) Does the linear system $A\mathbf{x} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ have...

- ☐ No solution
☐ One solution
☒ Infinitely many solutions
☐ Not enough information to answer.

consistent w/ free variables.

d. (Multiple Choice-choose one) For $T : \mathbb{R}^7 \rightarrow \mathbb{R}^3$ defined by $T(\mathbf{x}) = A\mathbf{x}$, is T onto?

- ☒ Yes
☐ No
☐ Not enough information to answer.

e. (Multiple Choice-choose one) For $T : \mathbb{R}^7 \rightarrow \mathbb{R}^3$ defined by $T(\mathbf{x}) = A\mathbf{x}$, is T one-to-one (also called injective)?

- ☐ Yes
☒ No
☐ Not enough information to answer.

Scratch paper page!! (Must be turned in with exam)

Back of scratch paper page! (Must be turned in with exam)