

Land Sharing or Land Sparing? Implications for Farmland Birds of Conservation Concern in France

Previously “Disentangling land sharing and land sparing effects on bird abundance in France using spatio-temporal Bayesian modeling”

Adélie Erard ^{1*}, Ottmar Cronie ^{2*},

Raphaël Lachière-Rey ^{3,1} and Romain Lorrillière ^{4,5}

^{1*} Université Paris Cité, CNRS, MAP5, F-75006, Paris, France.

^{2*} Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, Gothenburg, SE-412 96, Sweden.

³ Inria Paris, MathNet, Paris, F-75013, France.

⁴ MNHN, CRBPO, F-75005, Paris, France.

⁵ MNHN, Sorbonne Université, CNRS, Centre d’Écologie et des Sciences de la Conservation, CESCO, F-75005, Paris, France.

*Corresponding author(s). E-mail(s): aerard@math.cnrs.fr; ottmar.cronie@gu.se;

Contributing authors: raphael.lachieze-rey@math.cnrs.fr; romain.lorrilliere@mnhn.fr;

Abstract

Purpose: Farmland bird populations have experienced widespread declines across Europe, raising urgent questions about how agricultural landscapes should be structured to reconcile biodiversity conservation and food production. Land sparing and land sharing represent two contrasting strategies, yet empirical evidence remains mixed.

Objectives: We aim to quantify responses of bird communities along a land sharing–sparing gradient and identify which landscape management strategies best support different ecological groups.

Methods: Using the French Breeding Bird Survey, we analyzed 810 locations

monitored between 2001–2024. A land sharing–sparing gradient was quantified using forest area, agricultural heterogeneity, and landscape diversity indices. Bird counts were modeled with negative binomial spatio-temporal Bayesian models (INLA), incorporating climate, spatial random fields, and temporal autocorrelation.

Results: Forest specialists responded positively to greater forest area, supporting land sparing. Generalists benefited from higher agricultural heterogeneity, consistent with land sharing. Unexpectedly, farmland specialists did not benefit from increased heterogeneity, and conservation-priority farmland species showed negative associations with natural elements.

Conclusion: No single land-use strategy maximizes abundance across all bird groups. Land sparing favors forest specialists and conservation-priority species, while land sharing benefits generalists. Context-dependent landscape planning and customised agricultural practices are essential.

Keywords: Land sharing and land sparing, farmlands birds, INLA, conservation

Acknowledgment

This work was supported by the Chaire “Modélisation Mathématique et Biodiversité” of Veolia-École polytechnique-Muséum national d’Histoire naturelle-Fondation X, through Adélie Erard’ Ph.D. scholarship. We thank the thousands of volunteers who took part in the French breeding bird surveys.

1 Introduction

With the current global biodiversity decline (Burns et al. 2021; Inger et al. 2015), land use allocation is a major challenge for biodiversity conservation. Agricultural landscapes are central to this issue, as they cover a large proportion of European territory and harbour a substantial part of biodiversity.

Several studies show that farmland bird populations are declining in Europe (Donald et al. 2001; Bowler et al. 2019). This decline is mainly explained by the intensification of agricultural practices (Rigal et al. 2023; Genty et al. 2026) and pesticide use (Monnet et al. 2026; Perrot et al. 2025). Heldbjerg et al. (2018) also shows that ground-nesting birds on the ground are in significant decline in Denmark whereas birds nesting elsewhere are less impacted. This raises the question of how land should be allocated to maximize biodiversity? Two main land-use strategies are commonly discussed (Fischer et al. 2014; Grass et al. 2019; Fahrig 2020, 2013): land sparing and land sharing. The **land sparing** approach separates highly intensive agricultural production from conservation areas in order to minimize trade-offs between food production and biodiversity preservation. In such landscapes, agricultural areas are typically characterized by large, homogeneous patches and intensive farming practices, which often result in low biodiversity. In contrast, biodiversity is concentrated in relatively large protected areas such as national nature reserves or forests. By contrast, the **land sharing** approach aims to support both food security and biodiversity by

promoting their coexistence within the same landscapes. These landscapes are generally composed of many small agricultural fields interspersed with trees, hedgerows, or semi-natural elements. As a result, they tend to be more heterogeneous, with higher connectivity between habitat patches, and farming practices are usually less intensive.

Today, there is no consensus on how to allocate land use to maximize biodiversity and agricultural output. Indeed, on the one hand (Phalan et al. 2011; Edwards et al. 2015) show that land sparing is better to conserve biodiversity in Ghana, India and Colombia. On the other hand, (Riva and Fahrig 2022) show that small patches contain higher diversity. Moreover, (Lamb et al. 2019) show that land sparing would more likely only benefit forest birds and would disadvantage farmland birds in the United Kingdom. (Finch et al. 2019) proposes a solution based on three compartments enabling some large forest patches to conserve woodland birds, some low-intensity agriculture to conserve farmland birds and some big fields to ensure food security.

These previous results suggest that the effect of land sharing or land sparing on bird abundance depends on both the landscape and species' needs. These needs correspond to the ecological niches of the species which may be narrow for specialist species and broader for generalists (Devictor et al. 2010).

To resolve this uncertainty in the French agricultural landscapes, which covers more than half the country with highly diverse farming practices (Nagy and Soós 2025), we need robust monitoring approaches. Birds serve as ideal indicators because they are easy to sample, cover a wide range of life-history traits, are present in all habitats and trophic levels (Padoa-Schioppa et al. 2006; Karen Aghababayan 2024), and are sensitive to environmental stressors (Canterbury et al. 2000; Gregory et al. 2003; Rigal et al. 2023).

Moreover, citizen science facilitates bird data collection because many volunteers are skilled at bird identification. Hence, thanks to thousands of skilled French bird-watchers we gathered an extensive database from a breeding bird survey called the French Breeding Bird Surveys (Jiguet et al. 2012; Fontaine et al. 2021).

As indicators of land sharing and land sparing are complex, we need to take into account many variables that structure the landscape and bird populations. Moreover, to account for the spatio-temporal structure, a Bayesian framework and inference with INLA (Rue et al. 2009) appears to be particularly suitable because it efficiently handles the hierarchical structure of spatio-temporal data with multiple covariates.

In this study, we use spatio-temporal Bayesian modeling to disentangle the effects of land sharing and land sparing scenario on bird populations, with a focus on farmland and forest birds with a focus on species of conservation concern. We formulated the following hypotheses:

1. Land sharing is expected to benefit generalist and farmland species. In land-sharing landscapes, the higher diversity of habitat patches may provide a wider range of nesting opportunities for generalist species. For farmland species, lower agricultural intensity under land sharing is predicted to promote breeding success and survival.
2. Land sparing is expected to benefit forest species, particularly those of conservation concern. Forest specialists are generally sensitive to human disturbance; therefore, larger contiguous areas allocated to biodiversity conservation under land sparing are predicted to enhance their abundance.

139 2 Materials and Methods

140

141 2.1 Data

142

143 2.1.1 Bird abundance data

144 The data are provided by the French Breeding Bird Surveys (Jiguet et al. 2012;
145 Fontaine et al. 2021). It aims to survey the common French breeding bird populations
146 over time and space in France. This is a participatory program where thousands of
147 skilled volunteers will count birds. These volunteers get randomly assigned a 2 km by
148 2 km square, next to the place they live, by the French National Museum of Natural
149 History. Once this square is assigned, the volunteer needs to choose ten evenly dis-
150 tributed points inside the square. Then, each year the volunteer has to visit the square
151 twice during breeding season to record bird abundance. A visit of the square consists
152 of staying 5 minutes at each point and recording every seen or heard bird (Jiguet et al.
153 2012).

154 In order to have clean data, we removed every square that has not been observed
155 in the right times and dates. We also removed squares that do not have ten points in
156 it (around 2% of the data). Then we chose to aggregate data to the scale of the square
157 (sum of all points) to feasibly carry out statistical analysis made by INLA and remove
158 detectability bias (Nabias et al. 2024).

159 To have enough temporal depth in our analysis we only kept time series of at least
160 five years. As our goal is to study the effect of different kinds of agricultural practices
161 on bird population, we removed all squares that have three or more point classified as
162 urban by the observer or at least 25% of urban land use in the square.

163 It left us with a total of 810 squares that have been observed at least five years
164 in a row. We used the STOC classification (Fontaine et al. 2021) of bird to categorize
165 birds with their main types of habitat adding two species to agricultural birds: *Passer*
166 *montanus* (Barlow et al. 2020) and *Burhinus oedicnemus* (Hume and Kirwan 2020).

167 To disentangle the effects of land sharing and land sparing on French bird popula-
168 tions, we divided birds into groups of habitat specialization (agricultural, forest and
169 generalist birds) given by (Fontaine et al. 2021; Julliard et al. 2006). Then, we used the
170 French IUCN list (UICN France et al. 2016) to classify the species in two categories:
171 least concerned and birds with conservation concern. In the French IUCN list (UICN
172 France et al. 2016) they mention birds for which the category changes between 2008
173 and 2016. As our dataset goes from 2001 to 2024, we considered that a species is with
174 conservation concern if it is not "least concerned" in 2008 or 2016.

175 Maps of distribution of birds' observation by group are shown in Supplementary
176 material Section 1 (Figure SI-1 for generalist species, Figure SI-2 for forest species,
177 Figure SI-3 for agricultural species, Figure SI-4 for forest species with conservation
178 concern and Figure SI-5 for agricultural species with conservation concern).

179

180 2.1.2 Environmental data

181

182 Climate data is taken from WordClim (Harris et al. 2020). For each square and year
183 we retrieved minimum and maximum temperatures and total amount of precipitation
184 of the current and last spring (March to June) as well as minimum and maximum

of last summer (July and August). Then we added the latitude, recentered to be positive when in the North and negative otherwise. This variable also serves through interactions with the location and climate variables, allowing us to check if our model works well by finding the same results as in (Thepault et al. 2025).

We used CORINE land cover (European Environment Agency 2019) to retrieve land cover variables. This database contains all land uses in France in a raster format for years 2000, 2006, 2012 and 2018. The different kind of land uses are described in 44 groups (Section 5 of Supplementary material). For each year, we retrieved in each square the surface of agricultural (all CORINE codes starting by 2) and water areas (all CORINE codes starting by 5), forest (all CORINE codes starting by 31), natural (all CORINE codes starting by 32 or 33), water (all CORINE codes starting by 4 or 5) and artificial lands (all CORINE codes starting by 1). This provides a general assessment of the use of each square. We did a linear interpolation of the data when it was not constant between two years of data.

2.1.3 Land sharing and sparing indicators

To create indicator of land sharing and land sparing we took the intersection of CORINE patches and a disk of radius $10km$ centered in the middle of each square. This allowed us to obtain, for each square, the number of patches intersecting the disk, their area within the disk and their type. As the area of forest around the square is a great indicator of land sparing we retrieved the total area of forest patches intersecting a disk of 10 km radius around the center of the square (without any distinction between different forest types).

In order to have an indicator of the heterogeneity of agricultural practices, we computed a Shannon index (Shannon 1948) on agricultural land uses around the squares where the “species” is the kind of patch and the abundance the number of each kind of patch. Shannon’s indices will give us an idea of the diversity of agricultural land uses around each square. If this index is high then there is a high diversity in land uses, saying we are more in a land sharing scenario. Otherwise, it means that we are in a monocultural scenario meaning that we are in a land sparing scenario. An example of the land use around a square is given in Figure 1.

The last two variables we added are the forest area and agriculture with significant amount of nature in a radius of 10 kilometers around the point of interest. A higher forest area implies a land sparing scenario whereas a higher amount of agriculture with significant amount of nature implies a land sharing scenario.

2.1.4 Correlation study and Principal Component Analysis (PCA) on the land cover variables

The land use of each square consists of six highly correlated variables as they all sum up to the area of the square (4 square kilometers). We then use a PCA to summarize these variables onto fewer variables that are not correlated. After computations, we can summarize into three component explaining 80.8% of the variance of the data (Figure SI-7c). The axes are represented in Figures SI-7a and SI-7b. The first axis separates the agricultural squares (negative) from the natural and forest squares (positive). The




Fig. 1 Example of the land use in a 10 km radius disk around a square (black line). Each color represents a different kind of land use following the table in Supplementary material Section 5 (colors used are the official CORINE colors). Shannon's indices are calculated only with agricultural patches (codes starting with a two). In this case there are 68 agricultural patches spread in four categories. The black dots represent ten points of samples.

second axis separates urban (and water) squares (positive) with the others. The third axis separates closed squares (forest and cities) from the open squares (flat lands). We call these axes PCA1, PCA2 and PCA3.

2.2 Statistical approach

2.2.1 Model

Bird abundance may vary due to multiple drivers such as land-use change, climate variability, agricultural practices, and interactions with other species. To quantify these effects, we modelled bird counts using a negative binomial regression model, which accounts for overdispersion in the data due to the lack of observation as well as for real bird distribution. Formally, the expected abundance at location s and year t is defined as:

$$\log \lambda(s, t) = \beta_0 + \sum_k \beta_k X_k(s, t) + \sum_{i,j} \beta_{i,j} X_i(s, t) X_j(s, t) + w_{s,t},$$

where $X_k(s, t)$ denote environmental covariates (e.g. climate, land cover), and $w_{s,t}$ represents residual spatio-temporal variability. Interaction terms, $X_i(s, t) X_j(s, t)$, allow us to capture non-additive effects between environmental drivers. In our case, we used the separable case for the spatio-temporal variability:

$$w_{t,s} = w_s + w_t,$$

where w_t is an autoregressive model and w_s is a Matérn random field. For the temporal effect, there is a correlation between years, modelled in the following way:

$$w_t = \phi w_{t-1} + \epsilon_t,$$

$$\epsilon_t \sim \mathcal{N}(0, \tau^{-1}),$$

where ϕ is the auto-correlation parameter and τ is the precision parameter. For the spatial Matérn random field, the covariance is as following (as in (Krainski et al. 2018)):

$$\text{Cov}(w_s, w_{s'}) = \sigma^2 \frac{1}{2^{\nu-1} \Gamma(\nu)} (\kappa \|s - s'\|)^\nu K_\nu(\kappa \|s - s'\|);$$

where: σ^2 is the marginal variance of the process; ν is the smoothness parameter (usually set to 1); κ is a scaling parameter (linked to the range ρ of the process by: $\rho = \sqrt{8\nu/\kappa}$); and K_ν is the modified Bessel function of the second kind. The range ρ of the process gives information on the spatial correlation range between points.

2.2.2 Inference

We relied on a Bayesian framework for inference. Instead of using the traditional Monte Carlo Markov Chain algorithm to get posterior marginals, we used the Integrated Nested Laplace Approximation (INLA) developed by (Rue et al. 2009). It provides fast and accurate deterministic approximations of posterior distributions for latent Gaussian models.

In particular, it computes posterior marginals for latent effects (regression coefficients) as well as for hyperparameters (variance of spatial and temporal random effects). This approach makes it possible to fit computationally demanding spatio-temporal models efficiently.

Prior choice. As it is a Bayesian framework, we have to choose priors. They are summarized in Table 1. For the spatial field, we chose to use penalized complexity (PC) priors (Simpson et al. 2017) because it helps in choosing less complex model by penalizing the complexity of the latent Gaussian random field.

The spatial random field needs two parameters: the range, giving information about the spatial correlation range between two points and the variance of the distance of interaction σ (in the Matérn field). We chose to set the interaction range at 100 km because birds return to the same nesting sites (Jiguet et al. 2012; Fontaine et al. 2021). For the variance we chose $\sigma = 1$ as a weakly informative prior.

All the other priors are default INLA priors that are weakly informative and are summarized in Table 1.

Mesh choice. The first thing to do is to choose a mesh for the spatial random field. The mesh is automatically constructed by INLA, but we need to choose three parameters: the inner edge, the outer edge and the cutoff value. To avoid any boundary effects in the analysis, the mesh should be extended further away from the spatial domain (Lindgren et al. 2011). The inner edge sets the maximum length of an edge inside the spatial domain. In (Dambly et al. 2023) they suggest to use inner edge as 1/5 the prior range, we assumed an interaction range of approximately 100 km, so we

Table 1 Summary of prior distributions used in the INLA spatio-temporal models. PC stands for Penalized Complexity

Model component	Prior distribution	Interpretation / Notes
<i>Spatial random field</i>	PC on range and σ	$\Pr(\text{range} < 100\,000) = 0.99$, $\Pr(\sigma > 1) = 0.01$
<i>SPDE smoothness</i>	Fixed to $\alpha = 2$	Corresponds to a Matérn field with moderate smoothness ($\nu = 1$)
<i>Temporal random effect</i>	PC priors on ϕ and τ	Shrinks toward $\phi = 0$ (no auto-correlation) and small variance
<i>Fixed effects</i>	$\mathcal{N}(0, 10^6)$	Weakly informative / nearly flat prior
<i>Negative binomial dispersion</i>	$\log(\theta) \sim \text{Gamma}(1, 0.00005)$	Weak prior on overdispersion parameter

took 20 km. The outer edge sets the maximum length of an edge outside the spatial domain. In (Righetto et al. 2020) they show that the outer range does not make so much difference and shall be bigger than the prior range, we chose 200 km. The cutoff value sets the minimal distance between two points in the mesh. As we cannot have observations that are less than 2 km away we chose the cut-off to be 2 km.

In (Krainski et al. 2018) they mention that the mesh needs to be even in size and angle of the triangles of the mesh. It is the case with the values we chose (see Figure SI-6).

2.2.3 Model selection

We have five groups of species of interest : woodland species, farmland species, generalist species, woodland species with conservation concern and farmland species with conservation concern. We want to test the following set of variables: latitude, minimum temperature of last spring and last summer, maximum temperature of last spring and last summer, total amount of rain of last spring, the first three variables from the PCA on the land cover of each square (PCA1, PCA2, PCA3 from Section 2.1.4), the Shannon index on agricultural practices (Section 2.1.3), total surface of forest in a radius of 10 kilometers around the center of the square and the surface of farmland with significant amount of nature in a radius of 10 kilometers around the center of the square. To these variables, we added interactions between on the one hand maximum temperatures (of last spring and summer) and on the other hand latitude, surface of forest 10 kilometers around, surface of agriculture with significant amount of nature 10 kilometers around and Shannon index. This yielded a total of 12 variables and 8 interactions to test.

Testing possible model combinations for each group of species would be computationally intensive (around 10 seconds per model with more than 4000 models by group). We thus chose to perform stepwise selection to find the covariates that explain best the data, as in (Illian et al. 2013). The criterion we used to select is the Widely Applicable Information Criterion (WAIC) as for example in (Carson and Mills Fleming 2014). This criterion decreases when the fit of the model is better and penalizes the number of covariates. The WAIC gives a criterion of the goodness of fit of the model. However, it is a criterion that we can only use to compare models that are fitted to the exact same data and mesh parametrization (Dambly et al. 2023).

To test a wide range of models, we performed stepwise selection in three ways (Burnham and Anderson 2004): forward, backward and both. The forward procedure starts with the model with only random effects and then adds variables one by one. At each step, it selects the best variable to add to the model, according to the decrease of the WAIC score. When the two variables of an interaction are added, the corresponding interaction variable is added. Then, the backward procedure starts with the full model (all covariates and interaction) and remove one by one the covariates or interaction term along the same principle. Note that a covariate cannot be removed if an interaction term between this variable and another one is still in the formula. Finally, the "both" procedure starts as the forward procedure. After choosing a new variable to add, we add a backward step: we try to remove one by one each variable added in previous steps. We keep the model with the lowest WAIC after this backward step.

3 Results

We performed model selection for our five categories of birds. Detailed results can be found in Supplementary Table SI-1 for agricultural birds, Supplementary Table SI-2 for forest birds, Supplementary Table SI-3 for generalist birds, Supplementary Table SI-4 for agricultural birds with conservation concern and Supplementary Table SI-5 for forest birds with conservation concern.

First, we observed that in each category PCA1 has been chosen and is significant. Recall that this variable represents farmland squares when negative and forest squares when positive (Supplementary Figure SI-7a). Then all variables have been chosen at least in one group of birds. Regarding interactions, three of them have never been chosen by the selection procedure: Shannon index with last spring maximum temperature, Shannon index with last summer maximum temperature, and latitude with last summer maximum temperature.

Then, Table 2 and 3 indicate that, for all groups of species, there is a strong temporal autocorrelation ($\phi \approx 0.9$) and that the range of the Matérn field is not bigger than 40 km, meaning that the probability that two points more than 40 km apart depend on each other is very low. This distance aligns with the knowledge about bird breeding ecology as the breeding site fidelity and the distance of natal dispersal is known to be around 14 km (with 16 km of standard deviation) (Barbet-Massin et al. 2012). We can also observe that the overdispersion parameter is relatively high (from 11.64 to 47.25 Table 2 and 3).

Fixed effects are shown in Figure 2 for agricultural birds (A), forest birds (B) and generalist birds (C) and in Figure 3 for agricultural (A) and forest (B) birds with conservation concern.

The models use two kinds of **control variables** (land use and climate). Firstly, the **land uses** are described by the three first PCA axes: PCA1 (negative for agricultural squares and positive for forest squares), PCA2 (positive for urban square) and PCA3 (negative for open squares and positive for closed squares) (see 2.1.4). PCA1 is significant (when the confidence interval does not include zero) for all group of birds, positive for forest birds (0.44, with Confidence Interval [0.40, 0.49]), forest birds with

conservation concern (**0.78** [0.67, 0.88]) and generalist birds (0.03 [0.01, 0.06]), and negative for farmland birds (**-0.52** [-0.58, -0.46]) and farmland birds with conservation concern (**-0.62** [-0.68, -0.55]). PCA2 is selected and significant for forest birds (-0.07 [-0.10, -0.04]) and forest birds with conservation concern (0.08 [0.01, 0.16]). PCA3 is selected and significant for agricultural birds (-0.23 [-0.27, -0.18]) agricultural birds with conservation concern (-0.30 [-0.35, -0.26]) and generalist birds (0.03 [0.01, 0.05]). These results are in line with the usual hypotheses that the abundance of a group of birds increases in the habitat they are specialized in.

Then the other set of control variables was composed of **climate variables** and latitude. Last spring minimum temperature was selected and significant for forest birds (0.03 [0.01, 0.04]). Last spring maximum temperature was selected and significant for forest birds with conservation concern (-0.08 [-0.14, -0.03]). It was also selected but not significant for forest birds, agricultural birds and agricultural birds with conservation concern. Last spring total amount of rain was selected and significant for agricultural birds with conservation concern (-0.02 [-0.04, -0.01]). Last summer minimum temperature was selected but not significant for forest birds with conservation concern. Last summer maximum temperature was selected but not significant for agricultural birds, forest birds, and forest birds with conservation concern. The interaction between last spring maximum temperature and latitude was selected and significant for forest birds with conservation concern (-0.05 [-0.08, -0.02]). Although some climate variables were selected and significant for some models, we observe that these have limited effects because of the very small coefficients. However, when climate variables were selected and significant, results are globally in line with those of (Thepault et al. 2025).

In our approach the **land sharing and sparing** gradient is described thanks to three variables: agriculture with significant amount of nature, Shannon index and forest area in a radius of 10 kilometers around.

For agricultural birds (Figure 2-A) agriculture with a significant amount of nature and forest area 10 kilometers around were selected but not significant (respectively -0.05 [-0.11, 0.01] and -0.03 [-0.11, 0.04]). Some interaction between these variables and climate variables were also selected but not significant.

For forest birds (Figure 2-B), two variables of interest were selected and significant: forest area and agriculture with a significant amount of nature 10 kilometers around (respectively 0.17 [0.11, 0.23] and 0.13 [0.08, 0.18]). Interactions between overall forest area and last spring and last summer maximum temperature were selected and significant (respectively -0.1 [-0.02, -0.01] and 0.01 [0.01, 0.02]).

For generalist birds (Figure 2-C) the Shannon index on agricultural diversity was selected and significant (0.15 [0.12, 0.17]).

These results on all species primarily indicate that a land sparing scenario is better for forest and agricultural species and a land sharing scenario is better for generalist species.

We next examined the responses of species of conservation concern (Figure 3). For agriculture birds with conservation concern (Figure 3-A), the only land sharing/sparing variable selected was agriculture with a significant amount of nature and is significant (-0.11 [-0.18, 0.05]). For forest birds with conservation concern all three variables have been selected but only the area of forest 10 kilometers around was

Model component		Agricultural birds	Forest birds	Generalist birds
Spatial Random field	ρ (m)	17435.26	18336.21	13071.93
	σ	0.79	0.72	0.39
Temporal model	ϕ	0.89	0.60	0.89
	τ	111.48	367.41	359.25
Negative binomial dispersion		11.64	47.25	42.30

Table 2 Mean values of the posterior distribution of hyperparameters

significant (**0.5** [0.37,0.70]). These results on birds with conservation concern showed that land sharing is not the preferred solution in order to conserve these species.

Model component		Agricultural birds	Forest birds
Spatial Random field	ρ (m)	18480.75	36334.33
	σ	1.01	1.46
Temporal model	ϕ	0.92	0.92
	τ	129.18	24.74
Negative binomial dispersion		29.51	12.87

Table 3 Mean values of the posterior distribution of hyperparameters for birds with conservation concern

4 Discussion

In this study, we aimed to disentangle the impacts of land sharing and land sparing on French bird populations. Note that in a land sparing scenario, the goal is to divide landscape into a zone of high production agriculture and an area dedicated to biodiversity conservation whereas in a land sharing scenario, the landscape is shared between areas of food production and biodiversity conservation (with for example natural structures inside farmland areas).

First, our results showed that in the French bird community there is a diversity of response to the land sharing and land sparing gradient depending on habitat specialization (farmland, woodland, generalist) and conservation concern. Hence, land sharing seems to be beneficial for **generalist species**, as the abundance of this group increases if there is more diversity in agricultural practices. This result aligns with the main hypothesis because a land sharing scenario includes small and many kinds of habitats that offer a wide variety of nesting sites and resources that benefit species that do not have strict requirements in terms of resources such as habitat or food (Martin and Fahrig 2018). At the scale of our study (a 10 km buffer around 810 sites across France) land sparing seems to be the best scenario for all other species group with no conservation status. Indeed, for **forest species** our results showed that land sparing scenario is better, because the abundance of this group increases with a greater forest area, which aligns with our hypothesis. The variable “agriculture with a significant amount of nature” was also selected with a positive coefficient. This result can be explained by the study by (Monnet et al. 2026) that showed pesticides impact all




Fig. 2 Fixed effects of the best model for farmland species (A), woodland species (B) and generalist species (C). The variables highlighted in red are the land sharing and land sparing variables, the ones in green are about the land use of the squares, the one in yellow represent climate variables and the orange one are interactions between land sharing and sparing variables and climate. Significant variables are represented in plain color and with a star whereas non-significant variables are transparent.




Fig. 3 Fixed effects of the best model for farmland species with conservation concern (A) and woodland species with conservation concern(B). The variables highlighted in red are the land sharing and land sparing variables, the ones in green are about the land use of the squares, the one in yellow represent climate variables and the orange one are interactions between land sharing and sparing variables and climate. Significant variables are represented in plain color and with a star whereas non-significant variables are transparent.

birds and the fewer constraint forest species face, the better they perform. Regarding **farmland species**, unexpectedly, our results showed that a land sparing scenario is also preferable. This may be explained by the fact that in Europe we are already mostly in a land sharing scenario (Karner et al. 2019) and that being further on the land sharing spectrum would imply a habitat that is not the one they are specialized in.

599 Our results showed that the land sparing scenario is better for **forest species**
600 **with conservation concern** because the abundance increases with a greater forest
601 area.

602 For **agricultural species with conservation concern** it may appear counterin-
603 tuitive that their highest abundances are observed in land sparing scenarios, *i.e.*, areas
604 with a low proportion of natural habitats such as bushes and hedges. This pattern
605 might suggest that farmland birds of conservation concern do not benefit from these
606 semi-natural elements. However, this interpretation is likely misleading. We argue that
607 ground-nesting farmland bird species, which are now the primary focus of conserva-
608 tion efforts, have become particularly vulnerable in intensive agricultural landscapes.
609 This vulnerability stems from decades of agricultural mechanization and intensifica-
610 tion. Our analysis of land use within survey squares further supports this hypothesis
611 because birds with conservation concern are more abundant in farmland and open
612 landscapes. Hence, as shown in recent literature (Monnet et al. 2026; Rigal et al. 2023),
613 farmland birds are declining because of the intensification of agricultural practices and
614 pesticide use. Consequently, for conservation of these species the agricultural policies
615 should encourage a scenario with large open fields but with a low-yield agriculture
616 (Perrot et al. 2025).

617 In order to answer the question of how to allocate land use to conserve biodiver-
618 sity our results showed that there is not one agricultural practice to prefer to conserve
619 all groups of birds. Indeed, land sharing is better for generalist species whereas land
620 sparing is better for forest and agricultural species. However, as shown in (Rigal et al.
621 2023) farmland birds are declining; therefore, even if land sharing is not the primary
622 solution, we still consider a more biodiversity friendly agriculture to conserve bird
623 populations. As shown in (Monnet et al. 2026) pesticides are one of the drivers of this
624 decline, so reducing their use could therefore be a key solution. Overall, we cannot
625 conclude that farmland birds with conservation concern prefer land sparing. Despite
626 their apparent specialization in intensive agricultural landscapes, our findings suggest
627 that conserving these species should prioritize reducing chemical inputs and adopt-
628 ing wildlife-friendly practices within larger field structures, rather than intensifying
629 management further.

630 A main limitation of our analyses is the long time series of the STOC database.
631 Indeed, over this timescale there are no data on pesticides use, as the ones used in
632 (Monnet et al. 2026) only start in 2015 and the ones used in (Perrot et al. 2025) start
633 in 2008. There are also no data available on small natural structures inside (hedgerow
634 monitoring started in 2022 (Preux 2025)) fields. The indicators we used are thus,
635 the most appropriate variables available for this study period to account for land
636 use heterogeneity (with the Shannon index on agriculture practices), the presence of
637 natural structures in fields (agriculture with a significant amount of nature), as well
638 as big conserved forest. These three variables summarize well different characteristics
639 Then, we did not take into account detection bias, but we aggregated observations at
640 the survey-square scale to reduce randomness of detectability bias (Nabias et al. 2024).

641 In conclusion, our findings underscore a widespread challenge in agricultural land-
642 scapes akin to those in France: striking a balance between biodiversity conservation
643
644

and food security, particularly for species of conservation concern that depend on habitats shaped by intensive farming.

References

- Barbet-Massin M, Thuiller W, Jiguet F (2012) The fate of European breeding birds under climate, land-use and dispersal scenarios. *Global Change Biology* 18(3):881–890. <https://doi.org/10.1111/j.1365-2486.2011.02552.x>
- Barlow JC, Leckie SN, Pyle P, et al (2020) Eurasian Tree Sparrow (*Passer montanus*). In: Billerman SM, Keeney BK, Rodewald PG, et al (eds) *Birds of the World*. Cornell Lab of Ornithology, <https://doi.org/10.2173/bow.eutspa.01>
- Bowler DE, Heldbjerg H, Fox AD, et al (2019) Long-term declines of European insectivorous bird populations and potential causes. *Conservation Biology* 33(5):1120–1130. <https://doi.org/10.1111/cobi.13307>
- Burnham KP, Anderson DR (eds) (2004) *Model Selection and Multimodel Inference*. Springer, New York, NY, <https://doi.org/10.1007/b97636>
- Burns F, Eaton MA, Burfield IJ, et al (2021) Abundance decline in the avifauna of the European Union reveals cross-continental similarities in biodiversity change. *Ecology and Evolution* 11(23):16647–16660. <https://doi.org/10.1002/ece3.8282>
- Canterbury GE, Martin TE, Petit DR, et al (2000) Bird Communities and Habitat as Ecological Indicators of Forest Condition in Regional Monitoring. *Conservation Biology* 14(2):544–558. <https://doi.org/10.1046/j.1523-1739.2000.98235.x>
- Carson S, Mills Flemming J (2014) Seal encounters at sea: A contemporary spatial approach using R-INLA. *Ecological Modelling* 291:175–181. <https://doi.org/10.1016/j.ecolmodel.2014.07.022>
- Dambly LI, Isaac NJB, Jones KE, et al (2023) Integrated species distribution models fitted in INLA are sensitive to mesh parameterisation. *Ecography* 2023(7):e06391. <https://doi.org/10.1111/ecog.06391>
- Devictor V, Clavel J, Julliard R, et al (2010) Defining and measuring ecological specialization. *Journal of Applied Ecology* 47(1):15–25. <https://doi.org/10.1111/j.1365-2664.2009.01744.x>
- Donald PF, Green RE, Heath MF (2001) Agricultural intensification and the collapse of Europe’s farmland bird populations. *Proceedings of the Royal Society of London Series B: Biological Sciences* 268(1462):25–29. <https://doi.org/10.1098/rspb.2000.1325>
- Edwards DP, Gilroy JJ, Thomas GH, et al (2015) Land-Sparing Agriculture Best Protects Avian Phylogenetic Diversity. *Current Biology* 25(18):2384–2391. <https://doi.org/10.1016/j.cub.2015.08.022>

691 [//doi.org/10.1016/j.cub.2015.07.063](https://doi.org/10.1016/j.cub.2015.07.063)
692
693 European Environment Agency (2019) CORINE Land Cover 2018 (raster 100
694 m), Europe, 6-yearly - version 2020_20u1, May 2020. [https://doi.org/10.2909/](https://doi.org/10.2909/960998C1-1870-4E82-8051-6485205EBBAC)
695 [960998C1-1870-4E82-8051-6485205EBBAC](https://doi.org/10.2909/960998C1-1870-4E82-8051-6485205EBBAC)
696
697 Fahrig L (2013) Rethinking patch size and isolation effects: the habitat amount
698 hypothesis. *Journal of Biogeography* 40(9):1649–1663. [https://doi.org/10.1111/jbi.](https://doi.org/10.1111/jbi.12130)
699 [12130](https://doi.org/10.1111/jbi.12130)
700
701 Fahrig L (2020) Why do several small patches hold more species than few large
702 patches? *Global Ecology and Biogeography* 29(4):615–628. [https://doi.org/10.1111/](https://doi.org/10.1111/geb.13059)
703 [geb.13059](https://doi.org/10.1111/geb.13059)
704
705 Finch T, Gillings S, Green RE, et al (2019) Bird conservation and the land
706 sharing-sparing continuum in farmland-dominated landscapes of lowland England.
707 *Conservation Biology* 33(5):1045–1055
708
709 Fischer J, Abson DJ, Butsic V, et al (2014) Land Sparing Versus Land Sharing: Moving
710 Forward. *Conservation Letters* 7(3):149–157. <https://doi.org/10.1111/conl.12084>
711
712 Fontaine B, Moussy C, Carricaburu JC, et al (2021) Suivi des oiseaux communs en
713 France 1989-2019 : 30 ans de suivis participatifs. Pages: 45 pages
714
715 Genty L, Meynard CN, Bopp MC, et al (2026) Intensive management negatively
716 impacts field margin ecosystem service indicators at both field and landscape levels.
717 *Ecological Applications* 36(1):e70161. <https://doi.org/10.1002/eap.70161>
718
719 Grass I, Loos J, Baensch S, et al (2019) Land-sharing/-sparing connectivity landscapes
720 for ecosystem services and biodiversity conservation. *People and Nature* 1(2):262–
721 272. <https://doi.org/10.1002/pan3.21>
722
723 Gregory RD, Noble D, Field R, et al (2003) Using birds as indicators of biodiversity.
724 *Ornis Hungarica*
725
726 Harris I, Osborn TJ, Jones P, et al (2020) Version 4 of the CRU TS monthly high-
727 resolution gridded multivariate climate dataset. *Scientific Data* 7(1):109. [https://](https://doi.org/10.1038/s41597-020-0453-3)
728 doi.org/10.1038/s41597-020-0453-3
729
730 Heldbjerg H, Sunde P, Fox AD (2018) Continuous population declines for specialist
731 farmland birds 1987-2014 in Denmark indicates no halt in biodiversity loss in agri-
732 cultural habitats. *Bird Conservation International* 28(2):278–292. [https://doi.org/](https://doi.org/10.1017/S0959270916000654)
733 [10.1017/S0959270916000654](https://doi.org/10.1017/S0959270916000654)
734
735 Hume R, Kirwan GM (2020) Eurasian Thick-knee (*Burhinus oedicnemus*). In: Biller-
736 man SM, Keeney BK, Rodewald PG, et al (eds) *Birds of the World*. Cornell Lab of
Ornithology, <https://doi.org/10.2173/bow.eutkne1.01>

Illian JB, Martino S, Sørbye SH, et al (2013) Fitting complex ecological point process models with integrated nested Laplace approximation. <i>Methods in Ecology and Evolution</i> 4(4):305–315. https://doi.org/10.1111/2041-210x.12017	737 738 739 740
Inger R, Gregory R, Duffy JP, et al (2015) Common European birds are declining rapidly while less abundant species' numbers are rising. <i>Ecology Letters</i> 18(1):28–36. https://doi.org/10.1111/ele.12387	741 742 743 744
Jiguet F, Devictor V, Julliard R, et al (2012) French citizens monitoring ordinary birds provide tools for conservation and ecological sciences. <i>Acta Oecologica</i> 44:58–66. https://doi.org/10.1016/j.actao.2011.05.003	745 746 747 748
Julliard R, Clavel J, Devictor V, et al (2006) Spatial segregation of specialists and generalists in bird communities. <i>Ecology Letters</i> 9(11):1237–1244. https://doi.org/10.1111/j.1461-0248.2006.00977.x	749 750 751 752
Karen Aghababayan (2024) Birds as potential bioindicators for terrestrial ecosystems. <i>International Journal of Life Science Research Archive</i> 6(1):001–022. https://doi.org/10.53771/ijlsra.2024.6.1.0112	753 754 755 756
Karner K, Cord AF, Hagemann N, et al (2019) Developing stakeholder-driven scenarios on land sharing and land sparing – Insights from five European case studies. <i>Journal of Environmental Management</i> 241:488–500. https://doi.org/10.1016/j.jenvman.2019.03.050	757 758 759 760
Krainski E, Gómez-Rubio V, Bakka H, et al (2018) <i>Advanced Spatial Modeling with Stochastic Partial Differential Equations Using R and INLA</i> , 0th edn. Chapman and Hall/CRC, https://doi.org/10.1201/9780429031892	761 762 763 764
Lamb A, Finch T, Pearce-Higgins JW, et al (2019) The consequences of land sparing for birds in the United Kingdom. <i>Journal of Applied Ecology</i> 56(8):1870–1881. https://doi.org/10.1111/1365-2664.13362	765 766 767 768
Lindgren F, Rue H, Lindström J, et al (2011) An explicit link between Gaussian fields and Gaussian Markov random fields: the stochastic partial differential equation approach [with Discussion]. <i>Journal of the Royal Statistical Society Series B (Statistical Methodology)</i> 73(4):423–498	769 770 771 772
Martin AE, Fahrig L (2018) Habitat specialist birds disperse farther and are more migratory than habitat generalist birds. <i>Ecology</i> 99(9):2058–2066	773 774 775 776
Monnet AC, Cairo M, Deguines N, et al (2026) Common birds have higher abundances in croplands with lower pesticide purchases. <i>Proceedings of the Royal Society B: Biological Sciences</i> 293(2062):20252370. https://doi.org/10.1098/rspb.2025.2370	777 778 779 780
Nabias J, Barbaro L, Fontaine B, et al (2024) Reassessment of French breeding bird population sizes using citizen science and accounting for species detectability. <i>PeerJ</i>	781 782

12:e17889. <https://doi.org/10.7717/peerj.17889>

Nagy TMG, Soós A (2025) Land Sharing and Land Sparing in Agriculture – Can We Change Our Attitudes? *EuroChoices* 24(1):43–49. <https://doi.org/10.1111/1746-692X.12459>

Padoa-Schioppa E, Baietto M, Massa R, et al (2006) Bird communities as bioindicators: The focal species concept in agricultural landscapes. *Ecological Indicators* 6(1):83–93. <https://doi.org/10.1016/j.ecolind.2005.08.006>

Perrot T, Wolfram J, Schulz R, et al (2025) Temporal trends in pesticide pressure differ among non-target species groups but show spatially consistent patterns across France. *Journal of Hazardous Materials* 497:139695. <https://doi.org/10.1016/j.jhazmat.2025.139695>

Phalan B, Onial M, Balmford A, et al (2011) Reconciling Food Production and Biodiversity Conservation: Land Sharing and Land Sparing Compared. *Science* 333(6047):1289–1291. <https://doi.org/10.1126/science.1208742>

Preux T (2025) Cartographie de la densité de haies en France métropolitaine en 2022. <https://doi.org/10.34847/NKL.1726BU27>

Rigal S, Dakos V, Alonso H, et al (2023) Farmland practices are driving bird population decline across Europe. *Proceedings of the National Academy of Sciences* 120(21):e2216573120. <https://doi.org/10.1073/pnas.2216573120>

Righetto AJ, Faes C, Vandendijck Y, et al (2020) On the choice of the mesh for the analysis of geostatistical data using R-INLA. *Communications in Statistics - Theory and Methods* 49(1):203–220. <https://doi.org/10.1080/03610926.2018.1536209>

Riva F, Fahrig L (2022) The disproportionately high value of small patches for biodiversity conservation. *Conservation Letters* 15(3):e12881. <https://doi.org/10.1111/conl.12881>

Rue H, Martino S, Chopin N (2009) Approximate Bayesian Inference for Latent Gaussian Models by Using Integrated Nested Laplace Approximations. *Journal of the Royal Statistical Society Series B (Statistical Methodology)* 71(2):319–392

Shannon CE (1948) A mathematical theory of communication. *The Bell System Technical Journal* 27(3):379–423. <https://doi.org/10.1002/j.1538-7305.1948.tb01338.x>

Simpson D, Rue H, Riebler A, et al (2017) Penalising Model Component Complexity: A Principled, Practical Approach to Constructing Priors. *Statistical Science* 32(1):1–28

Thepault A, Adenot T, Lorrilliere R, et al (2025) Warm or Dry Springs (Still) Boost the Reproduction of Most Temperate Songbirds. *Global Ecology and Biogeography*

34(10):e70138. https://doi.org/10.1111/geb.70138	829
UICN France, MNHN, LPO, et al (2016) La liste rouge des espèces menacées en France	830
- Chapitre des Oiseaux de France métropolitaine. Paris, France	831
	832
	833
	834
Declarations	835
	836
Funding	837
This work was supported by the Chaire “Modélisation Mathématique et Biodiver-	838
sité” of Veolia-École polytechnique-Muséum national d’Histoire naturelle-Fondation	839
X, through a Ph.D. scholarship.	840
	841
	842
Competing interest	843
The authors have no relevant financial or non-financial interests to disclose.	844
	845
	846
Author contributions	847
All authors contributed to the study conception and design. Material preparation and	848
(online) data collection was performed by Adélie Erard. Data analysis was performed	849
by Adélie Erard and Romain Lorrillière. The first draft of the manuscript was written	850
by Adélie Erard and all authors commented on previous versions of the manuscript.	851
All authors read and approved the final manuscript.	852
	853
Data availability	854
The bird abundance data analyzed during the current study are available on reasonable	855
request on Vigie-Nature website . Climate data are available on WordClim website .	856
Land use data are available on Copernicus Website .	857
	858
	859
Code availability	860
R code used for analysis is available at https://github.com/adelleerard/birds_inla .	861
	862
	863
	864
	865
	866
	867
	868
	869
	870
	871
	872
	873
	874

Land Sharing or Land Sparing? Implications for Farmland Birds of Conservation Concern in France.

Supplementary material

Landscape Ecology

Adélie Erard^{1,*}, Ottmar Cronie^{2,*}, Raphaël Lachière-Rey^{3,1}, and Romain Lorrillière^{4,5}

¹Université Paris Cité, CNRS, MAP5, F-75006, Paris, France

² Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg, Gothenburg, SE-412 96, Sweden

³Inria Paris, MathNet, Paris, F-75013, France

⁴MNHN, CRBPO, F-75005, Paris, France

⁵MNHN, Sorbonne Université, CNRS, Centre d'Écologie et des Sciences de la Conservation, CESCO, F-75005, Paris, France

*Corresponding authors: aerard@math.cnrs.fr and ottmar.cronie@gu.se

March 5, 2026

List of Figures

SI-1	Observed abundance of generalist species, by year	3
SI-2	Abundance of forest species observed in the STOC program, by year	4
SI-3	Observed abundance of agricultural species, by year	5
SI-4	Observed abundance of forest species with conservation concern, by year	6
SI-5	Observed abundance of agricultural species with conservation concern, by year	7
SI-6	Mesh choice. The observed points are represented in blue.	8
SI-7	Results of the PCA on the land cover of each square	9

23 **List of Tables**

24 SI-1 Results for agricultural species 10

25 SI-2 Results for forest species 11

26 SI-3 Results for generalist species 12

27 SI-4 Results for agricultural species with conservation concern 13

28 SI-5 Results for forest species with conservation concern 14

29 SI-6 Bird species by taxonomic order, habitat type and conservation status 15

30 SI-7 Corine Land Cover (CLC) Level 3 classification codes and class names 18

31 **1 Maps**




Figure SI-1: Observed abundance of generalist species, by year




Figure SI-2: Abundance of forest species observed in the STOC program, by year




Figure SI-3: Observed abundance of agricultural species, by year




Figure SI-4: Observed abundance of forest species with conservation concern, by year




Figure SI-5: Observed abundance of agricultural species with conservation concern, by year




Figure SI-6: Mesh choice. The observed points are represented in blue.

2 PCA on land use of the square




Figure SI-7: Results of the PCA on the land cover of each square

3 Tables of the results

	mean	sd	0.025quant	0.5quant	0.975quant	mode	kld
(Intercept)	1.45	0.05	1.36	1.45	1.55	1.45	0.00
Latitude	0.06	0.04	-0.01	0.06	0.13	0.06	0.00
Last spring max temperature	0.02	0.01	-0.00	0.02	0.04	0.02	0.00
Last summer max temperature	0.01	0.01	-0.01	0.01	0.03	0.01	0.00
PCA1	-0.52	0.03	-0.58	-0.52	-0.46	-0.52	0.00
PCA3	-0.23	0.02	-0.27	-0.23	-0.18	-0.23	0.00
Forest 10km around	-0.03	0.04	-0.11	-0.03	0.04	-0.03	0.00
Agriculture with signif nature	-0.05	0.03	-0.11	-0.05	0.01	-0.05	0.00
Last spring max temperature×Forest 10km around	0.01	0.01	-0.00	0.01	0.03	0.01	0.00
Last summer max temperature×Agriculture with signif nature	0.01	0.01	-0.00	0.01	0.02	0.01	0.00
Latitude×Last spring max temperature	0.02	0.01	0.01	0.02	0.03	0.02	0.00
	mean	sd	0.025quant	0.5quant	0.975quant	mode	
(1/overdispersion)	11.64	0.28	11.09	11.64	12.21	11.64	
Range for spatial (m)	17435.26	1365.42	14914.11	17377.90	20286.97	17255.39	
Stdev for spatial	0.79	0.02	0.74	0.79	0.84	0.79	
Precision for year	111.48	72.08	20.67	94.76	290.94	58.91	
Rho for year	0.89	0.08	0.70	0.91	0.98	0.95	

Table SI-1: Results for agricultural species

	mean	sd	0.025quant	0.5quant	0.975quant	mode	kld
(Intercept)	1.70	0.02	1.66	1.70	1.75	1.70	0.00
PCA1	0.44	0.02	0.40	0.44	0.49	0.44	0.00
PCA2	-0.07	0.02	-0.10	-0.07	-0.04	-0.07	0.00
Agriculture with signif nature	0.13	0.02	0.08	0.13	0.18	0.13	0.00
Latitude	0.23	0.03	0.17	0.23	0.30	0.23	0.00
Forest 10km around	0.17	0.03	0.11	0.17	0.23	0.17	0.00
Last spring max temperature	-0.00	0.01	-0.02	-0.00	0.02	-0.00	0.00
Last summer max temperature	-0.01	0.01	-0.03	-0.01	0.01	-0.01	0.00
Last spring min temperature	0.03	0.01	0.01	0.03	0.04	0.03	0.00
Forest 10km around×Last spring max temperature	-0.01	0.00	-0.02	-0.01	-0.01	-0.01	0.00
Forest 10km around×Last summer max temperature	0.01	0.00	0.01	0.01	0.02	0.01	0.00
	mean	sd	0.025quant	0.5quant	0.975quant	mode	
(1/overdispersion)	47.25	1.71	43.99	47.21	50.71	47.13	
Range for spatial (m)	18336.21	1369.44	15796.14	18282.59	21185.06	18170.46	
Stddev for spatial	0.72	0.02	0.68	0.72	0.76	0.72	
Precision for year	367.41	156.77	132.08	343.02	735.57	294.44	
Rho for year	0.60	0.16	0.26	0.62	0.86	0.65	

Table SI-2: Results for forest species

	mean	sd	0.025quant	0.5quant	0.975quant	mode	kld
(Intercept)	2.25	0.03	2.19	2.25	2.30	2.26	0.00
PCA1	0.03	0.01	0.01	0.03	0.06	0.03	0.00
Shanon index on agricultural patches	0.15	0.01	0.12	0.15	0.17	0.15	0.00
Latitude	0.05	0.01	0.02	0.05	0.08	0.05	0.00
PCA3	0.03	0.01	0.01	0.03	0.05	0.03	0.00
	mean	sd	0.025quant	0.5quant	0.975quant	mode	
(1/overdispersion)	42.30	0.99	40.40	42.29	44.30	42.24	
Range for spatial (m)	13071.93	1065.66	11080.66	13035.25	15273.40	12975.89	
Stddev for spatial	0.39	0.01	0.36	0.39	0.42	0.39	
Precision for year	359.25	243.30	67.33	300.15	980.45	185.71	
Rho for year	0.89	0.08	0.69	0.91	0.98	0.95	

Table SI-3: Results for generalist species

	mean	sd	0.025quant	0.5quant	0.975quant	mode	kld
(Intercept)	1.10	0.05	1.00	1.10	1.20	1.10	0.00
PCA3	-0.30	0.02	-0.35	-0.30	-0.26	-0.30	0.00
Last spring total rain	-0.02	0.01	-0.04	-0.02	-0.01	-0.02	0.00
Agriculture with signif nature	-0.11	0.03	-0.18	-0.11	-0.05	-0.11	0.00
Last spring max temperature	-0.01	0.01	-0.03	-0.01	0.02	-0.01	0.00
PCA1	-0.62	0.03	-0.68	-0.62	-0.55	-0.62	0.00
Agriculture with signif nature×Last spring max temperature	-0.02	0.01	-0.03	-0.02	-0.00	-0.02	0.00
	mean	sd	0.025quant	0.5quant	0.975quant	mode	
(1/overdispersion)	29.51	1.37	26.92	29.47	32.31	29.38	
Range for spatial (m)	18480.75	1459.90	15740.59	18434.65	21484.56	18366.57	
Stddev for spatial	1.01	0.03	0.95	1.01	1.07	1.01	
Precision for year	129.18	76.50	25.76	113.24	312.73	74.57	
Rho for year	0.92	0.05	0.80	0.93	0.99	0.96	

Table SI-4: Results for agricultural species with conservation concern

	mean	sd	0.025quant	0.5quant	0.975quant	mode	kld
(Intercept)	-0.48	0.12	-0.72	-0.48	-0.25	-0.48	0.00
Latitude	0.28	0.10	0.08	0.28	0.48	0.28	0.00
Last summer min temperature	-0.05	0.03	-0.12	-0.05	0.01	-0.05	0.00
Last spring max temperature	-0.08	0.03	-0.14	-0.08	-0.03	-0.08	0.00
Last summer max temperature	0.05	0.02	-0.00	0.05	0.09	0.05	0.00
PCA1	0.78	0.05	0.67	0.78	0.88	0.78	0.00
PCA2	0.08	0.04	0.01	0.08	0.16	0.08	0.00
Shanon index on agricultural patches	-0.11	0.08	-0.26	-0.11	0.04	-0.11	0.00
Forest 10km around	0.54	0.08	0.37	0.54	0.70	0.54	0.00
Agriculture with signif nature	0.14	0.07	-0.00	0.14	0.28	0.14	0.00
Last summer max temperature ×Shanon index on agricultural patches	0.02	0.01	-0.01	0.02	0.04	0.02	0.00
Last spring max temperature ×Forest 10km around	0.02	0.02	-0.01	0.02	0.06	0.02	0.00
Latitude × Last spring max temperature	-0.05	0.02	-0.08	-0.05	-0.02	-0.05	0.00
	mean	sd	0.025quant	0.5quant	0.975quant	mode	
(1/overdispersion)	12.87	1.47	10.24	12.78	16.00	12.60	
Range for spatial (m)	36334.33	3889.39	29280.72	36127.94	44576.32	35719.07	
Stdev for spatial	1.46	0.07	1.33	1.46	1.60	1.46	
Precision for year	24.74	15.83	5.49	21.02	65.21	14.18	
Rho for year	0.92	0.06	0.76	0.93	0.98	0.96	

Table SI-5: Results for forest species with conservation concern

34 **4 List of species with their habitat and conservation status**

Table SI-6: Bird species by taxonomic order, habitat type and conservation status

Order	Species	Habitat type	Status
Accipitriformes	<i>Buteo buteo</i>	Agricultural	Least concerned
Passeriformes	<i>Emberiza cirrus</i>	Agricultural	Least concerned
Falconiformes	<i>Falco tinnunculus</i>	Agricultural	Conservation stake
Passeriformes	<i>Lanius collurio</i>	Agricultural	Conservation stake
Passeriformes	<i>Emberiza citrinella</i>	Agricultural	Conservation stake
Passeriformes	<i>Corvus frugilegus</i>	Agricultural	Least concerned
Passeriformes	<i>Sylvia communis</i>	Agricultural	Conservation stake
Charadriiformes	<i>Vanellus vanellus</i>	Agricultural	Conservation stake
Galliformes	<i>Coturnix coturnix</i>	Agricultural	Least concerned
Galliformes	<i>Alectoris rufa</i>	Agricultural	Least concerned
Upupiformes	<i>Upupa epops</i>	Agricultural	Least concerned
Passeriformes	<i>Saxicola rubicola</i>	Agricultural	Conservation stake
Passeriformes	<i>Passer montanus</i>	Agricultural	Conservation stake
Charadriiformes	<i>Burhinus oedicnemus</i>	Agricultural	Conservation stake
Passeriformes	<i>Lullula arborea</i>	Agricultural	Least concerned
Passeriformes	<i>Alauda arvensis</i>	Agricultural	Conservation stake
Passeriformes	<i>Emberiza calandra</i>	Agricultural	Conservation stake
Passeriformes	<i>Carduelis cannabina</i>	Agricultural	Least concerned
Galliformes	<i>Perdix perdix</i>	Agricultural	Least concerned
Passeriformes	<i>Motacilla flava</i>	Agricultural	Least concerned
Passeriformes	<i>Anthus pratensis</i>	Agricultural	Conservation stake
Passeriformes	<i>Phylloscopus collybita</i>	Forest	Least concerned
Passeriformes	<i>Turdus philomelos</i>	Forest	Least concerned
Piciformes	<i>Dendrocopos major</i>	Forest	Least concerned
Passeriformes	<i>Erithacus rubecula</i>	Forest	Least concerned
Passeriformes	<i>Regulus ignicapilla</i>	Forest	Least concerned
Passeriformes	<i>Troglodytes troglodytes</i>	Forest	Least concerned
Passeriformes	<i>Certhia familiaris</i>	Forest	Least concerned

Order	Species	Habitat type	Status
Passeriformes	<i>Regulus regulus</i>	Forest	Conservation stake
Passeriformes	<i>Sitta europaea</i>	Forest	Least concerned
Passeriformes	<i>Lophophanes cristatus</i>	Forest	Least concerned
Passeriformes	<i>Phylloscopus bonelli</i>	Forest	Least concerned
Passeriformes	<i>Poecile palustris</i>	Forest	Least concerned
Passeriformes	<i>Certhia brachydactyla</i>	Forest	Least concerned
Passeriformes	<i>Turdus viscivorus</i>	Forest	Least concerned
Passeriformes	<i>Phylloscopus sibilatrix</i>	Forest	Conservation stake
Passeriformes	<i>Phylloscopus trochilus</i>	Forest	Conservation stake
Passeriformes	<i>Pyrrhula pyrrhula</i>	Forest	Conservation stake
Passeriformes	<i>Coccothraustes coccothraustes</i>	Forest	Least concerned
Piciformes	<i>Dendrocopos medius</i>	Forest	Least concerned
Passeriformes	<i>Parus montanus</i>	Forest	Least concerned
Passeriformes	<i>Sylvia melanocephala</i>	Forest	Conservation stake
Columbiformes	<i>Columba palumbus</i>	Generalist	Least concerned
Passeriformes	<i>Corvus corone</i>	Generalist	Least concerned
Passeriformes	<i>Cyanistes caeruleus</i>	Generalist	Least concerned
Passeriformes	<i>Sylvia atricapilla</i>	Generalist	Least concerned
Passeriformes	<i>Turdus merula</i>	Generalist	Least concerned
Passeriformes	<i>Fringilla coelebs</i>	Generalist	Least concerned
Passeriformes	<i>Hippolais polyglotta</i>	Generalist	Least concerned
Passeriformes	<i>Parus major</i>	Generalist	Least concerned
Passeriformes	<i>Garrulus glandarius</i>	Generalist	Least concerned
Cuculiformes	<i>Cuculus canorus</i>	Generalist	Least concerned
Passeriformes	<i>Oriolus oriolus</i>	Generalist	Least concerned
Piciformes	<i>Picus viridis</i>	Generalist	Least concerned
Passeriformes	<i>Luscinia megarhynchos</i>	Generalist	Least concerned
Passeriformes	<i>Prunella modularis</i>	Generalist	Least concerned

5 Corine Land cover Codes

CLC Code	Class name
111	Continuous urban fabric
112	Discontinuous urban fabric
121	Industrial or commercial units
122	Road and rail networks and associated land
123	Port areas
124	Airports
131	Mineral extraction sites
132	Dump sites
133	Construction sites
141	Green urban areas
142	Sport and leisure facilities
211	Non-irrigated arable land
212	Permanently irrigated land
213	Rice fields
221	Vineyards
222	Fruit trees and berry plantations
223	Olive groves
231	Pastures
241	Annual crops associated with permanent crops
242	Complex cultivation patterns
243	Land principally occupied by agriculture, with significant areas of natural vegetation
244	Agroforestry areas
311	Broad-leaved forest
312	Coniferous forest
313	Mixed forest
321	Natural grasslands
322	Moors and heath land
323	Sclerophyllous vegetation
324	Transitional woodland-shrub
331	Beaches, dunes, sands

332	Bare rocks
333	Sparsely vegetated areas
334	Burnt areas
335	Glaciers and perpetual snow
411	Inland marshes
412	Peat bogs
421	Salt marshes
422	Salines
423	Intertidal flats
511	Water courses
512	Water bodies
521	Coastal lagoons
522	Estuaries
523	Sea and ocean

Table SI-7: Corine Land Cover (CLC) Level 3 classification codes and class names