

Original Articles

Recognition of stream drying based on benthic macroinvertebrates: A new tool in Central Europe



Michal Straka^{a,b,*}, Marek Polášek^{a,b}, Vít Syrovátka^b, Rachel Stubbington^c, Světlana Zahradková^b, Denisa Němejcová^a, Lenka Šikulová^b, Pavla Řezníčková^b, Libuše Opatřilová^{a,d}, Thibault Datry^e, Petr Pařil^b

^a T. G. Masaryk Water Research Institute, p. r. i., Brno Branch Office, Mojmírovo náměstí 16, 612 00 Brno, Czech Republic

^b Department of Botany and Zoology, Faculty of Science, Masaryk University, Kotlářská 2, 611 37 Brno, Czech Republic

^c School of Science and Technology, Nottingham Trent University, Nottingham NG11 8NS, UK

^d Povodí Vltavy, State Enterprise, Holečkova 3178/8, 150 00, Praha 5, Czech Republic

^e IRSTEA, UR RIVERLY, Centre de Lyon-Villeurbanne, France

ARTICLE INFO

Keywords:

Intermittent
Flow regime
Index
Invertebrates
Indicators

ABSTRACT

Many streams in the extensive Central European region have an intermittent flow regime. Conventional hydrological methods used to identify zero-flow conditions, and in particular drying events, have limited use when assessing large areas dominated by low-order streams. We developed a novel multimetric index to recognise antecedent stream drying based on the analysis of benthic macroinvertebrate communities. The data used to develop the index were collected in pristine streams with different flow permanence regimes between 2012 and 2014, using standard sampling methods for ecological status assessment. The data include 64 perennial, 19 near-perennial and 27 intermittent benthic macroinvertebrate samples. Metrics considered for the index included variables based on (i) the occurrence of indicator taxa, (ii) the proportion of biological and ecological traits, and (iii) structural community metrics. Linear discriminant analysis identified the metric combinations that best discriminated among the three flow permanence categories. Different metrics were used in the final multimetric index calculation for the autumn and spring season that followed stream drying. In both seasons, the index included the proportion of indicator taxa and the proportion of taxa with high body flexibility. In addition, the autumn index included the proportion of taxa with a preference for organic substrates, whereas in spring the index included total abundance. Independent data from regulatory monitoring activity were used to validate the accuracy of the index. The correct classification of independent samples was 92% and 96% for samples from perennial and non-perennial sites, respectively. The index can be calculated using data collected by routine monitoring programmes used to assess ecological status and provides information about stream intermittence where conventional hydrological monitoring is limited. As intermittent streams increase in extent in global regions including Central Europe, the tool may be of particular interest to those who use invertebrates to monitor or manage these ecosystems.

1. Introduction

Intermittent rivers and ephemeral streams (i.e. streams with periodic flow cessation and/or drying; IRES) are mostly associated with arid and semi-arid regions. However, they are also common and widespread in temperate and continental regions with cooler, wetter climates (Datry et al., 2014a; Stubbington et al., 2017). IRES are typically managed using methods developed for perennial waterways, or as if they were part of the terrestrial ecosystems (Acuña et al., 2014;

Stubbington et al., 2018), and the need for more effective policies and management strategies has been highlighted around the world (Datry et al., 2017; Marshall et al., 2018). In the European Union, a major objective for water management is the assessment of ecological status in water bodies, to meet the legislative requirements of the EU Water Framework Directive (European Commission, 2000). The characterisation of 'biological quality elements' such as benthic macroinvertebrates is one important approach used to evaluate ecological status.

* Corresponding author at: T. G. Masaryk Water Research Institute, p. r. i., Brno Branch Office, Mojmírovo náměstí 16, 612 00 Brno, Czech Republic
E-mail address: michal.straka@centrum.cz (M. Straka).

<https://doi.org/10.1016/j.ecolind.2019.105486>

Received 14 March 2019; Received in revised form 31 May 2019; Accepted 13 June 2019

1470-160X/ © 2019 Elsevier Ltd. All rights reserved.

Aquatic macroinvertebrates are routinely used as indicator organisms for determining ecological status and diagnosing specific environmental pressures (e.g. Rosenberg and Resh, 1993; Birk et al., 2012). However, even short-term (i.e. days to weeks) stream drying can substantially alter benthic macroinvertebrate community composition (Datry et al., 2014b; Hille et al., 2014; Lancaster and Ledger, 2015). The interpretation of ecological status based on metrics developed in perennial streams can thus be misleading (Munné and Prat, 2009; Menció and Mas-Pla, 2010; Wilding et al., 2018). Information about flow permanence is crucial to accurately interpret data used to evaluate status and thus inform effective IRES management. In addition, knowledge about stream intermittence is needed, to anticipate the effects of climate change, to assess human pressures such as water abstraction and to manage protected species and habitats (Wilby et al., 2010).

Using flow gauging stations to monitor the flow permanence of small streams is expensive and technically impractical, because small streams (Strahler orders 1–4) comprise a substantial part of the stream network length (e.g. 92% in Czech Republic; Zahrádková et al., 2015). Moreover, even where hydrological data are collected, they may not distinguish between lentic and dry zero-flow conditions, and alternative methods are therefore needed to recognize IRES and characterize their water regimes (Gallart et al., 2016; Beaufort et al., 2018), for example using stream biota. Loss of surface water acts as an ecological filter (Poff 1997) and benthic macroinvertebrates have taxon-specific quantitative responses to drying (Datry et al., 2014b; Leigh and Datry, 2017). Dry phases can therefore be detected through both structural (i.e. taxonomic) and functional (i.e. trait-based) changes in community composition (Bogan et al., 2013; Schriever et al., 2015; Leigh et al., 2016; Chadd et al., 2017). Presence/absence of indicator taxa of benthic invertebrates has also been used to recognize stream flow duration (NC Division of Water Quality, 2010; Nadeau, 2015), and both taxonomic structure and species trait information have been used to assess flow connectivity in Mediterranean regions (Cid et al., 2016).

Such tools may enable the data routinely collected during biomonitoring programmes to provide information about flow intermittence, even if hydrological data are absent. However, no method to detect drying events has been developed for the extensive continental-climate region of Central Europe.

Different metrics can be sensitive to different aspects of stream drying, and a combination of structural, functional and indicator-taxa metrics may therefore provide robust and sensitive insight into the responses of an assemblage to drying (Li et al., 2010). Here, our aim was to develop and test a new multimetric index to identify antecedent stream drying events based on the structural and functional composition of benthic macroinvertebrate communities. We evaluated three groups of metrics as indicators of stream drying: (i) the occurrence of indicator taxa; (ii) the proportion of biological and ecological traits; and (iii) structural community metrics. We identified the metrics and metric combinations that most effectively indicated antecedent stream drying.

2. Materials and methods

2.1. Study area

The study was conducted in 23 small (order 2–4) streams in the Czech Republic (Fig. 1). All were classified as reference streams, i.e. exposed to minimal anthropogenic impacts. Catchment land use was dominated by woodland, with smaller proportions of agricultural land, and with at least 10-m riparian buffer strips of deciduous bushes and trees. The study area has a warm-summer humid continental climate (Köppen-Geiger classification *Dfb*; Peel et al., 2007) and spans the Hercynian (17 streams) and West-Carpathian (6 streams) biogeographical subprovinces (Culek, 2013), which approximately correspond with ecoregions 9 (Central Highlands) and 10 (Carpathians; Illies, 1967). To facilitate comparison of macroinvertebrate communities from different sites, streams were paired, with each pair comprising one




Fig. 1. Location of the study sites in the Czech Republic. Open squares represent non-perennial (i.e. near-perennial and intermittent) sites and filled squares represent perennial sites.

non-perennial and one perennial stream of comparable altitude, size, geology and hydromorphology. In one case, a single perennial stream was paired with two non-perennial streams. A sampling site representative of typical conditions was selected in each stream. The altitude of the sampling sites ranged from 250 m to 560 m a.s.l. and the maximum distance between paired sites was 13 km.

2.2. Sampling strategy

Benthic macroinvertebrate assemblages were sampled in spring (March–April) in 2013 and 2014 and in autumn (September–October) in 2012, 2013 and 2014. Proportional multi-habitat three-minute kick samples were collected with a hand net (25 × 25 cm aperture, 0.5-mm mesh size) according to the standard method used for ecological status assessments in the Czech Republic (Kokeš et al., 2006; Kokeš and Němejcová, 2006). Samples were preserved in 4% formaldehyde and processed in the laboratory. All macroinvertebrates were identified to the lowest possible taxonomic level: 292 out of 421 to species level, 117 to genus level, and the remaining 12 to a higher taxonomic level.

Sites were classified as non-perennial or perennial *a priori*, based on expert knowledge. In addition, observed instream conditions were recorded from summer 2012 to autumn 2014, including determination of the dry period duration using a water-level logger (Solinst Levellogger Edge) and photo-trap (Acorn 5310MG) installed at each site. Each water-level logger was placed in the lowest part of the streambed, to account for the persistence of surface water in isolated pools. Photo-traps were installed in trees in the adjacent riparian zone (at 3 m height) and were facing a water-level gauge board. Logger and photo-trap data were collected every 15 min and every 4 h, respectively.

We divided individual samples into three intermittence categories: *perennial*, *intermittent* and an intermediate *near-perennial* category. The allocation of samples into the three categories was based primarily on the dry period duration: i) perennial sites were defined as those which experienced more than one year of continuous flow prior to sample collection; ii) near-perennial sites lost surface water for hundreds of meters for < 7 d in the summer prior to sampling, with disconnected pools occasionally present; and iii) intermittent sites dried for > 1 km for 7–86 d before sampling. The cut-off between near-perennial and intermittent sites was arbitrary, reflecting the gradual nature of stream drying. In total, we collected 27 perennial, 7 near-perennial and 12 intermittent samples in spring and 34 perennial, 12 near-perennial and 15 intermittent samples in autumn. Spring samples from intermittent and near-perennial streams were taken 175–224 d after flow resumption and autumn samples were taken 2–63 d after flow resumption.

2.3. Identification of perennial flow indicators and flow intermittence tolerant taxa

To identify taxa associated with either perennial or intermittent streams, we performed Indicator Species Analysis (IndVal; Dufrêne and Legendre, 1997) using samples from spring 2013 and autumn 2012 and 2013. IndVal was performed using the `multipatt` function in the R package `indicspecies` (De Cáceres and Legendre, 2009), which identifies taxa with a strong association with each group based on their abundance and frequency. The calculated indicator value (IV) of a given taxon for its preferred group ranges from zero (no affinity) to one (strongest affinity). Best-matching patterns were tested for statistical significance of the associations. Taxa significantly associated with either perennial or intermittent and near-perennial samples were considered as reliable indicators and were assigned a reliability value of 2 (see “Proportion of indicator taxa” section). Taxa with non-significant IV but found exclusively in either perennial or intermittent and near-

perennial streams in at least 10% of samples were considered as indicators with lower reliability and were assigned a reliability value of 1.

To support the IndVal analyses, we conducted a literature review of primary research papers comparing macroinvertebrate assemblages at perennial and non-perennial sites during flowing phases. From these studies, a taxon was identified as indicative of perennial flow if it was absent from intermittent stream assemblages. A taxon was considered as intermittence tolerant if it was found in intermittent stream assemblages. Literature sources describing the autecology of macroinvertebrate taxa were also reviewed to explore taxon-specific responses to intermittence. If a taxon's relationship with stream intermittence was described once in data-based published sources then the species was assigned a reliability value of 1, and if it was published two or more times, a value of 2 was assigned. The review focused on taxa living in the Czech Republic area; consulted literature is listed in Appendix A. If information in different published sources were contradictory or unclear, then a taxonomic expert decided the reliability of the taxa based on the available evidence.

2.4. Metrics used for index development

Three groups of potential metrics that indicate flow intermittence were calculated, to inform subsequent development of the multimetric index:

i) Proportion of indicator taxa

The proportion of indicator taxa was calculated as the sum of the reliability values of perennial flow indicators present in a sample divided by the sum of the reliability values of all (perennial + intermittence-tolerant) indicators present. The proportion of indicators was rescaled to zero and the resultant values are within the range −1 to +1.

Proportion of indicator taxa

$$= \left(\frac{\sum \text{values of perennial flow indicators}}{\sum \text{values of all indicators}} \times 2 \right) - 1$$

ii) Proportion of macroinvertebrate traits in community

The macroinvertebrate community was characterized using traits extracted from available databases (IS ARROW, 2014; Schmidt-Kloiber and Hering, 2015), supplemented by information from the literature review and from experts in different taxonomic groups. Information about fine-sediment-sensitive invertebrates was taken from Extence et al. (2013). To describe the ability to penetrate substrata, taxonomic experts assigned each taxon to one of three body flexibility modalities: (i) none, body flexibility is limited by shell or exoskeleton, e.g. molluscs, adult beetles; (ii) low, body is flexible but cannot turn around, e.g. mayflies, stoneflies; (iii) high, body can turn 360 degrees, e.g. Oligochaetes, Chironomidae (Omesová et al., 2008). Preferences for organic substrate were calculated as preferences for pelal (organic mud) and particulate organic matter. For each trait coded in a single-category assignment system or presence/absence assignment system (Schmidt-Kloiber and Hering, 2015), the community-level trait value was calculated as the number of individuals within the trait modality divided by the total macroinvertebrate abundance of a sample. The community-level trait value of traits coded in ten-point assignment system (Schmidt-Kloiber and Hering, 2015) was calculated as the sum of the trait values multiplied by the abundance of each taxon and divided by the total abundance (Schmera et al., 2014).

iii) Community structure metrics

The number of taxa, $\ln(x + 1)$ -transformed total abundance, and abundances and proportion in the community of the most abundant higher taxonomic units was calculated for each sample. This was done also for three drying-sensitive taxonomic groups (Bonada et al., 2007; Datry et al., 2014b): the family Heptageniidae, ET taxa (Ephemeroptera and Trichoptera) and EPT taxa (Ephemeroptera, Plecoptera and Trichoptera).

2.5. Index development

We performed exploratory data analysis to inform later identification of metrics that discriminate among intermittence classes. We used dot charts to assess metric distribution, and pair plots to assess their collinearity. We used nonmetric multidimensional scaling (NMDS) of a Bray-Curtis distance matrix calculated using $\ln(x + 1)$ -transformed abundance data to visualize variability in community composition in relation to year, season, subprovince and intermittence. We used the *adonis* function from the *vegan* package (Oksanen et al., 2017) to run permutational multivariate ANOVA (PERMANOVA; Anderson, 2001) to identify differences in community composition between seasons (*spring*, *autumn*), subprovinces (*Hercynian*, *West-Carpathian*) and intermittence categories (*intermittent*, *near-perennial*, *perennial*). Because only autumn samples were available for 2012, the difference between years was not tested. One-way ANOVA was used to examine differences in total abundance and taxa richness between perennial, near-perennial, and intermittent samples.

To reduce the number of potential metrics selected for inclusion in the final multimetric index (hereafter, the *Biodrought index*) a pre-selection process was conducted. To identify candidate metrics, values of each potential metric were plotted against intermittence categories. Those that effectively discriminated between intermittent and perennial samples were retained as candidate metrics (Table 1), with Mann-Whitney U tests used to identify differences in metric values between perennial and intermittent samples.

To identify the metric combination that best discriminated among perennial, near-perennial and intermittent sites, we identified linear combinations of candidate metrics using a sequence of linear

discriminant analyses (LDA). Because macroinvertebrate community composition differs among seasons (e.g. Straka et al., 2012) we expected different metrics to reflect stream intermittence in autumn and spring. Therefore, all possible combinations of 2–4 metrics were tested for their discriminatory power using the leave-one-out cross-validation procedure (*lda* function, *MASS* package; Venables and Ripley, 2002) in each season. The combination of metrics with the best discrimination power was selected and its equation noted. The final index is this equation centred to zero and can be used to calculate a ‘discrimination score’ for an independent sample. To relate the discrimination score to the intermittence categories, we calculated discrimination scores for all samples and fitted probability density functions of a normal distribution to each category, from which the probability of belonging to each category could be derived. Normality of distribution of LDA discriminant scores for each sample category was tested by Shapiro-Wilk tests. The effects of season and sample category on LDA discriminant score were tested using ANOVA. All statistical analyses and graphs were performed using R open-source software version 3.4.4 (R Core Team, 2018).

2.6. Validation by independent samples

The ability of the index to identify antecedent stream drying was tested on an external dataset, provided by the former Czech Agricultural Water Management Authority, and collected using the same method as the original dataset (Kokeš and Němejcová, 2006). The dataset comprised 117 (59 autumn + 58 spring) samples from 59 perennial sites and 26 (16 autumn + 10 spring) samples from 16 non-perennial sites. We could not distinguish between intermittent and near-perennial sites, and all samples for at which stream drying was observed were therefore classified as non-perennial. Samples were taken from 2nd to 4th order streams distributed across the Czech Republic (Fig. 2), at an altitude of 210 to 460 m a.s.l. The streams were not polluted or morphologically modified and monthly observations of hydrological instream conditions in the summer prior to sampling were available. The value of the Biodrought index was calculated for each sample and the probability of its classification as intermittent, near-perennial or perennial was calculated using the probability density function of the normal distribution. The sample was assigned to the category with the highest probability.




Fig. 2. Location of the sites used for validation within the Czech Republic. Open squares represent non-perennial sites and filled squares represent perennial sites.




Fig. 3. Total invertebrate abundance (number of individuals) and taxa richness (number of taxa) in samples classified into three flow intermittence categories: i) perennial; ii) near-perennial; iii) intermittent. The box area indicates the first and third quartiles, the central line indicates the median, whiskers represent 95% confidence intervals and circles indicates outliers.

3. Results

3.1. Community structure across years, seasons, and regions

In total, 532 552 benthic macroinvertebrate individuals from 421 taxa were collected (Appendix B). Individual sample abundance values ranged from 307 to 57 581 (mean $\pm$ SE, 4997 ± 591) individuals per sample and taxa richness values from 28 to 108 (60.7 ± 1.5) taxa per sample. Macroinvertebrate abundance decreased with increasing flow intermittence (Fig. 3), with significant differences among intermittence categories (one-way ANOVA, $F_{2,104} = 3.71$, $p = 0.028$). Taxa richness was also lowest in intermittent samples ($F_{2,104} = 15.11$, $p < 0.001$). The five most abundant taxa were *Gammarus fossarum* (30% of all individuals), *Micropsectra atrofasciata*-Gr. (9.5%), *Nemoura* sp. (3.5%), *Baetis rhodani* s.l. (3.1%) and *Habroleptoides confusa* (2.9%). Twenty-six taxa were identified as significant indicators of intermittent and near-perennial samples and 33 were identified as significant indicators of perennial samples (Appendix A). The five taxa with the highest IV were *Eiseniella tetraedra*, *Brachyptera risi*, *Parametriocnemus stylatus*, *Paraphaenocladus* sp. and *Marionina* sp. for non-perennial sites and *Dugesia*

gonocephala, *Baetis muticus*, *Baetis rhodani* s.l., *Leuctra* sp. and *Hydropsyche* sp. for perennial sites.

Non-metric multidimensional scaling analyses produced a two-dimensional solution (stress = 0.1998; Fig. 4). Community composition differed among perennial, near-perennial and intermittent samples (PERMANOVA; $F_{2,104} = 7.4$, $p < 0.001$). Despite considerable overlap, we detected differences in community composition both between seasons (PERMANOVA; $F_{1,105} = 6.0$, $p < 0.001$) and subprovinces ($F_{1,105} = 5.6$, $p < 0.001$).

3.2. Discrimination of samples from each intermittence category

Because PERMANOVA identified differences in community composition among perennial, near-perennial and intermittent samples, we investigated the metrics that most effectively discriminated between these intermittence categories. Because many potential metrics differed between intermittent and perennial samples, we plotted all calculated potential metrics and selected a subset of candidate metrics based on their ability to discriminate between categories (Table 1).




Fig. 4. Non-metric multidimensional scaling ordinations of all samples, showing differences in community composition between: (a) years (2012, 2013, 2014); (b) seasons (autumn, spring); (c) intermittence categories (intermittent, near-perennial, perennial); and (d) biogeographical subprovinces (Hercynian, West-Carpathian). Each point (sample) is connected to the group centroid.

Table 1

Candidate metrics tested as discriminators of macroinvertebrate communities from sites in different intermittence categories. Differences between perennial/intermittent samples were tested with Mann-Whitney U-tests. *** < 0.001, ** < 0.01, * < 0.05.

Metrics selected for index development	Autumn	Spring	Observed state at intermittent sites
% of lithal preference	***	***	lower
% of organic substrate preference	**	**	higher
% of rheophilic preference	***	n.s.	lower
% taxa with respiration with gills	***	***	lower
% taxa with high body flexibility	***	**	higher
% of rhithral preference	***	*	lower
% of epirhithral preference	***	*	lower
Weighted mean of sediment-sensitive invertebrates	***	n.s.	higher
Number of taxa	**	**	lower
Number of ET taxa	***	***	lower
Number of EPT taxa	***	***	lower
Total abundance	***	*	lower
Abundance of Heptageniidae	***	***	lower
Abundance of EPT	***	n.s.	lower
Proportion of Heptageniidae	***	***	lower
Proportion of Oligochaeta	***	n.s.	higher
Proportion of indicator taxa	***	***	lower

From the set of candidate metrics (Table 1), successive LDA revealed important metrics for sample separation. Three metrics were selected for discrimination in autumn: the proportion of indicator taxa (IT), the proportion of individuals with high body flexibility (HBF), and the proportion of taxa with a preference for organic substrates (POS). The formula for calculating the autumn index is:

$$\text{Biodrought index} = (4.23 \cdot \text{IT}) + (-1.69 \cdot \text{HBF}) + (-2.94 \cdot \text{POS}) - 0.26$$

Three metrics were also selected as discriminators of spring samples: the proportion of indicator taxa, the proportion of individuals with

high body flexibility, and the total abundance (TA). The formula for calculating the spring season index is:

$$\text{Biodrought index} = (4.09 \cdot \text{IT}) + (-1.76 \cdot \text{HBF}) + (0.47 \cdot \text{TA}) - 4.01$$

The Biodrought index was calculated for each sample (Fig. 5b) and the probability function for perennial, near-perennial and intermittent samples was approximated based on the distribution of index values within the intermittence categories. LDA discriminant scores (i.e. the Biodrought index) for each category were normally distributed (Shapiro-Wilk tests, $p > 0.05$). Mean discriminant scores varied between samples in the three intermittence categories ($F_{2, 101} = 215.9$; $p < 0.001$) but not between spring and autumn samples ($F_{1, 101} = 0.009$; $p = 0.926$), and there was no interaction between intermittence category and season ($F_{2, 101} = 0.328$, $p = 0.721$). Because there was no difference in the distribution of sample discriminant scores between spring and autumn, it was possible to calculate probability functions independently of season (Fig. 5a). Values for the probability function were derived from the dataset for intermittent (mean = -3.36, SD = 1.398), near-perennial (mean = -0.231, SD = 0.926) and perennial (mean = 1.559, SD = 0.777) sites.

3.3. Validation by independent samples

To evaluate the performance of the method, we calculated the Biodrought index values for a dataset not used for index development (Fig. 5c). Of 117 perennial samples, 108 (92%) had a probability > 50% of being classified correctly by the index, with eight and one samples classified as near-perennial and intermittent, respectively. The index classified 1 of 26 samples from non-perennial streams incorrectly, with 15 and 10 correctly classified as near-perennial and intermittent, respectively. Because it was not possible to distinguish between intermittent and near-perennial categories in this independent dataset, the correct classification of non-perennial samples was 96%.




Fig. 5. Biodrought index values for each flow intermittence category: (a) Probability of classification to each category according to the index; index values for each category, for (b) samples used for index development and (c) independent samples used for index validation. INT – intermittent, NPE – near-perennial, PER – perennial.

4. Discussion

Our results agree with previous research demonstrating that benthic macroinvertebrate community composition reflects stream flow intermittence (Arscott et al., 2010; Datry, 2012; Soria et al., 2017) and that the effects of relatively short (7–86 day) drying events can be detected even after seven months after flow resumption (Ledger and Hildrew, 2001; Chester and Robson, 2011). Previous studies from other regions have distinguished perennial and non-perennial streams using the proportional representation of different macroinvertebrate taxonomic groups (Mazzacano and Black, 2009; Nadeau, 2015; Cid et al., 2016; Cañedo-Argüelles et al., 2016) and using traits (Serra et al., 2017; Kelso and Entekin, 2018). Our results confirm that macroinvertebrate communities in Central Europe can also be explored to identify stream intermittence, and specifically drying events. Our Biodrought index provides a new tool to identify antecedent drying events, which may facilitate interpretation of ecological data including the results of ecological status assessments, especially when hydrological data are missing. The method was developed and validated using Czech Republic data, but it has considerable potential for wider uptake due to the extensive distribution of the continental climate zone across Central Europe.

4.1. Metrics used for index calculations

Biotic responses to environmental stress (e.g. water loss) can be species-specific (Lake, 2003) and we examined available information about responses to flow continuity mainly at the species level, and summarized the most sensitive/tolerant indicator taxa (Appendix A). The proportion of these indicator taxa was one of the most effective metrics to distinguish among intermittence categories. Flow intermittence tolerant taxa usually possess behavioural, morphological, physiological and/or life-history adaptations to survive dry phase (Lytle and Poff, 2004). For example, one species strongly associated with intermittent streams in this study was *Brachyptera risi*. This stonefly has the eggs with summer diapause, allowing the species to remain in the sediments in a viable egg stage during dry phases (Khoo, 1964). However, the selective loss of taxa sensitive to flow intermittence (and specifically drying) rather than selection for desiccation-resistant specialists is the primary driver of differences in community composition between perennial and non-perennial sites (Datry, 2012; Vidal-Abarca et al., 2013; Cid et al., 2016).

Although high body flexibility is not typically reported as a crucial trait in relation to stream drying (e.g. Bonada et al., 2007; Díaz et al., 2008; Walters, 2011) our results indicated its relevance. Organisms with flexible body shapes can respond to drying by moving into the subsurface interstices, which may maintain high humidity and can thus act as refuge for aquatic invertebrates during dry phases (Stubbington, 2012; Strachan et al., 2015). Moreover, a highly flexible body enables organisms including oligochaetes and leeches to form globular, desiccation-tolerant cysts (Montalto and Marchese, 2005; Shikov, 2011). Intermittent streams also supported a higher proportion of invertebrates with a preference for organic substrates, which may reflect their dry-phase survival in moisture-retaining patches of organic matter (Stubbington et al., 2009). Invertebrates that use such organic-rich habitats during flowing conditions can therefore persist in this substrate, which acts as a refuge that limits desiccation during dry phases (Boulton, 1989).

The Biodrought index calculated for the spring season also included total abundance. The relationship between total abundance and flow permanence in streams has previously been indicated by several studies (Rüegg and Robinson, 2004; Fenoglio et al., 2007; Datry, 2012). The lower abundance in our study was caused in particular by high mortality of *Gammarus* species, known to be sensitive to stream drying (Meyer and Meyer, 2000; Smith et al., 2003; Pařil et al., 2019).

4.2. Utility of the developed tool

Our index addresses the recognized need (Datry et al., 2011; Cid et al., 2016) for methods to identify antecedent drying events using biological metrics, and thus without using gauging stations or other prohibitively costly infrastructure. Even where gauging station data are available, biotic approaches such as our index complement the continuous, long-term hydrological information by distinguishing dry from lentic zero-flow states (Gallart et al., 2012). Sampling and processing of benthic invertebrate samples is methodologically well-managed and is routinely practised within the scope of monitoring programmes (Smith et al., 1999; Birk et al., 2012; Hill et al., 2017). The method used for taking samples (Kokeš et al., 2006) is comparable to other commonly used European methods such as AQEM/STAR method (Lorenz and Clarke, 2006), and the Biodrought index therefore has high potential for testing and use in other countries within the extensive continental climate zone that spans Central and Eastern Europe. The study dataset spans two biogeographical subprovinces (Hercynian and West-Carpathian; Culek, 2013) and the discriminatory power of the index was sufficient to correctly classify evaluated samples from the independent dataset from whole Czech Republic with high probability. Although working at a finer geographical resolution (e.g. within one subprovince) may achieve higher levels of correct assignment, we also recommend testing of our index in other Central European countries to explore the geographical limits of its reliability. A saprobic system based on common taxa is used in many Central and Eastern European countries (Rolauffs et al., 2004) and the Biodrought index may therefore be useable in Central European region without extensive taxonomic adjustment.

Our index was developed in pristine streams where only a single, natural stressor (drying) was known to influence macroinvertebrate community composition. Elsewhere, factors that may affect index performance include interactions between the hydrological stressor of drying (both natural and anthropogenic) with other human pressures, interactions which can have antagonistic or additive effects on biological communities (Folt et al., 1999; Matthaei et al., 2010; Ormerod et al., 2010). In particular, the typical sequence of hydrological conditions from low flows, to flow cessation and gradual drying can reduce oxygen concentrations, and intermittence and organic pollution can therefore have comparable effects by eliminating sensitive invertebrates (Pardo and García, 2016). Moreover, a reduction in oxygen availability and low flow can have synergistic effects on benthic invertebrates (Calapez et al., 2017). Disentangling the effects of stream drying and anthropogenic stressors (including organic pollution) is therefore challenging, but consideration of the Biodrought index alongside physico-chemical quality elements may improve the accuracy of ecological status assessments that might otherwise incorrectly classify IRES as polluted. Wider testing is needed to explore how the index performs in streams exposed to human stressors.

The spatial distribution of intermittent reaches in relation to perennial reaches and other refuges may have a considerable influence on the rate and extent of community recovery after drying events (Sedell et al., 1990; Bogan et al., 2017; but see Datry et al., 2014b). The spatial proximity of extensive refuges, in particular upstream perennial reaches, may enable rapid recolonization and community reestablishment within weeks to obscure the occurrence of an antecedent dry phase (Fowler, 2004; Fritz and Dodds, 2004; Pařil et al., 2019). In our dataset, the intermittent sites were at least 0.5 km downstream of a perennial reach and the samples were taken at most 7 months after flow resumed. However, recovery at highly isolated sites (e.g. > 10 km from perennial refuges) may take multiple years with repeated drying events often preventing aquatic communities from reaching a stable state (Bogan and Lytle, 2011; Bogan et al., 2017). Our index and comparable tools thus have higher potential for strong performance in IRES with relatively short dry phases, compared to those which may remain dry for years.

4.3. Conclusions and recommendations for index use

We developed the Biodrought index: a novel index to characterize the effect of stream drying on aquatic invertebrate communities. The index enable calculation of the probability of the antecedent stream drying, based on benthic invertebrates sampled using standard methods. We demonstrated the index as robust at national scale. The robustness of our index is evidenced by its registration by the Ministry of Environment of the Czech Republic, and we recommend its testing and adoption (with required adaptation) in other European countries within continental-climate region. In particular, the Biodrought index is intended for use by those with responsibility for monitoring and management of river ecosystems, including water management boards, environmental agencies and private water companies. We recommend adoption of the Biodrought index by such managers, to improve their interpretation of the macroinvertebrate assemblage data collected during ecological status assessments. The index can help avoid misinterpreting a deviation of community composition from reference status caused by a preceding dry phase (Reyjol et al., 2014; Cid et al., 2016; Stubbington et al., 2018). Environmental agencies could employ this tool for characterization of dry phase effects on macroinvertebrate communities in protected areas or for evaluation of the effectiveness of projects intended to restore naturally perennial flow at sites impacted by water resource pressures such as over-abstraction. The index can also be useful when identifying the impacts of water resource use in naturally perennial streams. However, since the method was developed in naturally intermittent streams, its use in non-natural IRES must be carefully assessed and the performance of Biodrought index in such conditions remains to be tested.

Our approach was essentially qualitative, identifying if a drying event had or had not occurred in the the one-year period preceding sample collection. Further research is needed to explore variability within 'intermittent', 'near-perennial' and other flow intermittence categories, in particular to identify species-specific responses to dry phases of differing durations. Identification of such thresholds is crucial to predict biotic responses to increasing intermittence in the context of ongoing climate change, which is interacting with increasing water resource pressures.

Acknowledgements

This work was supported by Long-term development conception of Výzkumný Ústav Vodohospodářský TGM, p. r. i. no. IG/2019/1612 and by COST Action CA15113 (Science and Management of Intermittent Rivers and Ephemeral Streams (www.smires.eu) and INTER-COST grant no. LTC17017. M. Polášek, P. Pařil and T. Datry were supported by the joined French Barrande 489 Project no. 38114TL and Czech MOBILITY project no. 7AMB17FR011, both entitled 490 DRYGRAD. We thank Michal Pospíšil for creating of the graphical abstract, Ondra Hájek for creating the maps and reviewers for their helpful comments.

Declaration of Competing Interest

None.

Appendices A and B. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ecolind.2019.105486>.

References

Acuña, V., Datry, T., Marshall, J., Barceló, D., Dahm, C.N., Ginebreda, A., McGregor, G., Sabater, S., Tockner, K., Palmer, M.A., 2014. Why should we care about temporary waterways? *Science* 343, 1080–1081. <https://doi.org/10.1126/science.1246666>.
Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of

variance. *Austral Ecol.* 26, 32–46. <https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x>.
Arscott, D.B., Larned, S., Scarsbrook, M.R., Lambert, P., 2010. Aquatic invertebrate community structure along an intermittence gradient: Selwyn River, New Zealand. *J. N. Am. Benthol. Soc.* 29, 530–545. <https://doi.org/10.1899/08-124.1>.
Beaufort, A., Lamouroux, N., Pella, H., Datry, T., Sauquet, E., 2018. Extrapolating regional probability of drying of headwater streams using discrete observations and gauging networks. *Hydrol. Earth Syst. Sci.* 22, 3033–3051. <https://www.hydrol-earth-syst-sci.net/22/3033/2018/>.
Birk, S., Bonne, W., Borja, A., Brucet, S., Courrat, A., Poikane, S., Solimini, A., Van De Bund, W., Zampoukas, N., Hering, D., 2012. Three hundred ways to assess Europe's surface waters: an almost complete overview of biological methods to implement the Water Framework Directive. *Ecol. Ind.* 18, 31–41. <https://doi.org/10.1016/j.ecolind.2011.10.009>.
Bogan, M.T., Lytle, D.A., 2011. Severe drought drives novel community trajectories in desert stream pools. *Freshw. Biol.* 56, 2070–2081. <https://doi.org/10.1111/j.1365-2427.2011.02638.x>.
Bogan, M.T., Boersma, K.S., Lytle, D.A., 2013. Flow intermittency alters longitudinal patterns of invertebrate diversity and assemblage composition in an arid-land stream network. *Freshw. Biol.* 58, 1016–1028. <https://doi.org/10.1111/fwb.12105>.
Bogan, M.T., Chester, E.T., Datry, T., Murphy, A.L., Robson, B.J., Ruhi, A., Stubbington, R., Whitney, J.E., 2017. Resistance, resilience, and community recovery in intermittent rivers and ephemeral streams. In: Datry, T., Bonada, N., Boulton, A. (Eds.), *Intermittent Rivers and Ephemeral Streams. Ecology and Management*. Elsevier, Netherlands, pp. 349–376.
Bonada, N., Rieradevall, M., Prat, N., 2007. Macroinvertebrate community structure and biological traits related to flow permanence in a Mediterranean river network. *Hydrobiologia* 589, 91–106. <https://doi.org/10.1007/s10750-007-0723-5>.
Boulton, A.J., 1989. Over-summer refuges of aquatic macroinvertebrates in two intermittent streams in central Victoria. *Trans. R. Soc. South Australia* 113, 22–34.
Calapez, A.R., Branco, P., Santos, J.M., Ferreira, T., Hein, T., Brito, A.G., Feio, M.J., 2017. Macroinvertebrate short-term responses to flow variation and oxygen depletion: a mesocosm approach. *Sci. Total Environ.* 599–600, 1202–1212. <https://doi.org/10.1016/j.scitotenv.2017.05.056>.
Cañedo-Argüelles, M., Bogan, M.T., Lytle, D.A., Prat, N., 2016. Are Chironomidae (Diptera) good indicators of water scarcity? Dryland streams as a case study. *Ecol. Ind.* 71, 155–162. <https://doi.org/10.1016/j.ecolind.2016.07.002>.
Chadd, R.P., England, J.A., Constable, D., Dunbar, M.J., Extence, C.A., Leeming, D.J., Murray-Bligh, J.A., Wood, P.J., 2017. An index to track the ecological effects of drought development and recovery on riverine invertebrate communities. *Ecol. Ind.* 82, 344–356. <https://doi.org/10.1016/j.ecolind.2017.06.058>.
Chester, E.T., Robson, B.J., 2011. Drought refuges, spatial scale and recolonisation by invertebrates in non-perennial streams. *Freshw. Biol.* 56, 2094–2104. <https://doi.org/10.1111/j.1365-2427.2011.02644.x>.
Cid, N., Verkaik, I., García-Roger, E.M., Rieradevall, M., Bonada, N., Sánchez-Montoya, M.M., Gómez, R., Suárez, M.L., Vidal-Abarca, M.R., Demartini, D., Buffagni, A., Erba, S., Karaouzas, I., Skoulidakis, N., Prat, N., 2016. A biological tool to assess flow connectivity in reference temporary streams from the Mediterranean Basin. *Sci. Total Environ.* 540, 178–190. <https://doi.org/10.1016/j.scitotenv.2015.06.086>.
Culek, M., 2013. Biogeographical provinces, subprovinces and bioregions of the Czech Republic. *J. Landscape Ecol.* 6, 5–16. <https://doi.org/10.2478/s10285-012-0065-5>.
Datry, T., 2012. Benthic and hyporheic invertebrate assemblages along a flow intermittence gradient: effects of duration of dry events. *Freshw. Biol.* 57, 563–574. <https://doi.org/10.1111/j.1365-2427.2011.02725.x>.
Datry, T., Arscott, D.B., Sabater, S., 2011. Recent perspectives on temporary river ecology. *Aquat. Sci.* 73, 453–457. <https://doi.org/10.1007/s00027-011-0236-1>.
Datry, T., Larned, S.T., Tockner, K., 2014a. Intermittent rivers: a challenge for freshwater ecology. *Bioscience* 64, 229–235. <https://doi.org/10.1093/biosci/bit027>.
Datry, T., Larned, S.T., Fritz, K.M., Bogan, M.T., Wood, P.J., Meyer, E.I., Santos, A.N., 2014b. Broad-scale patterns of invertebrate richness and community composition in temporary rivers: effects of flow intermittence. *Ecography* 37, 69–104. <https://doi.org/10.1111/j.1600-0587.2013.00287.x>.
Datry, T., Boulton, A.J., Bonada, N., Fritz, K., Leigh, C., Sauquet, E., Tockner, K., Huguén, B., Dahm, C.N., 2017. Flow intermittence and ecosystem services in rivers of the Anthropocene. *J. Appl. Ecol.* 55, 353–364. <https://doi.org/10.1111/1365-2664.12941>.
De Cáceres, M., Legendre, P., 2009. Associations between species and groups of sites: indices and statistical inference. *Ecology* 90, 3566–3574. <https://doi.org/10.1890/08-1823.1>.
Díaz, A.M., Alonso, M.L.S., Gutiérrez, M.R.V.-A., 2008. Biological traits of stream macroinvertebrates from a semi-arid catchment: patterns along complex environmental gradients. *Freshw. Biol.* 53, 1–21. <https://doi.org/10.1111/j.1365-2427.2007.01854.x>.
Dufrene, M., Legendre, P., 1997. Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecol. Monogr.* 67, 345–366. <https://doi.org/10.2307/2963459>.
European Commission, 2000. Directive 2000/60/EC of the European Parliament of the Council of 23rd October 2000 establishing a framework for community action in the field of water policy. *Official J. Eur. Commun.* 327, 1–72.
Extence, C.A., Chadd, R.P., England, J., Dunbar, M.J., Wood, P.J., Taylor, E.D., 2013. The assessment of fine sediment accumulation in rivers using macro-invertebrate community response. *River Res. Appl.* 29, 17–55. <https://doi.org/10.1002/rra.1569>.
Fenoglio, S., Bo, T., Cucco, M., Malacarne, G., 2007. Response of benthic invertebrate assemblages to varying drought conditions in the Po river (NW Italy). *Italian J. Zool.* 74, 191–201. <https://doi.org/10.1080/11250000701286696>.
Folt, C.L., Chen, C.Y., Moore, M.V., Burnaford, J., 1999. Synergism and antagonism

- among multiple stressors. *Limnol. Oceanogr.* 44, 864–877. <https://doi.org/10.4319/lo.1999.44.3.part.2.0864>.
- Fowler, R.T., 2004. The recovery of benthic invertebrate communities following dewatering in two braided rivers. *Hydrobiologia* 523, 17–28. <https://doi.org/10.1023/B:HYDR.0000033077.13139.7f>.
- Fritz, K.M., Dodds, W.K., 2004. Resistance and resilience of macroinvertebrate assemblages to drying and flood in a tallgrass prairie stream system. *Hydrobiologia* 527, 99–112. <https://doi.org/10.1023/B:HYDR.0000043188.53497.9b>.
- Gallart, F., Prat, N., García-Roger, E.M., Latron, J., Rieradevall, M., Llorens, P., Barberá, G.G., Brito, D., De Girolamo, A.M., Lo Porto, A., Buffagni, A., Erba, S., Neves, R., Nikolaidis, N.P., Perrin, J.L., Querner, E.P., Quiñero, J.M., Tournoud, M.G., Tzoraki, O., Skoulikidis, N., Gómez, R., Sánchez-Montoya, M.M., Froebrich, J., 2012. A novel approach to analysing the regimes of temporary streams in relation to their controls on the composition and structure of aquatic biota. *Hydrol. Earth Syst. Sci.* 16, 3165–3182. <https://doi.org/10.5194/hess-16-3165-2012>.
- Gallart, F., Llorens, P., Latron, J., Cid, N., Rieradevall, M., Prat, N., 2016. Validating alternative methodologies to estimate the regime of temporary rivers when flow data are unavailable. *Sci. Total Environ.* 565, 1001–1010. <https://doi.org/10.1016/j.scitotenv.2016.05.116>.
- Hill, R.A., Fox, E.W., Leibowitz, S.G., Olsen, A.R., Thornbrugh, D.J., Weber, M.H., 2017. Predictive mapping of the biotic condition of conterminous U.S. rivers and streams. *Ecol. Appl.* 27, 2397–2415. <https://doi.org/10.1002/eap.1617>.
- Hille, S., Kristensen, E.A., Graeber, D., Riis, T., Jørgensen, N.K., Baattrup-Pedersen, A., 2014. Fast reaction of macroinvertebrate communities to stagnation and drought in streams with contrasting nutrient availability. *Freshw. Sci.* 33, 847–859. <https://doi.org/10.1086/677554>.
- Illies, J., 1967. *Limnofauna Europaea. Eine Zusammenstellung aller die europäischen Binnengewässer bewohnenden mehrzelligen Tierarten m. Angaben über d. Verbreitung u. Ökologie*, first ed. Gustav Fischer Verlag, Stuttgart, pp. 474.
- IS ARROW (Assessment and Reference Reports of Water Monitoring), 2014. Praha, Český hydrometeorologický ústav. <http://hydro.chmi.cz/isarrow/> (accessed 20.06.2014).
- Kelso, J.E., Entekhabi, S.A., 2018. Intermittent and perennial macroinvertebrate communities had similar richness but differed in species trait composition depending on flow duration. *Hydrobiologia* 807, 189–206. <https://doi.org/10.1007/s10750-017-3393-y>.
- Khoo, S.G., 1964. *Studies on the Biology of Stoneflies*. Unpublished Ph.D. thesis. University of Liverpool.
- Kokeš, J., Němejcová, D., 2006. Metodika odběru a zpracování vzorků makrozoobentosu tekoucích vod metodou PERLA [Methodology for macrozoobenthos sampling in running waters by PERLA method]. Metodika VÚV TGM Praha.
- Kokeš, J., Zahradková, S., Němejcová, D., Hodovský, J., Jarkovský, J., Soldán, T., 2006. The PERLA system in the Czech Republic: a multivariate approach to assess ecological status of running waters. *Hydrobiologia* 566, 343–354. https://doi.org/10.1007/978-1-4020-5493-8_24.
- Lake, P.S., 2003. Ecological effects of perturbation by drought in flowing waters. *Freshw. Biol.* 48, 1161–1172. <https://doi.org/10.1046/j.1365-2427.2003.01086.x>.
- Lancaster, J., Ledger, M.E., 2015. Population-level responses of stream macroinvertebrates to drying can be density-independent or density-dependent. *Freshw. Biol.* 60, 2559–2570. <https://doi.org/10.1111/fwb.12643>.
- Ledger, M.E., Hildrew, A.G., 2001. Recolonization by the benthos of an acid stream following a drought. *Arch. Hydrobiol.* 152, 1–17. <https://doi.org/10.1127/archiv-hydrobiol/152/2001/1>.
- Leigh, C., Bonada, N., Boulton, A.J., Huguely, B., Larned, S.T., Vander Vorste, R., Datry, T., 2016. Invertebrate assemblage responses and the dual roles of resistance and resilience to drying in intermittent rivers. *Aquat. Sci.* 78, 291–301. <https://doi.org/10.1007/s00027-015-0427-2>.
- Leigh, C., Datry, T., 2017. Drying as a primary hydrological determinant of biodiversity in river systems: a broad-scale analysis. *Ecography* 40, 487–499. <https://doi.org/10.1111/ecog.02230>.
- Li, L., Zheng, B., Liu, L., 2010. Biomonitoring and bioindicators used for river ecosystems: definitions, approaches and trends. *Procedia Environ. Sci.* 2, 1510–1524. <https://doi.org/10.1016/j.proenv.2010.10.164>.
- Lorenz, A., Clarke, R.T., 2006. Sample coherence – a field study approach to assess similarity of macroinvertebrate samples. *Hydrobiologia* 566, 461–476. https://doi.org/10.1007/978-1-4020-5493-8_32.
- Lytle, D.A., Poff, N.L., 2004. Adaptation to natural flow regimes. *Trends Ecol. Evol.* 19, 94–100. <https://doi.org/10.1016/j.tree.2003.10.002>.
- Marshall, J.C., Acuña, V., Allen, D.C., Bonada, N., Boulton, A.J., Carlson, S.M., Dahm, C.N., Datry, T., Leigh, C., Negus, P., Richardson, J.S., 2018. Protecting U.S. temporary waterways. *Science* 361, 856–857. <https://doi.org/10.1126/science.aav0839>.
- Matthaei, C.D., Piggott, J.J., Townsend, C.R., 2010. Multiple stressors in agricultural streams: interactions among sediment addition, nutrient enrichment and water abstraction. *J. Appl. Ecol.* 47, 639–649. <https://doi.org/10.1111/j.1365-2664.2010.01809.x>.
- Mazzacano, C., Black, S.H., 2009. *Using Aquatic Macroinvertebrates as Indicators of Stream Flow Duration*. The Xerces Society for Invertebrate Conservation, Portland, USA.
- Menció, A., Mas-Pla, J., 2010. Influence of groundwater exploitation on the ecological status of streams in a Mediterranean system (Selva Basin, NE Spain). *Ecol. Ind.* 10, 915–926. <https://doi.org/10.1016/j.ecolind.2010.02.001>.
- Meyer, A., Meyer, E.L., 2000. Discharge regime and the effect of drying on macroinvertebrate communities in a temporary karst stream in East Westphalia (Germany). *Aquat. Sci.* 62, 216–231. <https://doi.org/10.1007/PL00001333>.
- Montalvo, L., Marchese, M., 2005. Cyst formation in Tubificidae (Naidinae) and Opisthocystidae (Annelida, Oligochaeta) as an adaptive strategy for drought tolerance in fluvial wetlands of the Paraná River, Argentina. *Wetlands* 25, 488–494. [https://doi.org/10.1672/0097-0706\(2005\)025\[0488\]2.0.CO;2](https://doi.org/10.1672/0097-0706(2005)025[0488]2.0.CO;2).
- Munné, A., Prat, N., 2009. Use of macroinvertebrate-based multimetric indices for water quality evaluation in Spanish Mediterranean rivers: an intercalibration approach with IBMWP index. *Hydrobiologia* 628, 203–225. <https://doi.org/10.1007/s10750-009-9757-1>.
- NC Division of Water Quality, 2010. Methodology for Identification of Intermittent and Perennial Streams and their Origins, Version 4.11. North Carolina Department of Environment and Natural Resources, Division of Water Quality. Raleigh, NC.
- Nadeau, T.-L., 2015. *Streamflow Duration Assessment Method for the Pacific Northwest*. US Environmental Protection Agency, region 10, Seattle, WA.
- Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2017. *vegan: Community Ecology Package*. R package version 2.4-3. <https://CRAN.R-project.org/package=vegan>.
- Omesová, M., Horsák, M., Helešic, J., 2008. Nested patterns in hyporheic meta-communities: the role of body morphology and penetrability of sediment. *Naturwissenschaften* 95, 917–926. <https://doi.org/10.1007/s00114-008-0399-3>.
- Ormerod, S.J., Dobson, M., Hildrew, A.G., Townsend, C., 2010. Multiple stressors in freshwater ecosystems. *Freshw. Biol.* 55, 1–4. <https://doi.org/10.1111/j.1365-2427.2009.02395.x>.
- Pardo, I., García, I., 2016. Water abstraction in small lowland streams: unforeseen hypoxia and anoxia effects. *Sci. Total Environ.* 568, 226–235. <https://doi.org/10.1016/j.scitotenv.2016.05.218>.
- Pařil, P., Leigh, C., Polášek, M., Sarremejane, R., Řezníčková, P., Dostálová, A., Stubbington, R., 2019. Short-term streambed drying events alter amphipod population structure in a central European stream. *Fundamental Appl. Limnol.* <https://doi.org/10.1127/fal/2019/1164>.
- Peel, M.C., Finlayson, B.L., McMahon, T.A., 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrol. Earth Syst. Sci.* 11, 1633–1644. <https://doi.org/10.5194/hess-11-1633-2007>.
- Poff, N.L., 1997. Landscape filters and species traits: towards mechanistic understanding and prediction in stream ecology. *J. N. Am. Benthol. Soc.* 16, 391–409. <https://doi.org/10.2307/1468026>.
- R Core Team, 2018. R: A language and environment for statistical computing. Retrieved from <http://www.R-project.org/>.
- Reyrol, Y., Argillier, Ch., Bonne, W., Borja, A., Buijse, A.D., Cardoso, A.C., Daufresne, M., Kernan, M., Ferreira, M.T., Poikane, S., Prat, N., Solheim, A.-L., Stroffek, S., Usseglio-Polatera, P., Villeneuve, B., van de Bund, W., 2014. Assessing the ecological status on the context of the European Water Framework Directive: where do we go now? *Sci. Total Environ.* 497–498, 332–344. <https://doi.org/10.1016/j.scitotenv.2014.07.119>.
- Rolauffs, P., Stubaer, I., Zahradková, S., Brabec, K., Moog, O., 2004. Integration of the saprobic system into the European Union Water Framework Directive. *Hydrobiologia* 516, 285–298.
- Rosenberg, D.M., Resh, V.H. (Eds.), 1993. *Freshwater Biomonitoring and Benthic Macroinvertebrates*. Chapman and Hall, New York.
- Rüegg, J., Robinson, Ch.T., 2004. Comparison of macroinvertebrate assemblages of permanent and temporary streams in an Alpine flood plain, Switzerland. *Arch. Hydrobiol.* 161, 489–510. <https://doi.org/10.1127/0003-9136/2004/0161-0489>.
- Schmera, D., Podani, J., Erős, T., Heino, J., 2014. Combining taxon-by-trait and taxon-by-site matrices for analysing trait patterns of macroinvertebrate communities: a rejoinder to Monaghan & Soares (2014). *Freshw. Biol.* 59, 1551–1557. <https://doi.org/10.1111/fwb.12369>.
- Schmidt-Kloiber, A., Hering, D., 2015. www.freshwaterecology.info – an online tool that unifies, standardises and codifies more than 20,000 European freshwater organisms and their ecological preferences. *Ecol. Ind.* 53, 271–282. <https://doi.org/10.1016/j.ecolind.2015.02.007>.
- Schriever, T.A., Bogan, M.T., Boersma, K.S., Cañedo-Argüelles, M., Jaeger, K.L., Olden, J.D., Lytle, D.A., 2015. Hydrology shapes taxonomic and functional structure of desert stream invertebrate communities. *Freshw. Sci.* 34, 399–409. <https://doi.org/10.1086/680518>.
- Sedell, J.R., Reeves, G.H., Hauer, F.R., Stanford, J.A., Hawkins, Ch.P., 1990. Role of refugia in recovery from disturbances: modern fragmented and disconnected river systems. *Environ. Manage.* 14, 711–724. <https://doi.org/10.1007/BF02394720>.
- Serra, S.R.Q., Graça, M.A.S., Dolédec, S., Feio, M.J., 2017. Discriminating permanent from temporary rivers with traits of chironomid genera. *Ann. Limnol.* – Int. J. Limnol. 53, 161–174. <https://doi.org/10.1051/limn/2017004>.
- Shikov, E.V., 2011. *Haemopsis sanguisuga* (Linnaeus, 1758) (Hirudinea) – the first observation of a leech predation on terrestrial gastropods. *Folia Malacol.* 19, 103–106. <https://doi.org/10.2478/v10125-011-0016-5>.
- Smith, M.J., Kay, W.R., Edward, D.H.D., Papas, P.J., Richardson, K. StJ, Simpson, J.C., Pinder, A.M., Cale, D.J., Horwitz, P.H.J., Davis, J.A., Yung, F.H., Norris, R.H., Halse, S.A., 1999. AusRivAS: using macroinvertebrates to assess ecological condition of rivers in Western Australia. *Freshw. Biol.* 41, 269–282. <https://doi.org/10.1046/j.1365-2427.1999.00430.x>.
- Smith, H., Wood, P.J., Gunn, J., 2003. The influence of habitat structure and flow permanence on invertebrate communities in karst spring systems. *Hydrobiologia* 510, 53–66. <https://doi.org/10.1023/B:HYDR.0000008501.55798.20>.
- Soria, M., Leigh, C., Datry, T., Bini, L.M., Bonada, N., 2017. Biodiversity in perennial and intermittent rivers: a meta-analysis. *Oikos* 126, 1078–1089. <https://doi.org/10.1111/oik.04118>.
- Strachan, S.R., Chester, E.T., Robson, B.J., 2015. Freshwater invertebrate life history strategies for surviving desiccation. *Springer Sci. Rev.* 3, 57–75. <https://doi.org/10.1007/s40362-015-0031-9>.
- Straka, M., Syrovátková, V., Helešic, J., 2012. Temporal and spatial macroinvertebrate variance compared: crucial role of CPOM in a headwater stream. *Hydrobiologia* 686, 119–134. <https://doi.org/10.1007/s10750-012-1003-6>.

- Stubbington, R., 2012. The hyporheic zone as an invertebrate refuge: a review of variability in space, time, taxa and behaviour. *Freshw. Res.* 63, 293–311. <https://doi.org/10.1071/MF11196>.
- Stubbington, R., Wood, P.J., Boulton, A.J., 2009. Low flow controls on benthic and hyporheic macroinvertebrate assemblages during supra-seasonal drought. *Hydrol. Process.* 23, 2252–2263. <https://doi.org/10.1002/hyp.7290>.
- Stubbington, R., England, J., Wood, P.J., Sefton, C.E.M., 2017. Temporary streams in temperate zones: recognizing, monitoring and restoring transitional aquatic-terrestrial ecosystems. *Wiley Interdisciplinary Rev., Water* 4 <https://doi.org/10.1002/wat2.1223>. Article e1223.
- Stubbington, R., Chadd, R., Cid, N., Csabai, Z., Miliša, M., Morais, M., Munné, A., Pařil, P., Peřić, V., Tziortzis, I., Verdonschot, R.C.M., Detry, T., 2018. Biomonitoring of intermittent rivers and ephemeral streams in Europe: current practice and priorities to enhance ecological status assessments. *Sci. Total Environ.* 618, 1096–1113. <https://doi.org/10.1016/j.scitotenv.2017.09.137>.
- Venables, W.N., Ripley, B.D., 2002. *Modern Applied Statistics with S*, fourth ed. Springer, New York.
- Vidal-Abarca, M.R., Sánchez-Montoya, M.M., Guerrero, C., Gómez, R., Arce, M.I., García-García, V., Suárez, M.L., 2013. Effects of intermittent stream flow on macroinvertebrate community composition and biological traits in a naturally saline Mediterranean stream. *J. Arid Environ.* 99, 28–40. <https://doi.org/10.1016/j.jaridenv.2013.09.008>.
- Walters, A.W., 2011. Resistance of aquatic insects to a low-flow disturbance: exploring a trait-based approach. *Freshw. Sci.* 30, 346–357. <https://doi.org/10.1899/10-041.1>.
- Wilby, R.L., Orr, H., Watts, G., Battarbee, R.W., Berry, P.M., Chadd, R., Dugdale, S.J., Dunbar, M.J., Elliott, J.A., Extence, C., Hannah, D.M., Holmes, N., Johnson, A.C., Knights, B., Milner, N.J., Ormerod, S.J., Solomon, D., Timlett, R., Whitehead, P.J., Wood, P.J., 2010. Evidence needed to manage freshwater ecosystems in a changing climate: turning adaptation principles into practice. *Sci. Total Environ.* 408, 4150–4164. <https://doi.org/10.1016/j.scitotenv.2010.05.014>.
- Wilding, N.A., White, J.C., Chadd, R.P., House, A., Wood, P.J., 2018. The influence of flow permanence and drying pattern on macroinvertebrate biomonitoring tools used in the assessment of riverine ecosystems. *Ecol. Ind.* 85, 548–555. <https://doi.org/10.1016/j.ecolind.2017.10.059>.
- Zahrádková, S., Hájek, O., Tremel, P., Pařil, P., Straka, M., Němejcová, D., Polášek, M., Ondráček, P., 2015. Hodnocení rizika vysychání drobných vodních toků v České republice. *VTEI* 6, 4–16.